



The Ethanolic Extract of *Trichosanthes dioica* Leaves Can Ameliorate the Liver Damage in Paracetamol Induced Liver Toxicity in SD Rats

**Md. Masum Billaha¹, Md. Abir Hasan¹, Uma Rani Sarker¹, Ratri Banik¹,
Md. Imran Hossen¹ and Md. Nahid Hasan^{1,2*}**

¹Department of Pharmacy, Gono Bishwabidyalay, Savar, Dhaka-1344, Bangladesh.

²Department of Pharmaceutical Sciences, North South University, Dhaka-1229, Bangladesh.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2018/46204

Editor(s):

(1) Dr. Barkat Ali Khan, Department of Pharmaceutics, Gomal University, Dera Ismail Khan, Pakistan.

Reviewers:

(1) M. Gunjegaonkar Shivshankar, JSPM's Charak College of Pharmacy and Research, India.

(2) S. Murugesan, Pachaiyappa's College, University of Madras, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46204>

Original Research Article

**Received 06 October 2018
Accepted 24 December 2018
Published 23 January 2019**

ABSTRACT

Traditional plants have been used to treat hepatotoxicity by folk medicine practitioners. *Trichosanthes dioica* (TD) has been used in folk medicine to alleviate several diseases. In the present study, the ethanolic extract of *Trichosanthes dioica* leaves has been utilized to study its activity on paracetamol induced hepatotoxicity in rats.

Swiss albino mice (25-30 g) and SD Rats (140–200 g) were used. Acute hepatotoxicity was induced by paracetamol (800 mg/kg body weight), administered once daily for one week whereas the test extract was given orally throughout the whole experiment at 250 and 500 mg/kg body weight. Silymarin (100 mg/kg b.w.) was given orally as standard hepatoprotective drug. The degree of hepatoprotection was determined by the estimation of biochemical parameters like ALT, AST, ALP, bilirubin, total protein and albumin level.

The increased plasma levels of hepatic marker enzymes including AST, ALT, ALP and bilirubin found in the paracetamol control group, which may be due to the liver cell destruction or changes in the cell membrane permeability indicating severity of hepatocellular damage induced by

*Corresponding author: E-mail: nahidhasan.nsu2018@gmail.com;

paracetamol. Pre-treatment with TD as well as standard hepatoprotective agent silymarin recovered the increased plasma levels of these hepatic enzymes to the reduced levels. The results of the study provide evidence that the *Trichosanthes dioica* leaves has shown hepatoprotective activity against paracetamol induced hepatotoxicity in rats.

Keywords: *Trichosanthes dioica*; hepatoprotective activity; paracetamol induced hepatotoxicity; liver enzymes.

1. INTRODUCTION

The liver is “Great Chemical Factory” of body as involved in regulation, synthesis, storage and secretion of many important proteins, nutrients and chemicals. Liver is also involved in detoxification of toxic chemicals and xenobiotics. It is exposed to various toxins, changing liver functions and eventually leading to liver ailments like hepatitis, cirrhosis and alcoholic liver disease [1]. During many biochemical processes, free radical reactive oxygen species are generated that cause liver injury [2].

A large number of drugs have been reported to have potentially hepatotoxicity [3]. Paracetamol induces hepatotoxicity which results with several cases of cirrhosis, hepatitis and suicidal attempts [4]. An overdose of the analgesic drug paracetamol can lead to severe liver injury in humans and in experimental animals [5].

Herbal drugs have gained importance and popularity in recent years because of their safety, efficacy and cost effectiveness. One of the important and well -documented uses of plant products is their use as hepatoprotective agents. Hence, there is an ever increasing need for safe hepatoprotective agent [6]. The biodiversity of flora of Bangladesh is very broad, and several native Bangladeshi medicinal plant species have a long tradition of use with great phytotherapeutic potential [7]. So, research in medicinal plants is a vital sector for the discovery of promising drugs in Bangladesh [8].

Trichosanthes dioica Roxb. L. (family: Cucurbitaceae) is an annual or perennial herb. It is called pointed gourd in English, Potol in Bengali and Patola in Sanskrit, is a dioecious climber distributed in tropical Asia, Polynesia, and Australia [9,10,11]. In an earlier chemical study revealed the presence of the toxic bitter principles cucurbitacins. The main phytochemical groups that were present were alkaloids, glycosides, flavonoids, carbohydrates, fixed oils, steroids, saponins, tannins, and phenols. The various chemical constituents present in *T. dioica*

are vitamin A, vitamin C, tannins, and saponin [12,13,14,15]. The various parts of TD have been used in several diseases like as tonic, febrifuge, in alcoholism, jaundice, edema, alopecia; also as antipyretic, diuretic, cardiotoxic, hypoglycemic and laxative [16,17,18].

The purpose of this work was to explore the possible hepatoprotective activity of *T. dioica* leaves extract in experimental animals.

2. MATERIALS AND METHODS

2.1 Plant Collection and Extraction

The plant *Trichosanthes dioica* (TD) leaves were collected from Savar area and was identified and authenticated by the Dept. of Botany, Jahangirnagar University, Savar, Dhaka. The collected materials were thoroughly washed in water, cut into smaller parts and shed dried at 35°–40°C and pulverized in electric grinder to get extractable powder. Then powder was extracted in soxhlet apparatus with ethanol (96%).

2.2 Experimental Animals

For the experiment, Sprague Dawley rats of either sex, weighing between 140-200 g, were collected from the animal research lab in the Department of Pharmacy, Jahangirnagar University, Savar, Dhaka. Animals were maintained under standard environmental conditions (temperature: 27.0±1.0°C, relative humidity: 55-65% and 12 h light/12 h dark cycle) and had free access to feed and water *ad libitum*. The animals were acclimatized to laboratory condition for one week prior to experiments. All protocols for animal experiment were approved by the institutional animal ethical committee.

2.3 Toxicity Studies

Toxicity studies of the extracts were carried out in Swiss Albino mice of either sex weighing between 20 and 25 g. The extract was found to be safe till 4000 mg/kg p.o. [19].

2.4 Experimental Design for the Assessment of Liver Functions

Animal study was performed at Pharmacology Laboratory, Department of Pharmacy, Jahangirnagar University, Savar, Dhaka-1342. The rats were housed in polypropylene cages at room temperature (27±2°C). The rats were divided into five groups of 6 animals (n = 6) each.

Group I: Received water (10 mL/kg p.o.) once daily for 7 days, and served as normal control.

Group II: Received water (10 mL/kg p.o.) and paracetamol 800 mg/kg, p.o. once daily for 7 days and served as positive control.

Group III: Received standard drug silymarin (100 mg/ kg p.o.) and paracetamol 800 mg/kg, p.o. once daily for 7 days, serving as STD.

Group IV and V: Received *Trichosanthes dioica* (TD) leaves extract (250 and 500 mg/kg respectively) and paracetamol 800 mg/kg p.o. once daily for 7 days [20].

Rats were anesthetized using ketamine. After sacrifice, blood samples were collected and the serums were separated by centrifugation. Serum samples were subjected to liver function tests of enzymes such as glutamate pyruvate transaminase (GPT/ALT), glutamate-oxaloacetate transaminase (GOT/AST), alkaline phosphatase (ALP), total bilirubin, total protein, albumin and globulin by using the standard enzymatic colorimetric method.

2.5 Statistical Analysis

Statistical analysis of animal experiments was carried out using One way ANOVA following Dunnet's post hoc test using SPSS 16.0 for windows. Data were presented as Mean ± SEM. The results obtained were compared with the vehicle control group. *p* values <0.05, <0.01 and <0.001 were considered to be statistically significant, highly significant and very highly significant respectively.

3. RESULTS AND DISCUSSION

Acetaminophen (AAP) is a frequently used analgesic that can cause hepatic necrosis in high doses by the formation of NAPQI which causes the hepatic GSH depletion. At high doses, more NAPQI will bind covalently to

cellular macromolecules [21,22,23,24]. Among the macromolecules that leak from the damaged tissues, enzymes, because of tissue specificity and catalytic activity, are released into the bloodstream, [25] and a study of these enzyme activities in plasma has been found to be of great importance in the assessment of liver damage [26].

The increased levels of AST and ALT are indicative of cellular damage and loss of functional integrity of the cell membrane in the liver [27]. The increase in ALP in liver disease is the result of increased synthesis of the enzyme through cells lining the canaliculi, usually either intra- or extrahepatic, which reflects the pathological alteration in biliary flow [28].

The present study demonstrated the ability of the TD extract to decrease the ALT and AST level significantly (*p*<0.05) at 500 mg/kg dose and also the ALP level at 250 mg/kg (*p*<0.01) and 500 mg/kg dose (*p*<0.001) (Table 2).

Reduction of the enhanced level of serum ALT, AST, ALP and total bilirubin by TD extract seemed to offer protection and maintain the functional integrity of hepatic cells.

An abnormal increase in the levels of bilirubin in plasma indicates hepatobiliary disease and severe disturbance of hepatocellular function [29]. Prior oral administration of *Trichosanthes dioica* (TD) extract exhibited significant protection against AAP-induced hepatotoxicity. It decreased the levels of bilirubin significantly (*p*<0.05) at 500 mg/kg which is an indication of protection against hepatic damage caused by AAP.

Most proteins found in plasma are produced by the liver, the principle exception being immunoglobulins. Severe liver damage has been associated with decreased production of various proteins resulting in reduced serum levels of total protein, albumin, and/ or globulin [30,31]. Decreased protein production may render other abnormal test values. e.g. depletion of coagulation factors (all are globulins) may result in prolonged prothrombin or activated partial thromboplastin times [32]. And, increased loss of protein via urine or feces due to renal or gastrointestinal disease will reduce serum protein levels. Inflammation anywhere in the body often results in increased production of specific globulin proteins produced by the liver [30].

Table 1. Effect of *Trichosanthes dioica* leaves on the body weight and liver weight in paracetamol-induced hepatotoxicity in rats

Group	Body Weight (gm) (Mean±SEM)		Liver Weight (gm) (Mean±SEM)
	Initial Day	Final Day	
Normal Control	148.33±2.95	162.17±2.36	4.83±0.23
Paracetamol Control	149.33±3.85	141.0±3.27	6.10±0.25
STD (Sylimarin 100 mg/kg)	150.0±4.66	145.67±4.70	5.25±0.18*
TD 250 mg/kg	148.33±4.78	141.83±5.01	5.87±0.10
TD 500 mg/kg	150.0±2.63	145.83±2.47	5.40±0.14

N.B: Data were analyzed by one way ANOVA following Bonferroni post hoc test. Values are expressed as Mean±SEM, n=6. *(p< 0.05) = significant, ** (p< 0.01) = highly significant, *** (p< 0.001) = very highly significant compared to paracetamol control

Table 2. Effect of *Trichosanthes dioica* leaves on plasma ALT, AST, ALP and bilirubin level in paracetamol-induced hepatotoxicity in rats

Group	ALT (IU/L) (Mean±SEM)	AST (IU/L) (Mean±SEM)	ALP (IU/L) (Mean±SEM)	Bilirubin (g/dl) (Mean±SEM)
Normal Control	36.37±6.82	43.33±1.86	167.17±4.89	1.17±0.13
Paracetamol Control	74.67±3.01	76.0±4.91	291.0±11.04	3.02±0.14
STD (Sylimarin 100 mg/kg)	44.67±2.22***	51.50±2.72***	138.67±2.74***	1.20±0.18***
TD 250 mg/kg	63.0±2.08	70.0±1.42	218.83±8.67***	2.57±0.17
TD 500 mg/kg	51.0±1.63**	54.50±1.96***	180.17±6.80***	1.80±0.22***

N.B: Data were analyzed by one way ANOVA following Bonferroni post hoc test. Values are expressed as Mean±SEM, n=6. *(p< 0.05) = significant, ** (p< 0.01) = highly significant, *** (p< 0.001) = very highly significant compared to paracetamol control

Table 3. Effect of *Trichosanthes dioica* leaves on plasma total protein, albumin and globulin in paracetamol-induced hepatotoxicity in rats

Group	T. Protein (g/dl) (Mean±SEM)	Albumin (g/dl) (Mean±SEM)	Globulin (g/dl) (Mean±SEM)
Normal Control	6.02±0.18	4.05±0.28	1.97±0.35
Paracetamol Control	4.43±0.15	2.40±0.21	2.03±0.32
STD (Sylimarin 100mg/ kg)	5.70±0.17***	3.93±0.37**	1.77±0.42
TD 250 mg/kg	4.77±0.11	3.02±0.15	1.75±0.24
TD 500 mg/kg	5.68±0.18***	3.77±0.16**	1.92±0.33

N.B: Data were analyzed by one way ANOVA following Bonferroni post hoc test. Values are expressed as Mean±SEM, n=6. *(p< 0.05) = significant, ** (p< 0.01) = highly significant, *** (p< 0.001) = very highly significant compared to paracetamol control

The results indicate that the protein level was slightly decreased at 250 mg/kg dose whereas it was slightly increased at 500 mg/kg dose. But the plasma albumin level was decreased in both the doses although it was not significant (Table 3).

Through this study also the consumption of TD for 7 days was found to reduce the body weight as well as the liver weight of rats (Table 1).

4. CONCLUSION

The overall significance of this study was that, the treatment with *Trichosanthes dioica* leaf extract was able to protect the changes induced

by AAP. On the basis of the above results it can be concluded that *Trichosanthes dioica* has significant value as hepatoprotective plant against paracetamol induced liver injury model. Further studies are required to determine the possible hepatoprotective mechanism(s) involved and, to isolate and identify the responsible bioactive compounds.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The study was approved by the institutional animal ethical committee.

ACKNOWLEDGEMENTS

Authors are very much thankful to Pharmacology laboratory, Dept. of Pharmacy, Jahangirnagar University, Savar, Dhaka.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ahsan R, Islam MK, Musaddik A, Haque E. Hepatoprotective activity of methanol extract of some medicinal plants against carbon tetrachloride induced hepatotoxicity in Albino rats. *Global J Pharmacol.* 2009;3: 116-22.
2. Ali M, Ramachandram R, Rafiullah MR, Singh O, Siddiqui AW, Mir SR. Prevention of carbon tetrachloride-induced hepatotoxicity by the ethanol extract of *Capparis mooniifruits* in rats. *Pharma Bio.* 2004;42: 286-88.
3. Ajith TA, Hema U, Aswathy MS. *Zingiber officinale* Roscoe prevents acetaminophen-induced acute hepatotoxicity by enhancing hepatic antioxidant status. *Food and Chemical Toxicology.* 2007;45:2267-2272.
4. Avila DS, Palma AS, Colle D, Scolari R, Manarin F, Da Silveira AF, Nogueira CW, Rocha JB, Soares FA. Hepatoprotective activity of a vinylic telluride against acute exposure to acetaminophen. *Eur J Pharmacol*; 2011.
5. Lee WM. Acetaminophen and the U.S. Acute Liver Failure Study Group: Lowering the risks of hepatic failure. *Hepatology.* 2004;40(1):6-9.
6. Saumendu Deb Roy, Sumit Das, Dibyendu Shil, Koushik Nandan Dutta. Herbal hepatoprotective agents: A review. *World Journal of Pharmaceutical Research.* 2012;1(2):87-99.
7. Karunakar Shukla. Bioassay- An uncomplicated methodologies for ensure safety of traditional formulations. *Research Journal of Pharmacognosy and Phytochemistry (RJPP).* 2009;1(1).
8. Mia AW, Ghani A. In: Ghani A (ed). *Traditional medicine, pharmacy department, Jahangirnagar University, Savar, Dhaka, Bangladesh.* 1990;10-12.
9. Kirtikar KR, Basu BD. *Indian medicinal plants.* New Delhi: Bishen Singh Mahendra Pal Singh. 1935;1110-1111.
10. Nadkarni KM. *Indian Materia Medica.* Bombay: Popular Prakashan. 1976;1236-1237.
11. Sharma PC, Yelne MB, Dennis TJ. *Database on medicinal plants used in Ayurveda.* New Delhi. Central Council for Research in Ayurveda and Siddha. 2002; 269-272.
12. Singh K, Pointed gourd (*Trichosanthes dioica* Roxb.). *Indian Hort.* 1989;33:35-38.
13. Mukherjee PK. *Quality control of herbal drugs.* New Delhi: Business Horizons. 2002;164-71.
14. Ghaisas MM, Tanwar MB, Ninave PB, Navghare VV, Deshpande T. Hepatoprotective activity of aqueous & ethanolic extract of *T. dioica* in ferrous sulphate induced liver injury. *Pharmacologyonline.* 2008;3:127-135.
15. Chopra RN, Nayar SL, Chopra IC. *Glossary of Indian medicinal plants.* CSIR, New Delhi. 2002;340.
16. Nadkarni AK, *Indian Materia Medica*, 3rd Edn, Bombay Popular Prakashan Mumbai. 1982;1236-1237.
17. Khare CP. *Encyclopedia of Indian medicinal plants.* Berlin, Heidelberg; New York: Springer-Verlag. 2004;458.
18. Rai PK, Rai DK, Jaiswal D, Sharma B, Watal G. Effect of water extract of *Trichosanthes dioica* fruits in streptozotocin induced diabetic rats. *Indian J Clin Biochem.* 2008;23:387-90.
19. Ji Su Kim, Jung Bong Ju, Chang Won Choi, Sei Chang Kim. Hypoglycemic and antihyperlipidemic effect of four Korean Medicinal plants in alloxan induced diabetes rats. *American Journal of Biochemistry & Biotechnology.* 2006;2(4): 154-160.
20. Mallhi TH, Qadir MI, Khan YH, Ali M. Hepatoprotective activity of aqueous methanolic extract of *Morus nigra* against paracetamol-induced hepatotoxicity in mice. *Bangladesh J Pharmacol.* 2014;9: 60-66.
21. Black M. Acetaminophen hepatotoxicity. *Gastroenterology.* 1980;78:382-392.
22. Pacifici GM, Back DJ, Orme ML. Sulphation and glucuronidation of paracetamol in human liver: Assay conditions. *Biochem Pharmacol.* 1988;37: 4405-4407.

23. Jollow DJ, Mitchell JR, Potter WZ, Davis DC, Gillette JR, Brodie BB. Acetaminophen-induced hepatic necrosis: II. Role of covalent binding *in vivo*. J Pharmacol Exp Ther. 1973;187:195–202.
24. Potter DW, Hinson JA. Reactions of N-acetyl-p-benzoquinoneimine with reduced glutathione, acetaminophen and NADPH. Mol Pharmacol. 1986;30:33–41.
25. Hearse DJ. Cellular damage during myocardial ischaemia: Metabolic changes leading to enzyme leakage. In: Enzymes in Cardiology (Hearse DJ, De Leiris J, Loisanse D, Eds.) John Wiley and Sons Ltd. New York. 1979;1–21.
26. Plaa GL, Zimmerman HJ. Evaluation of hepatotoxicity: Physiological and biochemical measures of hepatic function. In: Comprehensive Toxicology, (McCuskey RS, Earnest DL, Eds.) Cambridge University Press, Cambridge, UK. 1997;9: 97–109.
27. Drotman RB, Lowhorn GT. Serum enzymes as indicators of chemical induced liver damage. Drug Chem Toxicol. 1978;1: 163–171.
28. Plaa GL, Hewitt WR. Detection and evaluation of chemically induced liver injury. In: Principles and methods of toxicology, 2nd Ed. (Wallace Hayes A, Ed.), Raven Press, New York. 1989;399–428.
29. Martin P, Friedman LS. Assessment of liver function and diagnostic studies. In: Handbook of Liver Disease (Friedman LS, Keeffe EB, Eds.), Churchill Livingstone, Philadelphia. 1998;1–14.
30. Alper CA. Plasma protein measurement as a diagnostic aid. N. Eng. J. Med. 1974;291: 287-290.
31. Killingsworth LM. The role of high resolution electrophoresis in the clinical evaluation of protein status. Freehold, New Jersey: Worthington Diagnostics; 1981.
32. Badylak SF, Van Vleet JF. Alteration of prothrombin time and activated partial thromboplastin time in dogs with hepatic disease. Am. J. Vet. Res. 1981;42:2053-2056.

© 2018 Billaha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46204>