



## **Effect of Agronomic Management on Soil Microclimatic Parameters and Root Yield of Sida Hemp (*Sida alnifolia* L.)**

**P. T. Vidhu Priyadarsini<sup>a\*</sup>, P. V. Sindhu<sup>a</sup> and Meera V. Menon<sup>a</sup>**

<sup>a</sup> College of Agriculture, KAU P.O, Kerala Agricultural University, Thrissur 680656, Kerala, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2022/v34i232601

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94369>

**Original Research Article**

**Received 02 October 2022**  
**Accepted 07 December 2022**  
**Published 12 December 2022**

### **ABSTRACT**

Agronomic management exhibits great influence on growth and yield of crops by modifying physical, chemical and biological properties of soil and plants. Field experiment was conducted during June to December, 2018 to assess the effect of agronomic management viz. light intensity, application of manuring and weed management on soil microclimatic variations and its effects on root yield of Sida hemp (*Sida alnifolia* L.). The treatments included two levels of light intensity (open and 50 per cent shade), two levels of manuring (no manure and FYM@10 t ha<sup>-1</sup>), and four weed management practices (black polythene mulching, organic mulching with paddy straw @ 5 t ha<sup>-1</sup>, hand weeding at 1,3,5 MAP (Month after planting) and no weeding). Higher soil temperature at 10 cm depth was observed in open condition and under black polythene mulching throughout the growing period. Higher soil moisture content at 15 cm depth was recorded under shade and black polythene mulching. Black polythene mulching resulted in increase in soil temperature and soil moisture content by almost two percent as compared to bare soil in open condition. The highest root yield of 14.66 t ha<sup>-1</sup> was noticed in plots with black polythene mulching with FYM under open condition. Soil temperature and light intensity showed positive correlation with root yield.

**Keywords:** Soil temperature; soil moisture; light intensity; mulching; root yield.

## 1. INTRODUCTION

Plants accomplish a very vibrant role in sustaining and refining quality of human life. Plants which are commonly used for treating or preventing various ailments are generally considered as medicinal plants. Quality of raw drug is as important as its quantity in medicinal plant cultivation. Since bulk of the present requirement is met by wild collection from natural habitats, when the crop is brought under cultivation, with improved management techniques, it is indispensable to ensure its quality. Cultivating plants under a micro climate similar to its niche original is found to be the viable solution for ensuring its therapeutic properties.

*Sida alnifolia* is a species found in tropical and subtropical regions of India, belonging to the family Malvaceae. According to National Medicinal Plant Board, Sida is the 3<sup>rd</sup> most widely consumed drug in Ayurveda pharmaceutical industry [1]. Because of its high commercial value, the crop is included in the group of high volume traded medicinal plants sourced from waste lands. At present bulk of the present requirement is met in wild collection from natural habitats.

Management methods exhibit great influence on growth and yield of crops by way of modifying physical, chemical and biological properties of soil and plants. Organic manures provide a better environment for crop growth and root development by improving the soil physical, chemical, biological properties and supplying plant nutrients including micronutrients. According to Upadhyaya et al. [2], the yield, total phenol and total flavanoid contents of medicinal plant *Adhatoda vasica*, improved significantly by the application of organic manures.

Light is a physical factor which can influence growth, yield and secondary metabolites production. Both deficient and excessive light intensities may be injurious to plants and it will affect plant growth, development and yield [3].

Crop-weed competition is a common interaction occurring in cropped field significantly influencing crop growth and yield. Mulching is a non-chemical weed management method which modifies the plant microclimate by modifying soil temperature, soil moisture and evaporation and the modified microclimate in turn affects growth and development of crops. According to Lalitha et al. [4], moisture content, soil temperature and

nutrient availability increased under plastic mulching. Gunasekaran and Shakila [5] reported significant influence of mulching on tuber characters such as number of tubers, tuber length, tuber girth, and fresh tuber weight of medicinal coleus *Coleus forskohlii*.

As the information on influence of management methods on *Sida* hemp (*Sida alnifolia* L.) is limited, the present study was undertaken to assess the effect of agronomic management viz. light intensity, manuring and weed management on soil microclimatic variations and its consequent effects on root yield of *Sida* hemp (*Sida alnifolia* L.)

## 2. MATERIALS AND METHODS

The study was conducted at Agronomy farm, College of Horticulture, Kerala Agricultural University, Vellanikkara, Thrissur (India) during June to December, 2018. Monthly weather data for the experimental period was given in the Chart 1. The area located at 13° 32'N latitude and 76° 26'E longitude. The altitude of the place is 40 m above mean sea level. The experiment was laid out in Randomized Block Design (Factorial), with 3 factors and three replications. Factor A consist of two light intensity (open and 50 per cent shade), factor B consist of manuring (no manure and FYM@10 t ha<sup>-1</sup>), and factor C consist of four weed management practices (black polythene mulching, organic mulching with paddy straw @ 5 t ha<sup>-1</sup>, hand weeding at 1,3,5 MAP and no weeding). One month old healthy, uniform sized seedlings were selected and transplanted in the main field on 2<sup>nd</sup> June 2018 at a spacing of 50 cm x 25 cm. Soil temperature at 10 cm depth and soil moisture at 0-15 cm depth were recorded at weekly intervals using soil thermometer and gravimetric method respectively. The observation on root yield was recorded six months after planting (4<sup>th</sup> December 2018) at seed maturation stage by uprooting of whole plants. Field view under open and shaded condition were given in Fig.1 and Fig. 2 respectively.

The data collected were statistically analysed and presented as per procedure given by Sheoran et al. [6].

## 3. RESULTS AND DISCUSSION

### 3.1 Soil Temperature

The average soil temperature at 10 cm depth during the experiment period ranged from 24.1°C to 28.5°C. Throughout the growing period higher

soil temperature was observed in open condition as compared to shaded condition (Fig. 3). As per Onwuka and Mang [7], the soil temperature is a function of amount of solar radiation from the sun that reached on the soil surface and amount that absorbed by the soil. Martias and Musil [8] also reported increase in soil temperature with increased solar radiation that reached the soil surface. Significant influence of manuring on soil temperature have not been observed in this

experiment. Among different weed management practices, black polythene mulched soil recorded higher temperature (Fig. 4). The rise in soil temperature may due to solar energy trapped inside the mulch material through greenhouse effect [9]. Organic mulched soil showed lower soil temperature as compared to hand weeded soil (Fig. 4). The reduction in soil temperature may due to retaining soil moisture by reducing soil evaporation through organic mulching [10].




**Fig. 1. Field view under open growing condition**




**Fig. 2. Field view under shaded condition**

**Chart 1. Monthly weather data during experimental period (June 2018-December 2018)**

Months	Max temp. (°C)	Min temp. (°C)	RH %	Rainfall (mm)	Rainy days	Total Evp. (mm)	Sunshine hours
June	29.8	23.2	89	730.0	23	65.7	51.2
July	29.6	22.5	88	793.2	22	79.6	58.0
Aug	29.2	22.2	87	928.0	21	70.7	68.4
Sep	32.2	22.5	75	29.0	1	99.6	216.2
Oct	32.8	22.9	76	393.0	13	94.4	176.0
Nov	32.7	23.3	68	66.6	5	102.3	207.5
Dec	33.0	22.5	63	0.0	0	109.5	215.7



**Fig. 3. Effect of growing condition on soil temperature (°C) at 10 cm depth**



**Fig. 4. Effect of weed management on soil temperature (°C) at 10 cm depth**

### 3.2 Soil Moisture

The average soil moisture at 15 cm depth ranged from 12.7 per cent to 33.5 per cent throughout the growing season. Plots under shade recorded higher soil moisture content at 15 cm depth except from 2<sup>nd</sup> week to 13<sup>th</sup> week (Fig. 5). This exception was due to monsoon rains received during this period. Youn et al. [11] also observed higher soil moisture under shaded condition. Soil moisture content was unaffected by manuring. Weed management practices significantly influenced the soil moisture content at 15 cm depth. Black polythene mulching recorded higher moisture content as compared to other weed management practices except from 2<sup>nd</sup> week to 13<sup>th</sup> week (Fig. 6). According to Sakthivel [12], mulching with black polythene sheet conserved soil moisture by reducing rate of evaporation. Soil moisture content in organic mulched soil was found to be lesser as compared to no weeding soil from 14<sup>th</sup> week to 25<sup>th</sup> week (After south west monsoon). This may be due to increased weed biomass serve as protection for soil surface against weather aggressions to maintain soil moisture.

### 3.3 Root Yield

The highest root yield of 14.66 t ha<sup>-1</sup> was recorded in treatment combination, black polythene mulching with FYM under open condition followed by black polythene mulching without manure under open condition (Table 1). The lowest root yield was recorded in no weeding plots without manure under 50 per cent shade (5.14 t ha<sup>-1</sup>). Under best treatment combination of manuring and weed management (FYM @ 10 t ha<sup>-1</sup> x black polythene mulching), by altering only growing condition, a yield increase of 4.84 t ha<sup>-1</sup> could be observed. This

indicates the sun loving nature of *Sida alnifolia*. Optimum soil microclimatic conditions in this combination might have contributed to higher root yield.

### 3.4 Correlation Studies

Correlation between microclimatic factors and yield at vegetative and harvest stages is depicted in Table 2. At both stages, positive correlation was observed between light intensity, soil temperature and root yield. However, a negative correlation was observed between soil moisture and light intensity (-0.628 and -0.533 at vegetative and harvest stages respectively). Positive correlation between soil temperature and total yield using plastic mulch and row covers on cucumber was reported by Ibarra Jiménez et al. [13].

### 3.5 Economics

The data on economics (Rs ha<sup>-1</sup>) of cultivation of *Sida* hemp under different growing condition, manuring and weed management is furnished in Table 3. Highest cost of cultivation was observed with black polythene mulch with FYM @ 10t ha<sup>-1</sup> in open condition and in 50% shade, black polythene mulch with FYM @10t ha<sup>-1</sup> (Rs. 72250) and the lowest cost of cultivation was observed in no weeding without manure in open condition (Rs. 35,000). A higher B:C ratio of 1.73 was obtained from black polythene mulch with manure under open condition. Hand weeding with and without manure under shaded condition recorded lower B:C ratio (0.82).

- Labour charges (Rs.600/day)
- Cost of polythene sheet – Rs.7/m<sup>2</sup>
- Cost of FYM – Rs. 1200/tonne
- Cost of organic mulch – Rs.6/Kg

**Table 1. Effect of growing condition, manuring and weed management on root yield of *Sida* hemp**

Treatments	Root yield (t ha <sup>-1</sup> )	
	Harvest	
	Open	50 per cent shade
No manuring x Black polythene	11.92	8.06
No manuring x Organic mulch	7.45	5.95
No manuring x Hand weeding	7.09	5.50
No manuring x No weeding	6.05	5.14
FYM @ 10 t ha <sup>-1</sup> x Black polythene	14.66	9.82
FYM @ 10 t ha <sup>-1</sup> x Organic mulch	8.85	6.18
FYM @ 10 t ha <sup>-1</sup> x Hand weeding	7.67	6.10
FYM @ 10 t ha <sup>-1</sup> x No weeding	6.89	5.87
CD (0.05)	2.09	




Fig. 5. Effect of growing condition on soil moisture (%) at 0-15cm depth




Fig. 6. Effect of weed management on soil moisture (%) at 0-15cm depth

Table 2. Correlation between microclimate factors and root yield at vegetative and harvesting stage

Vegetative stage				
	Soil temperature	Soil moisture	Light intensity	Yield
Soil temperature	1.000			
Soil moisture	-0.391**	1.000		
Light intensity	0.180	-0.628**	1.000	
Yield	0.373**	-0.041	0.449**	1.000
Harvesting stage				
	Soil temperature	Soil moisture	Light intensity	Yield
Soil temperature	1.000			
Soil moisture	-0.357**	1.000		
Light intensity	0.254	-0.533**	1.000	
Yield	0.596**	0.161	0.452**	1.000

**Table 3. Interaction effect of growing condition, manuring and weed management on benefit: Cost ratio**

Treatments	Cost of cultivation (Rs ha <sup>-1</sup> )	Gross income (Rs ha <sup>-1</sup> )	Net income (Rs ha <sup>-1</sup> )	B:C ratio
A1B1C1	66250.00	101354.00	35104.00	1.53
A1B1C2	43688.00	63342.00	19654.00	1.45
A1B1C3	57250.00	60259.33	3009.33	1.05
A1B1C4	35000.00	51396.67	16396.67	1.47
A1B2C1	72250.00	124610.00	52360.00	1.73
A1B2C2	49688.00	65223.33	15535.33	1.31
A1B2C3	63250.00	75242.00	11992.00	1.19
A1B2C4	41000.00	58559.33	17559.33	1.43
A2B1C1	66250.00	68464.67	2214.67	1.03
A2B1C2	43688.00	50580.67	6892.67	1.16
A2B1C3	57250.00	46727.33	-10522.67	0.82
A2B1C4	35000.00	43644.67	8644.67	1.25
A2B2C1	72250.00	83447.33	11197.33	1.15
A2B2C2	49688.00	52598.00	2910.00	1.06
A2B2C3	63250	51850	-11400	0.82
A2B2C4	41000	49900.67	8900.67	1.22

A1 – Open; C1 – Black polythene sheet; A2 – 50% shade; C2 – Organic mulch; B1 – No manure; C3 – Hand weeding; B2 – FYM @ 10t/ha; C4 – No weeding

#### 4. CONCLUSION

The current research work was carried out to assess the effect of variations in growing conditions, manuring and weed management on soil microclimatic parameters and root yield of *Sida* hemp. Growing condition significantly influenced the soil temperature, soil moisture and root yield of *Sida* hemp. Eventhough, manuring had significant effect on root yield, soil microclimatic parameters were not influenced by manuring. Among different weed management methods, black polythene mulching showed higher soil temperature, soil moisture (except from 2<sup>nd</sup> week to 13<sup>th</sup> week) and root yield. From the present study, the combination of open condition, application of FYM @ 10 t ha<sup>-1</sup> and weed management by black polythene mulching can be recommended as optimum agronomic management for better soil microclimatic parameters and root yield of *Sida* hemp.

#### ACKNOWLEDGEMENTS

The authors are highly thankful to the College of Agriculture, Vellanikkara, Thrissur, Kerala for providing all assistance and research material.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Ved DK, Goraya GS. Demand and supply of medicinal plants in India. NMPB, New Delhi. 2007;18(85):10-52.
2. Upadhyaya S, Khativora E, Saikia LR. Comparison of total flavonoid content in *Adhatoda vasica* Nees. Grown using different organic manure. J Pharmacy Res. 2010;3(10):2408-2409.
3. Safeer PM, Sreekumar S, Krishnan PN, Biju CK, Seeja G. Influence of soil texture and bed preparation on growth performance in *Plectranthus vettiveroides*. J Agric Vete Sci. 2013;5:41-45.
4. Lalitha M, Thilagam KV, Balakrishnan N, Mansour M. Effect of plastic mulch on soil properties and crop growth-a review. Agr Rev. 2010;31(2):145-149.
5. Gunasekaran P, Shakila A. Effect of mulching on weed control and tuber yield of medicinal coleus (*Coleus forskholli* Briq.) Asian J Hort. 2014;9:124-127.

6. Sheoran OP, Tonk DS, Kaushik LS, Hasija RC, Pannu RS. Statistical Software Package for Agricultural Research Workers. CSHAU, Hisar; 1998.
7. Onwuka B, Mang B. Effects of soil temperature on some soil properties and plant growth. *Adv. Plants Agric. Res.* 2018;8(1):34-37.
8. Martias AD, Musil S. Temperature and thermal diffusivity within a range land soil near Oracle, Arizona. *J. Arizona–Nevada Academy Sci.* 2012;44(1):15–21.
9. Abhivyakthi. Kumari P. Ojha RK. Job M. Effect of plastic mulches on soil temperature and tomatoyield inside and outside the polyhouse. *Agric. Sci. Digest.* 2016;36:333–336.
10. Zhang J, Hang X, Lamine SM., Jiang Y, Afreh D, Qian H, Feng X, Zheng C, Deng A, Song Z, Zhang W. Interactive effects of straw incorporation and tillage on crop yield and greenhouse gas emissions in double rice cropping system. *Agric Ecosystems Environ.* 2017;250:37-43.
11. Youn WB, Hernandez JO, Park BB. Effects of shade and planting methods on the growth of *Heracleum moellendorffii* and *Adenophora divaricata* in different soil moisture and nutrient conditions. *Plants.* 2021;10(10):2203.
12. Sakthivel N. Impact of black polythene mulching in mulberry garden on weed infestation, soil moisture, plant growth and leaf yield under tropical conditions. *Int. J. Tropical Agric.* 2019;37(1):89-95.
13. Ibarra-Jiménez L, Quezada-Martín MR, Rosa-Ibarra M. The effect of plastic mulch and row covers on the growth and physiology of cucumber. *Aust. J Exp. Agr.* 2004;44(1):91-94.

© 2022 Priyadarsini et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/94369>