



Studies on the Effect of Foliar Spray of Micronutrients on Physical Parameters of Guava (*Psidium guajava* L.) cv. Allahabad Safeada

Usha Shukla ^a, Rubee Lata ^a, Shatrunjay Yadav ^{a*} and Pradeep Kumar ^b

^a Department of Horticulture, School of Agricultural Sciences and Technology, Babasaheb Bhimrao Ambedkar University (A Central University), Vidya-vihar, Rae Bareilly Road, Lucknow-226025, India.

^b Chandra Shekhar Azad University of Agriculture & Technology Kanpur, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2022/v41i2131753

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85195>

Original Research Article

Received 04 April 2022
Accepted 08 June 2022
Published 28 June 2022

ABSTRACT

A field experiment was conducted on Horticulture Research Farm –II of Department of Horticulture, School of Agricultural Sciences and Technology, Babasaheb Bhimrao Ambedkar University, Lucknow, India, during the academic year of 2019-20. to study the effect of micronutrient spray on the physical parameters of guava cv. Allahabad Safeda on 10 - year- old guava plants that fruit set (68.80 %) fruit retention (65.89%) fruit length (7.59 cm), fruit width (6.75 cm), fruit weight (136.8 g.) fruit volume (148.13cm³), specific gravity(0.92), fruit yield per plant (42.20kg/plant), and yield (q/ha) were obtained maximum with the foliar spray of zinc sulphate T₃ (0.4%) and borax T₉ (0.4%) respectively.

Keywords: Guava; micronutrients; physical parameters; *Psidium guajava*.

1. INTRODUCTION

"The superfruit guava (*Psidium guajava* L.), also known as the "Apple of the Tropics" is one of the

most popular fruits cultivated in India's tropical, sub-tropical, and dry areas. About 150 species of tiny shrubs and trees belong to the genus *Psidium*, which belongs to the Myrtaceae family"

*Corresponding author: E-mail: sattu0064@gmail.com;

[1]. "About 20 species have edible fruits of which the most commonly cultivated is the common guava (*Psidium guajava* L.) was introduced in India in the 17th century by Portuguese from Latin America [2] and has originated in tropical America perhaps from Mexico". "Guava is widely grown all over the country in the kitchen garden, near the well and tube well premises, and also on a commercial scale" [3]. "Guava fruits are also used for the preparation of jam, jelly, RTS, nectar, etc. The guava bear flowers and fruits on current season growing twinges and highly cross-pollinated crop and pollination occurs through honey bees and andirons insect. The guava fruit developed from the inferior ovary exhibited a double sigmoid growth curve, fruit with many seed berry, the fruit takes nearly 4-5 month from dark green to yellowish-green" [4]. "Nutritional deficiencies in fruit crops significantly hamper the physiological process of plants thus reducing yield and producing inferior fruit, and making the plant vulnerable to a number of biotic and abiotic stresses. Micronutrients have received greater attention for crop production nowadays, because of their widespread deficiency in soil due to intensive cropping production and insufficient use of organic manure compared to high mineral fertilization for fruit crop production" [5]. "It formed an outstanding source of antioxidants such as vitamin C (260mg/100gm), carotenoids, and polyphenols. The fruit comprises three to fourfold higher amounts of vitamin C as a single orange. Vitamin C is essential for immune system stimulation, connective tissue formation as well as to reduce the incidence of degenerative diseases such as arthritis, arteriosclerosis, and cancer. In addition, antioxidants are known to retard aging as well as prevent or delay oxidative damage of lipids, proteins, and nucleic acids caused by reactive oxygen species" [6] "It has also an important role in fruit quality. Boron has an effect on the cell wall structure and also has a major effect on cell elongation. Zinc is involved in many enzymatic reactions in the growth and development of plants. It is also involved in regulating protein and carbohydrate metabolism. Copper plays a very important role in plant growth and nutrition, as well as in cell wall deposition. It increases fruit weight and yields also. Foliar application of nutrients enhances the efficient use along with the quick response" [7]. "The experiment was undertaken to study the effect of foliar application of mineral nutrients viz., boron (B), zinc (Zn), and copper(Cu) sprayed at different growth stages viz., at fruit set and at two weeks after fruit set of guava (*Psidium guajava* L.). An increase in the

plant canopy and minimum seed index was in Zn (0.4%) sprayed at the fruit set. Fruit size, sugar: acid. Increased yield in Zn (0.4%) and pectin enhances with B (0.3%) was found at two weeks after the fruit set, respectively. Minimum physiological loss in weight of fruits was with Cu (0.4%) sprayed at two weeks after fruit set" [7].

2. MATERIALS AND METHODS

The experiment was conducted at Horticulture Research Farm –II of Department of Horticulture, School of Agricultural Sciences and Technology, Babasaheb Bhimrao Ambedkar University, Lucknow, India, during the academic year of 2019-20. The plants having uniform vigor, size with the age of 12 years old were selected for the study. The experimental site has a sub-tropical climate. The soil of the orchard is clay loam which is well-drained and well aerated. The soil texture was quite loose suited for the development of the plant root system. Nutrient application and other orchard management practices were followed as per recommended package and practices for guava. Manure and fertilizer were applied in each treatment before flowering in the month of July for taking winter season crop. The first spraying of micronutrients was done before flowering (first week of August) and second after fruit set (second week of September) during 2019-20. The experiment consisted of 22 treatment combinations of 3 micronutrients viz., zinc, copper, and boron at 0.2%, 0.3%, and 0.4% and is a combination of two and three except spray of plain tap water as control. Aqueous solutions of zinc, copper, boron, and tap water were sprayed at the time of full bloom on rainy season crops (Ambe Bahar). The experiment was laid out in Randomized Block Design (RBD) with 3 replications of all the treatments. Micronutrient sprays were done at the full bloom stage in the early morning with the help of a foot sprayer @ six liters per tree to ensure the maximum absorption of nutrients through the leaves. Zinc (as Zinc sulfate - $ZnSO_4$), Copper (as Copper sulfate - $CuSO_4$), Boron (as Boric acid - H_3BO_3) with concentrations of 0.2% (2000 ppm) 0.3% (3000 ppm) 0.4% (4000 ppm), Control-Spray of tap water.

Experimental Treatments and Measurements:

Treatment combinations are T_1 - Zinc sulphate (0.2%) , T_2 - Zinc sulphate (0.3 %), T_3 - Zinc sulphate (0.4 %), T_4 - Copper sulphate (0.2 %), T_5 - Copper sulphate (0.3%), T_6 -Copper sulphate (0.4 %), T_7 -Boric acid (0.2 %), T_8 -

Boric acid (0.3%), T₉ -Boric acid (0.4 %), T₁₀ - Zinc sulphate + copper sulphate (0.2 %), T₁₁ - Zinc sulphate + copper sulphate (0.3 %), T₁₂ Zinc sulphate + copper sulphate 90.4 %), T₁₃- Zinc sulphate + boric acid(0.2 %), T₁₄ -Zinc sulphate + boric acid(0.3 %), T₁₅ -Zinc sulphate + boric acid(0.4 %), T₁₆ Copper sulphate + boric acid(0.2 %), T₁₇- Copper sulphate + boric acid(0.3 %), T₁₈ -Copper sulphate + boric acid (0.4 %), T₁₉ Zinc sulphate + copper sulphate + boric acid(0.2 %), T₂₀- Zinc sulphate + copper sulphate+ boric acid(0.3 %), T₂₁- Zinc sulphate + copper sulphate+ boric acid(0.4 %), T₂₂- Control spray of tap water.

Measurement of Physical Properties of Fruit:

To determine fruit morphological characteristics like fruit size (length and width), fruit weight, fruit volume, and fruit retention. Twenty fruits were selected randomly from replicates. The specific gravity was calculated by dividing the weight of the fruit by its volume. Fruit firmness of guava was determined by a screw-type Penetrometer and the reading was expressed in kg/cm².

Statistical Analysis: The experimental data for two successive years 2019 and 2020 was subjected to statistical analysis. The critical differences (C.D.) at a 5% level of probability

were worked out for comparing treatment means. They were analyzed according to the procedure of analysis for Randomized Block Design (RBD) given by Cochran and Cox (1992).

3. RESULTS AND DISCUSSION

A perusal of data in a Table-1 that foliar spray of micronutrients proved significantly effective in improving the percentage of fruit set. The maximum fruit set (68.80 percent) was found with the foliar application of borax (0.4%) in T₉ followed by 66.70 percent in T₃ with foliar application of zinc sulfate (0.4%) in T₃. The plant under control showed a minimum fruit set (54.80%) in T₂₂. The application of various treatments significantly influenced the percentage of fruit retention over the control (T₂₂). The Maximum fruit retention % was found in spray of borax(0.2%) in T₇ (65.89%) followed by spray of Borax (0.4%) in T₉ (65.80%). The minimum fruit retention (57%) was found in control (T₂₂). Similar results were also observed by Ali et al. [8] and Prasad et al. [9]. In comparison to the control, foliar micronutrient spray improves fruit size (length and breadth). With foliar feeding of copper sulfate 0.4 percent (T₅) and borax 0.4 percent (T₉), the greatest fruit

Table 1. Effect of foliar application of micronutrients on physical parameters of winter season guava (*Psidium guajava* L)

Treatments	Fruit set(%)	Fruit retention(%)	Fruit length (cm)	Fruit width (cm)	Fruit weight (g.)	Fruit volume (cm ³)	Specific gravity	Fruit Yield (kg/plant)
Zn (0.2%)	64.85	55.50	5.60	5.90	125.54	142.60	0.93	39.30
Zn (0.3%)	64.23	55.36	5.88	5.96	123.45	140.20	0.91	41.00
Zn (0.4%)	66.70	61.50	6.36	6.25	132.50	144.60	0.92	41.55
Cu (0.2%)	64.66	56.23	6.30	5.93	126.06	141.21	0.91	40.12
Cu (0.3%)	63.25	55.36	7.59	5.92	130.60	140.31	0.91	41.25
Cu (0.4%)	62.87	56.23	6.75	5.89	130.07	142.21	0.92	39.98
B (0.2%)	65.91	65.89	7.10	6.10	127.42	141.71	0.93	40.30
B (0.3%)	65.38	55.23	5.67	6.00	131.81	141.70	0.92	33.65
B (0.4%)	68.80	65.80	7.10	6.75	137.70	148.70	0.93	42.30
Zn+Cu (0.2%)	65.00	54.36	5.25	5.96	118.23	144.36	0.91	39.21
Zn+Cu (0.3%)	65.27	56.36	5.36	5.98	111.27	136.21	0.91	41.36
Zn+Cu (0.4%)	65.36	56.14	5.66	6.12	131.23	135.23	0.92	40.36
Zn+B (0.2%)	64.69	60.36	5.45	6.13	131.32	132.65	0.92	40.98
Zn+B (0.3%)	64.65	60.95	5.64	6.10	117.43	140.23	0.91	41.21
Zn+B (0.4%)	64.32	57.95	5.85	5.80	121.64	141.25	0.92	40.32
Cu+B(0.2%)	60.36	59.36	5.13	5.94	120.94	142.65	0.92	40.65
Cu+B (0.3%)	62.45	58.23	5.24	5.96	117.03	141.65	0.92	41.32
Cu+B (0.4%)	62.54	61.21	5.46	6.10	120.61	142.85	0.91	40.22
Zn+Cu+B(0.2%)	64.25	63.21	5.34	5.99	113.16	142.58	0.91	41.32
Zn+Cu+B(0.3%)	63.45	60.35	5.61	5.89	128.13	143.54	0.92	41.81
Zn+Cu+B(0.4%)	64.00	60.34	5.78	5.94	130.36	143.21	0.91	41.50
Control	54.80	57.00	4.89	5.20	115.27	128.60	0.90	27.50
S Em.	1.65	1.30	0.033	0.19	0.480	1.75	0.00	1.35
CD at 5%	5.10	4.03	0.059	0.59	0.874	5.39	NS	4.17

size was reported in terms of fruit length (7.59cm) and fruit width (6.75cm). Increased size due to borax spraying, which delivers boron to the plant, may have controlled cell wall permeability, enabling more water mobilization in the fruit, resulting in higher fruit size. These results are in close conformity with the findings of Prasad et al., Pal et al. [10] in guava. All of the micronutrients considerably increased the weight of the fruit when compared to the control. The maximum fruit weight (136.8 g.) was achieved with foliar sprays of borax 0.4 percent (T₉) and zinc sulphate 0.4 percent (T₃), while the minimum fruit weight was achieved with the control. Increased fruit weight appears to have an indirect impact in hastening cell division and cell elongation, as well as increasing fruit size. Trivedi et al. [11] observed similar findings in guava. The results show that different treatments enhanced the fruit volume substantially above the control. The treatment T₉ produced the largest fruit volume (148.13 cm³) (borax 0.4 percent). This may be attributed to the role of borax in cell division, cell growth, and increased intercellular space in monocarpic cells, which results in increased fruit weight and size. Similar results were also obtained by Pal et al. [10] in guava. The maximum specific gravity (0.92) was found with foliar spray of borax 0.4 percent (T₉). The minimum specific gravity (0.90) was found in the control. The results are similar to the findings of Trivedi et al. [11] in guava. The maximum yield (42.20 kg/tree) was recorded with borax 0.4 percent (T₉) followed by zinc sulphate + copper sulphate 0.3 percent (T₁₁). These micronutrients improve the fruit length, width, and weight which ultimately increased the yield. These results are in close conformity with the findings of Prasad et al. [9], Trivedi et al. [11] in guava.

4. CONCLUSION

According to the present investigation, the physical characters of guava fruit with respect to fruit length, fruit width, fruit weight, fruit volume(cm³), etc were obtained maximum with the foliar spray of zinc sulphate T₃ (0.4%) and borax T₉(0.4%) respectively. Therefore, it may be concluded that foliar spray of zinc sulphate (0.4%) and borax (0.4%) can be recommended to the guava growers for obtaining a better quality of winter season guava fruits.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar R, Tiwari R, Kumawat BR. Quantitative and qualitative enhancement in guava (*Psidium guajava* L.) cv. 'Chittidar' through foliar feeding. *International Journal of Agricultural Sciences*. 2013;9(1):177–181.
2. Singh G, Singh AK, Pandey D. Effect of cropping pattern on fruiting behavior of guava (*Psidium guajava* L.) Tree American Agricultural Research. 2000;22:175–182.
3. Bal JS, Dhaliwal GS. Distribution and quality characteristics of graded guava fruits. *Haryana J. Hort. Sci*. 2004;33:53–54.
4. Akash Shukla RS, Verma, Som Prakash, Dharmpal Singh, and Rajesh Kumar. Effect of Foliar Application of Nutrients and Plant Growth Regulators on Quality Parameters of Guava (*Psidium guajava* L.) Fruit cv. Lalit. *International Journal of Current Microbiology and Applied Sciences*. 2019;8(5):956-959.
5. Sayan Sau, Sukamal Sarkar, Bikash Ghosh, Krishnedu Ray, Prahlad Deb and Dibakar Ghosh. Effect of Foliar Application of B, Zn and Cu on Yield, Quality and Economics of Rainy Season Guava Cultivation, *Current Journal of Applied Science and Technology*. 2018;28(1):1–10.
6. Bhatia SK, Yadav S, Ahlawat VP, Dahiya SS. Effect of foliar application of nutrients on the yield and fruit quality of winter season guava cv. L-49. *Hr. J Hort. Sci*. 2001;30(1&2):6-7.
7. Ajay Yadav, Verma RS, Ram RB, Viplaw Kumar, Ram Kewal Yadav. Effect of foliar application of micronutrients on physical parameters of winter season guava (*Psidium guajava* L) cv. Lalit. *Plant Archives*. 2017;17(2):1457-1459.
8. Ali W, Pathak RA, Yadav AL. Effect of foliar application of nutrients on guava (*Psidium guajava* L.) cv. Allahabad Safeda. *Prog. Hort*. 1993;23(1-4):14-21.
9. Prasad B, Das S, Chatterjee D, Singh UP. Effect of foliar application of urea, zinc, and boron on yield of guava. *J. of Applied Bio*. 2005; 15(1):44-47.
10. Pal, Anju; Pathak RK, Pal Krishna, Singh, Tejbir. Effect of foliar application of nutrients on yield and quality of guava

- (*Psidium guajava* L.) fruits cv. Sardar. *Prog. Res.* 2008;3(1):89-90.
11. Trivedi N, Singh D, Bahadur, V, Prasad VM, Collis JP. Effect of foliar application of zinc and boron on yield and fruit quality of guava (*Psidium guajava* L.). *Hort Flora Research Spectrum.* 2012;1(3):281-283.

© 2022 Shukla et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85195>