



# Effect of *Allium cepa* (Onion) Bulb and *Annona muricata* (Soursop) Pulp Juices on Body Weight, Prostate Weight, Selected Hormones and Prostate-Specific Antigen (PSA) in Testosterone and Oestradiol-Induced Benign Prostatic Hyperplasia in Albino Rats

Ibegbulem C. O.<sup>a</sup>, Ujowundu C. O.<sup>a</sup>, Ukairo D. I.<sup>a</sup>  
and Ihejieta H. A.<sup>b\*</sup>

<sup>a</sup> Department of Biochemistry, School of Biological Sciences, Federal University of Technology, Owerri, Nigeria.

<sup>b</sup> Department of Microbiology/Biochemistry, School of Industrial and Applied Sciences, Federal Polytechnic Nekede, Owerri, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. Author ICO designed the study. Author UCO handled the literature searches. Author UDI wrote the protocol. Author IHA managed the analyses of the study, performed the statistical analysis and wrote the first draft of the manuscript. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJRB/2023/v13i3264

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107355>

Original Research Article

Received: 05/08/2023  
Accepted: 09/10/2023  
Published: 17/10/2023

\*Corresponding author: E-mail: awuzieha1221p@gmail.com;

## ABSTRACT

**Aim:** This study investigated the effect of soursop pulp and onion bulb juice samples on body weight, prostate weight (PW), percentage prostate increase inhibition (PPII), dihydrotestosterone (DHT), testosterone (T), prolactin and PSA levels in testosterone and oestradiol-induced BPH in albino rats.

**Study Design:** The simple random design was used in this study.

**Place and Duration of Study:** Dave Research House, Federal University of Technology, Owerri, Nigeria, between April 2022 and July 2022.

**Methodology:** Fifty-four (54) male albino rats were randomly distributed into 9 groups (A-I) of 6 rats each, with B to I induced with 10 mg/kg b.w of exogenous hormones (1:1) every 2 days. They were treated as follows: A (normal control), B (BPH control), C (standard control), D (10 ml/kg b.w soursop), E (10 ml/kg b.w onion), F (10 ml/kg b.w, 1:1 soursop and onion), G (20 ml/kg b.w soursop), H (20 ml/kg b.w onion), and I (20 ml/kg b.w, 1:1 soursop and onion), for 28 days. The rats were weighed weekly. At the end of the treatment, they were euthanized, prostate and serum samples were collected for further assessments.

**Results:** The results showed that at low dose, the body weights of the rats were maintained. The single samples at low dose improved PW and PPII better. DHT and T levels reduced in all test groups, and the standard control. The single samples reduced DHT levels with increase in concentration, while the soursop juice and combination samples decreased testosterone levels with increase in concentration. Prolactin levels significantly ( $p < .05$ ) reduced in the test groups and standard control when compared with the BPH control. While, PSA levels also decreased in the test groups, but were not significantly ( $p > .05$ ) different from the controls.

**Conclusion:** The study suggested that soursop pulp and onion bulb juices exerted ameliorative effects on BPH, especially individually, and at the low dose studied. Hence, their consumption by older males should be encouraged.

**Keywords:** *Allium cepa bulb; Annona muricata pulp; benign prostatic hyperplasia; hormones; juice.*

## 1. INTRODUCTION

Benign prostatic hyperplasia (BPH), also referred to as benign prostatic hypertrophy [1], is a disease condition that is common in older men. It is a progressive condition identified by an enlarged prostate resulting from prostatic cells proliferation [2], and accompanied by lower urinary tract symptoms (LUTS) [3]. According to Lukacz *et al.* [4] and Lepor [5], LUTS can be grouped into two; filling/irritative symptoms (nocturia, dysuria, and increased urinary urgency and frequency) and obstructive symptoms in voiding (poor stream, hesitancy, incomplete voiding, terminal dribbling, and overflow incontinence). The pathophysiology of BPH is still not fully understood, but several pathways/mechanisms appear to be implicated in the onset and progression of BPH. Although, aging is the primary mechanism involved, research has demonstrated the critical involvement of hormonal changes/alterations, metabolic syndrome, and inflammation [6]. Prostate weight increases with age [7], while DHT, testosterone and prolactin have been noted to increase prostate volume [8][9].

During the treatment of BPH, the chemotherapeutic agents used are mainly the alpha blockers and the 5  $\alpha$ -reductase inhibitors. The alpha blockers relax prostate and bladder neck muscles allowing easy flow of urine, while the 5  $\alpha$ -reductase inhibitors stop further enlargement or even causes shrinking of prostate. These drugs have side effects. The alpha blockers produce dizziness, fainting, headaches, light-headedness, low blood pressure and occasionally, retrograde ejaculation, while the 5  $\alpha$ -reductase inhibitors produce low libido, erectile dysfunction, as well as, retrograde ejaculation, as side effects. More than 42 % of men between the ages of 51 to 60, 70 % of men between 61-70 years of age and as many as 90 % of men from 80 and above are affected by BPH [10]. LUTS adversely affects the quality of life of BPH patients. BPH still has no known medicinal cure. The primary goal of BPH management has been to provide relief-treatment for the symptoms of the condition. However, it is very common for the condition to reoccur once treated. Hence, there is pressing need to discover and develop phytotherapeutic agents with curative effect on BPH and related diseases.

Medicinal plants are one of the major source of complementary and alternative medicine. Phytotherapeutic agents are widely perceived to be more cost-effective, safer and have fewer side effects compared with conventional therapy for the management of ailments. This belief is responsible for the growing interest in their development [11][12]. *Allium cepa* Linn (Onion) is the most cultivated member of the genus *Allium*, native to Europe, Asia, North America and Africa. After tomatoes, it is the second most significant horticultural crop, with current production being around 44 million tonnes, worldwide. Onion is a common ingredient in many dishes. It is increasingly consumed due to its flavour and health benefits. Onions are rich in flavonoids and the alk(en)yl cysteine sulphoxides (ACSOs), which are beneficial to human health. The two flavonoid subgroups; anthocyanins and flavanols such as quercetin and its derivatives, found in onion play major role in its antioxidant activity [13]. Research has reported that compounds from onion possess anticarcinogenic properties, antiplatelet activity, antithrombotic activity, antiasthmatic and antibiotic effects [14].

*Annona muricata* L. (soursop or graviola) is a member of the Annonaceae family comprising about 130 genera and 2300 species [15]. It is native to the warmest tropical areas in South and North America and now widely distributed throughout tropical and subtropical parts of the world, including India, Malaysia and Nigeria [16]. *A. muricata* is an erect evergreen, terrestrial tree that can grow to a height of 5-8 m, featuring an open roundish canopy with large, glossy, dark green leaves. During fruiting season, the tree produces edible fruit that are large, heart shaped and green in colour, with diameter ranging from 15 to 20 cm [17]. In traditional medicine therapy, phytochemicals from soursop plant have a long history of use. Cancer, bacterial or parasitic infections, fever, high blood sugar, high blood pressure, inflammation, and anxiety, are a few of the conditions they are used to treat. Various scientific research results have shown that soursop extract contains antibacterial, antiprotozoan, anti-inflammatory, antioxidant, and anticancer effects [18].

This study reports the effect of *Annona muricata* (Soursop) pulp and *Allium cepa* (Onion) bulb juice samples on body weight, prostate weight (PW), percentage prostate increase inhibition (PPII), and serum levels of DHT, testosterone, prolactin and prostate-specific antigen (PSA) in testosterone and estradiol-induced BPH in albino rats.

## 2. METHODOLOGY

### 2.1 Sample Collection, Identification and Preparation

Bulbs of common onion and fresh ripe fruits of soursop were purchased at Relief Market, Owerri, Nigeria. They were identified by Dr. C. M. Duru, a taxonomist in the Department of Biology, Federal University of Technology, Owerri, Nigeria, with authentication numbers; FHI 110177 (*Annona muricata* L.) and FHI 107561 (*Allium cepa* L.). They were cleaned and washed with clean running water. The soursop fruit was cut open to collect the pulp, and remove all the seeds. The onion was cut into smaller pieces. The onion and soursop pulp were blended separately. They were then filtered with muslin cloth and the filtrates collected into separate specimen bottles for use and stored at 4 °C every other day as these samples were prepared every 2 days.

### 2.2 Animal Handling

A total of 54 male albino rats of Wistar strain were used for this study. The rats were acclimatized in different cages for 14 days. The simple random sampling technique was used in the grouping. The animals were maintained under standard environmental conditions (12 h light and 12 h dark cycle, uniform temperature of  $28 \pm 5$  °C), with free access to food (Vital Feed Finisher Pellets, Ibadan) and water, throughout the period of investigation. All investigations were conducted in accordance with the accepted principles for animal care and use [19].

### 2.3 Induction of Benign Prostatic Hyperplasia

The BPH was induced by administration of 10 mg/kg b.w of exogenous testosterone (CAS 57-85-2) and oestradiol (CAS 979-32-8) in the ratio of 1:1 intraperitoneally (every 2 days) for four weeks. The steroids were diluted in olive oil as solvent [20].

### 2.4 Animal Grouping and Treatment Administration

Treatments were administered daily for 28 d through oral gavage. At the end of the treatment, the rats were fasted overnight and euthanized by cervical dislocation. Blood was collected by ocular puncture into plain sample tubes and allowed to clot. Serum was separated within one

**Table 1. Grouping and treatment of the experimental animals**

Groups	Number of rats	Treatment for 28 days
A (normal control)	6	Water.
B (BPH control)	6	BPH induced rats + solvent. (Untreated)
C (standard control)	6	BPH induced rats + Finasteride (standard drugs - CAS 98319-26-7) at 10 mg/kg bw.
D	6	BPH induced rats + 10 ml/kg bw of soursop juice. (10 ml/kg AM)
E	6	BPH induced rats + 10 ml/kg bw of onion juice. (10 ml/kg AC)
F	6	BPH induced rats + 1:1 combination of soursop and onion juices (10 ml/kg bw). (10 ml/kg AMAC)
G	6	BPH induced rats + 20 ml/kg bw of soursop juice. (20 ml/kg AM)
H	6	BPH induced rats + 20 ml/kg bw of onion juice. (20 ml/kg AC)
I	6	BPH induced rats + 1:1 combination of soursop and onion juices (20 ml/kg bw). (20 ml/kg AMAC)

Key:AM - *Annona Muricata*; AC – *Allium Cepa*; AMAC - *Annona Muricata and Allium Cepa*

hour of blood clotting with a Pasteur pipette after centrifugation (centrifuge, MSLZL09) at 3000 x g for 20 min. the serum samples were used for biochemical tests. The prostates were excised and rinsed in ice-cold 1.15 % KCl, blotted and weighed using an electronic weighing balance (SUN-224CL-220 Gm) [21].

### 2.5 Determination of Percentage Change in Body Weight

The body weights of the rats were measured using an electronic weighing balance (SUN-224CL-220 Gm) and recorded before commencement of experiment, and subsequently once a week till sacrificed [22].

The percentage change in weight (per week) was calculated as follows:

$$\frac{\text{Final body weight} - \text{Initial body weight}}{\text{Final body weight}} \times 100$$

### 2.6 Calculation of Percentage Prostate Increase Inhibition (PPII)

$$\text{Percentage prostate increase inhibition} = 100 - \left[ \frac{(T - C)}{(B - C)} \times 100 \right]$$

Where C, B, and T are the prostate weights of the normal control, BPH control, and treatment group [23].

### 2.7 Biochemical Tests

Prostate Specific Antigen (PSA) level was measured using AccuBind® Immunoassay (TYPE 3) test kit.

Prolactin level was measured using AccuBind®, Immunoassay (TYPE 4) test kit. Testosterone level was measured using AccuBind®, Competitive Enzyme Immunoassay (TYPE 7) test kit. DHT was measured using enzyme-linked immunosorbent assay (ELISA) kits.

### 2.8 Statistical Analysis

Data were analyzed based on one-way analysis of variance (ANOVA) with Statistical Product and Service Solutions (SPSS) version 23. Results are expressed as mean ± SD and presented in Tables. Differences between means were considered to be significant at  $P = .05$  using Duncan multiple range test (DMRT).

## 3. RESULTS AND DISCUSSION

The body weights of experimental animals determined weekly, is usually used as a general index of overall health [24]. For the body weight (Table 2), the results indicated increase in the body weights of all the groups in week 1. It was noticed that there was a reduction in the body weight of albino rats in Group B (BPH control), and the high dose groups H (20ml /kg AC) and I (20 ml/kg AMAC) in week 2. Further reduction in the body weights of the experimental animals in these groups and Group G (20 ml/kg AC-high dose) was observed in week 3. While in week 4, an increase in the body weights of the experimental animals in all the groups was observed, except for Group G where there was further decrease in body weight. This indicated that at the low dose (10 ml/kg b.w) of the test

**Table 2. Effect of *Annona muricata* pulp and *Allium cepa* bulb juices on body weight of albino rats with testosterone and oestradiol-induced BPH**

Group	Body weight per week (g)				
	Initial weight	Week 1	Week 2	Week 3	Final weight (Wk 4)
NC	177.13 ± 19.13	196.31 ± 22.65 (9.70 ± 3.30 <sup>c</sup> %)	218.15 ± 26.18 (9.98 ± 1.18 <sup>b</sup> %)	235.18 ± 31.03 (7.13 ± 2.19 <sup>d</sup> %)	244.60 ± 47.56 (3.10 ± 6.47 <sup>ab</sup> %)
BPH	198.56 ± 18.26	200.04 ± 17.03 (0.78 ± 0.69 <sup>a</sup> %)	197.92 ± 13.50 (-1.00 ± 1.75 <sup>a</sup> %)	189.68 ± 20.61 (-4.67 ± 5.22 <sup>a</sup> %)	199.84 ± 15.48 (5.23 ± 3.28 <sup>b</sup> %)
SC	159.37 ± 1.98	166.52 ± 3.71 (4.24 ± 3.33 <sup>abc</sup> %)	167.23 ± 8.40 (0.18 ± 7.39 <sup>a</sup> %)	175.54 ± 8.38 (4.73 ± 1.59 <sup>cd</sup> %)	185.41 ± 9.70 (5.30 ± 1.51 <sup>b</sup> %)
10 ml/kg AM	148.47 ± 6.80	158.13 ± 5.21 (6.13 ± 1.67 <sup>abc</sup> %)	162.93 ± 3.20 (2.96 ± 1.94 <sup>a</sup> %)	168.55 ± 2.42 (3.32 ± 2.42 <sup>bcd</sup> %)	176.42 ± 4.99 (4.40 ± 3.41 <sup>ab</sup> %)
10 ml/kg AC	140.26 ± 17.45	153.75 ± 17.52 (8.85 ± 1.04 <sup>bc</sup> %)	161.30 ± 16.61 (4.75 ± 1.16 <sup>ab</sup> %)	162.23 ± 18.73 (0.47 ± 1.56 <sup>abcd</sup> %)	165.46 ± 20.21 (1.89 ± 0.77 <sup>ab</sup> %)
10 ml/kg AMAC	151.68 ± 13.57	159.98 ± 19.32 (4.88 ± 5.22 <sup>abc</sup> %)	167.43 ± 13.63 (4.62 ± 4.74 <sup>ab</sup> %)	170.65 ± 11.97 (1.92 ± 2.70 <sup>abcd</sup> %)	176.06 ± 15.92 (2.93 ± 3.33 <sup>ab</sup> %)
20 ml/kg AM	163.18 ± 9.27	171.36 ± 5.73 (4.83 ± 2.27 <sup>abc</sup> %)	172.70 ± 7.93 (0.73 ± 1.27 <sup>a</sup> %)	168.80 ± 10.42 (-2.44 ± 4.70 <sup>ab</sup> %)	165.88 ± 16.48 (-2.06 ± 5.18 <sup>a</sup> %)
20 ml/kg AC	179.80 ± 26.23	188.88 ± 22.33 (4.99 ± 3.05 <sup>abc</sup> %)	185.65 ± 14.00 (-1.54 ± 4.71 <sup>a</sup> %)	180.57 ± 3.00 (-2.75 ± 6.01 <sup>ab</sup> %)	189.69 ± 9.98 (4.69 ± 3.41 <sup>ab</sup> %)
20 ml/kg AMAC	177.95 ± 17.18	184.29 ± 12.48 (3.54 ± 3.54 <sup>ab</sup> %)	182.58 ± 13.84 (-0.98 ± 1.27 <sup>a</sup> %)	182.13 ± 10.75 (-0.18 ± 1.86 <sup>abc</sup> %)	187.81 ± 7.26 (3.07 ± 2.54 <sup>ab</sup> %)

Values are means ± standard deviations of triplicate determinations. Values on the same column bearing different superscript letters are significantly different ( $p < .05$ )

samples; soursop juice, onion juice, and combination of soursop and onion juices, the body weights of the animal models increased steadily throughout the weeks of the study, in the same manner observed in the normal and standard controls (Groups A and C, respectively). The weight gain in the affected groups throughout the weeks can be easily noticed with the positive values of percentage change in body weights per week. Hence, at low doses, the samples maintained the body weights of the albino rats.

In Table 3, the results showed that the PWs reduced in all test groups. Lower values were observed in the groups treated with low doses of the test samples, meaning that increasing the dosage of the test samples does not reduce the PWs any further. The results of PPII indicated that the low dose onion juice (10 ml/kg AC) inhibited prostate increase better among the test groups, followed by low dose soursop juice (10

ml/kg AM). Except for the PPII value of the high dose soursop juice (20 ml/kg AM), all the PPII values from the test groups were not significantly different from the standard control. The results of this study agreed with the findings of Ogbu *et al.* [25], which stated that actogenin-rich fraction of *Annona muricata* leaves extract reduced PW in treated rats.

Table 4 shows the results of the effect of juice samples of *Annona muricata* pulp and *Allium cepa* bulb on serum DHT, testosterone, prolactin and PSA levels of BPH induced albino rats. DHT and testosterone are androgens crucial for the growth and development of the internal organs of the human male reproductive system, and BPH [26][27][28]. The 5- $\alpha$  reductase enzyme catalyzes the conversion of testosterone to DHT. Due to its high affinity to the androgen receptors, DHT is the most potent androgen that promotes the growth of the epithelial and stromal cells of the prostate [8][29][30].

**Table 3. Effect of juices of *Annona muricata* pulp and *Allium cepa* bulb on prostate weight and percentage prostate increase inhibition of albino rats with testosterone and oestradiol-induced BPH**

Group	Prostate weight (g)	Percentage prostate increase inhibition (%)
NC	2.18 $\pm$ 0.26 <sup>a</sup>	-
BPH	3.31 $\pm$ 0.35 <sup>c</sup>	-
SC	2.53 $\pm$ 0.21 <sup>ab</sup>	69.32 $\pm$ 40.15 <sup>b</sup>
10mg/kg AM	2.85 $\pm$ 0.66 <sup>bc</sup>	41.00 $\pm$ 35.33 <sup>ab</sup>
10mg/kg AC	2.77 $\pm$ 0.13 <sup>abc</sup>	48.08 $\pm$ 24.44 <sup>ab</sup>
10mg/kg AMAC	2.98 $\pm$ 0.07 <sup>bc</sup>	29.79 $\pm$ 23.15 <sup>ab</sup>
20mg/kg AM	3.25 $\pm$ 0.22 <sup>c</sup>	5.61 $\pm$ 28.81 <sup>a</sup>
20mg/kg AC	2.95 $\pm$ 0.58 <sup>bc</sup>	32.45 $\pm$ 34.25 <sup>ab</sup>
20mg/kg AMAC	3.00 $\pm$ 0.07 <sup>bc</sup>	27.73 $\pm$ 24.50 <sup>ab</sup>

Values are means  $\pm$  standard deviations of triplicate determinations. Values on the same column bearing different superscript letters are significantly different ( $p < .05$ )

**Table 4. Effect of *Annona muricata* pulp and *Allium cepa* bulb juices on serum levels of dihydrotestosterone, testosterone, prolactin and PSA of albino rats with testosterone and oestradiol-induced BPH**

Group	DHT (ng/ml)	Testosterone (ng/ml)	Prolactin (ng/ml)	PSA (ng/ml)
NC	8.02 $\pm$ 0.20 <sup>bc</sup>	2.30 $\pm$ 0.40 <sup>a</sup>	1.00 $\pm$ 0.30 <sup>a</sup>	0.20 $\pm$ 0.00 <sup>ab</sup>
BPH	10.50 $\pm$ 0.60 <sup>d</sup>	10.20 $\pm$ 0.70 <sup>f</sup>	1.80 $\pm$ 0.10 <sup>b</sup>	0.40 $\pm$ 0.10 <sup>c</sup>
SC	8.70 $\pm$ 0.50 <sup>c</sup>	6.40 $\pm$ 0.20 <sup>b</sup>	1.00 $\pm$ 0.10 <sup>a</sup>	0.40 $\pm$ 0.20 <sup>c</sup>
10mg/kg AM	9.10 $\pm$ 0.10 <sup>c</sup>	9.80 $\pm$ 0.50 <sup>f</sup>	0.90 $\pm$ 0.30 <sup>a</sup>	0.30 $\pm$ 0.10 <sup>bc</sup>
10mg/kg AC	8.10 $\pm$ 0.80 <sup>bc</sup>	7.30 $\pm$ 0.60 <sup>c</sup>	1.10 $\pm$ 0.10 <sup>a</sup>	0.30 $\pm$ 0.00 <sup>bc</sup>
10mg/kg AMAC	7.10 $\pm$ 1.20 <sup>b</sup>	8.30 $\pm$ 0.20 <sup>de</sup>	1.00 $\pm$ 0.30 <sup>a</sup>	0.40 $\pm$ 0.10 <sup>c</sup>
20mg/kg AM	5.30 $\pm$ 0.80 <sup>a</sup>	9.60 $\pm$ 0.10 <sup>f</sup>	0.90 $\pm$ 0.00 <sup>a</sup>	0.20 $\pm$ 0.10 <sup>ab</sup>
20mg/kg AC	7.20 $\pm$ 0.30 <sup>b</sup>	8.70 $\pm$ 0.00 <sup>e</sup>	1.00 $\pm$ 0.01 <sup>a</sup>	0.10 $\pm$ 0.00 <sup>a</sup>
20mg/kg AMAC	8.40 $\pm$ 0.20 <sup>c</sup>	7.90 $\pm$ 0.10 <sup>cd</sup>	0.80 $\pm$ 0.20 <sup>a</sup>	0.20 $\pm$ 0.10 <sup>ab</sup>

Values are means  $\pm$  standard deviations of triplicate determinations. Values on the same column bearing different superscript letters are significantly different ( $p < .05$ )

PSA is predominantly produced in the prostate gland. Benign prostatic hyperplasia is an age-related condition, hence, its occurrence increases with increase in age, as the prostate volume increases with age. This increase in prostate volume with age leads to increase in PSA levels [31]. Although, PSA values greater than 4.0 ng/ml is considered abnormal in men above 60 years of age, yearly increase in PSA levels greater than 0.35 ng/ml is alarming and requires further testing [32]. Prolactin has also been implicated in development of BPH, as it targets the prostate through interactions with the prolactin receptors expressed in the prostate [33]. Chronic hyperprolactinemia induces enlargement and inflammation of the lateral rat prostate [34]. High estrogen and prolactin levels also increase DHT levels and stimulate growth of prostate cells [9].

From the results of this study, a reduction in DHT and testosterone levels were observed in all test groups, same as the standard control. It was also noted that the individual samples caused a reduction DHT levels with increase in concentration, while the soursop juice and combination samples decreased testosterone levels with increase in concentration. The results for serum prolactin levels showed a significant ( $p < .05$ ) reduction in both the test groups and standard control when compared with the disease control. While, there was also an observed decrease in serum PSA levels in the test groups, which also occurred in a dose-dependent manner, however, all the values obtained for both tests and controls did not vary significantly ( $p > .05$ ). Investigations carried out by Blessing [2], and Ebenyi *et al.* [35] also observed a dose-dependent decrease in PSA level, and PSA, DHT and testosterone levels, respectively. The increased values obtained in the BPH control further reiterated the discoveries of other researchers. The ability of soursop and onions in ameliorating BPH as observed in this study agreed with the findings of Hong *et al.* [36] on stem bark extracts of soursop and Elberry *et al.* [37] on red onions scales extract.

#### 4. CONCLUSION

The results of this study suggested that soursop pulp and onion bulb juice exerts ameliorative effects on BPH. Better effects were achieved with the use of single samples and at the low dose studied.

#### ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

#### ACKNOWLEDGEMENTS

The Tertiary Education Trust Fund (TETFund, TEFT/DASTD/POLY/IMO STAFF/TSAS/2020/VOL.1) of the Federal Government of Nigeria sponsored the Ph.D programme that birthed this research work.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. McVary KT. Patient education: benign prostatic hyperplasia (BPH) (beyond the basics). In: O'Leary MP, Givens J, Editors; 2020. Accessed 1 May 2023. Available: <https://www.uptodate.com/contents/benign-prostatic-hyperplasia-bph-beyond-the-basics/print>.
2. Blessing AN. Effect of ethanol leaf extract of *Gnetum africanum* on testosterone and oestradiol induced benign prostatic hyperplasia. *J Med Plant Res.* 2020;14(7):309-16. DOI:10.5897/JMPR2019.6901
3. Emberton M, Cornel EB, Bassi PF, Fourcade RO, Gómez JMF, Castro R. Benign prostatic hyperplasia as a progressive disease: a guide to the risk factors and options for medical management. *International Journal of Clinical Practice.* 2008;62(7):1076-86. DOI:10.1111/j.1742-1241.2008.01785.x. PMID: 18479366
4. Lukacz ES, Sampsel C, Gray M, MacDiarmid S, Rosenberg M, Ellsworth P, et al. A healthy bladder: A consensus statement. *International Journal of Clinical Practice.* 2011;65(10):1026-36. DOI:10.1111/j.1742-1241.2011.02763.x PMID: 21923844

5. Lepor H. Pathophysiology of benign prostatic hyperplasia in the aging male population. *Rev Urol.* 2005;7(4):S3-S12.PMID: 16986052.
6. Berger AP, Deibl M, Leonhartsberger N, Bektic J, Fritsche G, Steiner H, et al. Vascular damage as a risk factor for benign prostatic hyperplasia and erectile dysfunction. *BJU Int.* 2005;96(7):1073-8. DOI:10.1111/j.1464-410X.2005.0577.x. PMID: 16225531
7. Zhang SJ, Quian HN, Zhao Y, Sun K, Wang H-Q, Liang G-Q, et al. Relationship between age and prostate size. *Asian J Androl.* 2013;15(1):116-20. DOI:10.1038/aja.2012.127 PMID: 23223031
8. Carson III C, Rittmaster R. The role of dihydrotestosterone in benign prostatic hyperplasia. *Urol.* 2003;61(4):2-7. DOI:10.1016/s0090-4295(3)00045-1 PMID: 12657354
9. Cooper J. Symptoms, causes, natural treatment. *Benign prostatic hyperplasia (BPH)- enlarged prostate; 2019.* Accessed 30 April 2023. Available:<https://advancedfunctionalmedicine.com.au/benign-prostatic-hyperplasia-bph-enlarged-prostate/>
10. Park HJ, Won JE, Sorsaburu S, Rivera PD, Lee SW. Urinary tract symptoms (LUTS) secondary to benign prostatic hyperplasia (BPH) and LUTS/BPH with erectile dysfunction in Asian men: a systematic review focusing on Tadalafil. *World J Mens Health.* 2013;31(3):193-207. DOI:10.5534/wjmh.2013.31.3.193 PMID: 24459652
11. Calixto JB. Efficacy, safety, quality control, marketing and regulatory guidelines for herbal medicines (phytotherapeutic agents). *Braz J Med Biol Res.* 2000;33(2):179-89. DOI:10.1590/s0100-879x2000000200004. PMID: 10657057
12. Thompson IM, Goodman PJ, Tangen CM, Lucia MS, Miller GJ, Ford LG, et al. The influence of finasteride on the development of prostate cancer. *N Engl J Med.* 2003;349(3):215-24. DOI:10.1056/NEJMoa030660. PMID: 12824459.
13. Rodrigues AS, Almeida DPF, Simal-Gandara J, Perez-Gregorio MR. Onions: a source of flavonoids. *Flavonoids; 2017.* Accessed 1 May 2023 Available:<https://doi.org/10.5772/intechopen.69896>
14. Griffiths G, Trueman L, Crowther T, Thomas B. Onions -- a global benefit to health. *Phytother Res.* 2002;16(7):603-15. DOI:10.1002/ptr.1222 PMID: 12410539.
15. Mishra S, Ahmad S, Kumar N, Sharma B. *Annona muricata* (the cancer killer): A review. *Glob. J Pharm Res.* 2013;2:1613-8.
16. Adewole SO, Caxton-Martins EA. Morphological changes and hypoglycemic effects of *Annona muricata* Linn. (Annonaceae) leaf aqueous extract on pancreatic B-cells of streptozotocin-treated diabetic rats. *Afr J Tradit Complement Altern Med.* 2009;9:173-87.
17. Ribeiro de Souza EB, da Silva RR, Afonso S, Scarminio IS. Enhanced extraction yields and mobile phase separations by solvent mixtures for the analysis of metabolites in *Annona muricata* L. leaves. *J Sep Sci.* 2009;32(23-24):4176-85. DOI:10.1002/jssc.200900375 PMID: 19882621
18. Gyesei JN, Opoku R, Borquaye LS. Chemical composition, total phenolic content, and antioxidant activities of the essential oils of the leaves and fruit pulp of *Annona muricata* L. (Soursop) from Ghana. *Biochem Res Int.* 2019;4164576.
19. Animal Research Review Panel. ARRPP Guideline 20: Guidelines for the housing of rats in scientific institutions. 2007;16-56. Accessed 13 April 2022. Available:[https://www.animaethics.org.au/PDFGuidelines for the Housing of Rats in Scientific Institutions.](https://www.animaethics.org.au/PDFGuidelines%20for%20the%20Housing%20of%20Rats%20in%20Scientific%20Institutions.pdf)
20. Mbaka G, Anunobi C, Ogunsina S, Osiagwu D. Histomorphological changes in induced benign prostatic hyperplasia with exogenous testosterone and estradiol in adult male rats treated with aqueous ethanol extract of *Secamone afzelii*. *Egyptian Journal of Basic and Applied Sciences.* 2017;4:15-21. Available:<https://doi.org/10.1016/j.elbas.2016.11.003>
21. Akanni OO, Owumi SE, Olowofela OG, Adeyanju AA, Abiola OJ, Adaramonye OA. Protocatechuic acid ameliorates testosterone-induced benign prostatic hyperplasia through the regulation of



- inflammation and oxidative stress in castrated rats. *J Biochem Mol Toxicol.* 2020;34(8):e22502.  
DOI:10.1002/jbt.22502. PMID: 32227675
22. Majeed M, Bani S, Natarajan S, Pandey A, Naveed S. Evaluation of 90 day repeated dose oral toxicity and reproductive/developmental toxicity of 3'-hydroxypterostilbene in experimental animals. *PLoS One.* 2017;12(3):e0172770.  
DOI:10.1371/journal.pone.0172770. PMID: 28257483
  23. Sasidharan S, KP S, Bhaumik A, Kanti Das S, Nair JH. Administration of *Caesalpinia bonduc* seed extracts ameliorates testosterone-induced Benign Prostatic Hyperplasia (BPH) in male wistar rats. *Res Rep Urol.* 2022;14:225-39.  
DOI:10.2147/RRU.S365578. PMID: 35651598
  24. Brai BIC, Odetola AA, Agomo PU. Effects of *Persea americana* leaf extracts on body weight and liver lipids in rats fed hyperlipidaemic diet. *Afri J Biotechnol.* 2007;6(8):1007-11.
  25. Ogbu PN, Ugota EO, Onwuka RU, Ogbu IM, Alope C. Effect of acetogenin fraction of *Annona muricata* leaves on antioxidant status and some indices of benign prostatic hyperplasia in rats. *Redox Rep.* 2020;25(1):80-6.  
DOI:10.1080/13510002.2020.1804711. PMID: 32878595
  26. Andriole G, Bruchofsky N, Chung LW, Matsumoto AM, Rittmaster R, Roehrborn C, et al. Dihydrotestosterone and the prostate: The scientific rationale for 5alpha-reductase inhibitors in the treatment of benign prostatic hyperplasia. *J Urol.* 2004;172:1399-403.  
DOI:10.1097/01.ju.0000139539.94828.29. PMID: 15371854
  27. Lucia MS, Lambert JR. Growth factors in benign prostatic hyperplasia: Basic science implications. *Curr Urol Rep.* 2008;9(4):272-8.  
DOI:10.1007/s11934-008-0048-6. PMID: 18765125
  28. Miller J, Tarter TH. Combination therapy with dutasteride and tamsulosin for the treatment of symptomatic enlarged prostate. *Clin Interv Aging.* 2009;4:251-8.  
DOI:10.2147/cia.s4102. PMID: 19554096
  29. Mizokami A, Koh E, Izumi K, Narimoto K, Takeda M, Honma S, et al. Prostate cancer stromal cells and LNCaP cells coordinately activate the androgen receptor through synthesis of testosterone and dihydrotestosterone from dehydroepiandrosterone. *Endocr-Relat Cancer.* 2009;16(4):1139-55.  
DOI:10.1677/ERC-09-0070. PMID: 19608712.
  30. Nicholson TM, Ricke WA. Androgens and estrogens in benign prostatic hyperplasia: Past, present and future. *Differentiation.* 2011;82:184-99.  
DOI:10.1016/j.diff.2011.04.006. PMID: 21620560
  31. Putra IB, Hamid AR, Mochtar, CA, Umbas R. Relationship of age, prostate-specific antigen, and prostate volume in Indonesian men with benign prostatic hyperplasia. *Prostate Int.* 2016;4(2):43-8.  
DOI:10.1016/j.pmil.2016.03.002. PMID: 27358842
  32. Pavlovich CP. Prostate cancer: age-specific screening guidelines. *John Hopkins Medicine.* 2023. Accessed 1 May 2023.  
Available: <https://www.hopkinsmedicine.org/health/conditions-and-diseases/prostate-cancer/prostate-cancer-age-specific-screening-guidelines#:~:text=The%20normal%20range%20is%20between,doctor%20may%20recommend%20further%20testing.>
  33. La Vignera S, Condorelli RA, Russo GI, Morgia G, Calogero AE. Endocrine control of benign prostatic hyperplasia. *Androl.* 2016;4(3):404-11.  
DOI:10.1111/andr.12186 PMID: 27089546
  34. Van Coppenolle F, Slomianny C, Carpentier F, Le Bourhis X, Ahidouch A, Croix D, et al. Effects of hyperprolactinemia on rat prostate growth: Evidence of androgen-dependence. *Am J Physiol Endocrinol Metab.* 2001;280(1):E120-9.  
DOI:10.1152/ajpendo.2001.280.1. PMID: 11120666
  35. Ebenyi LN, Ominyi MC, Anyanwu CB, Ogah O, Ogbanshi ME. The leaf extract of *Carica papaya* alleviates benign prostate hyperplasia in male albino rats by antioxidative mechanisms. *Nigerian Journal of Biochemistry and*

- Molecular Biology. 2022;37(4): 298-302. DOI:10.1155/2014/640746 PMID: 24829522.  
Available:<https://doi.org/10.2659/njmb.136>
36. Elberry AA, Mufti S, Al-Maghrabi J, Sattar EA, Ghareib SA, Mosli HA, et al. Immunomodulatory effect of red onion (*Allium cepa* Linn) scale extract on experimentally induced atypical prostatic hyperplasia in Wistar rats. *Mediators Inflamm.* 2014;2014:640746.
37. Hong GU, Choi M, Chung M-H, Ro JY. Anti-inflammatory effects of graviola stem bark extracts on the testosterone-induced benign prostatic hyperplasia model in pats. *Food Supplements and Biomaterials for Health.* 2021;1(2):e18. DOI:10.52361/fsbh.2021.1.e18

© 2023 Ibegbulem et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/107355>