



# **Yield Enhancement of Field Pea (*Pisum sativum*) through Zinc Solubilizing Microorganism**

**Shraddha Somkuwar <sup>a</sup>, Rahul Aske <sup>b\*</sup>, Arvind Ahirwal <sup>b</sup>,  
Deepak Kher <sup>b</sup> and Smriti Chauhan <sup>a</sup>**

<sup>a</sup> Department of Microbiology, Barkatullah University Bhopal, (M.P.), India.

<sup>b</sup> School of Agriculture, Sanjeev Agrawal Global Educational (SAGE) University Bhopal (M.P.), India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/CJAST/2023/v42i404258

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108043>

**Original Research Article**

**Received: 16/08/2023**

**Accepted: 22/10/2023**

**Published: 01/11/2023**

## **ABSTRACT**

In the rabi season of 2022-23, a field experiment was conducted at Instructional Farm, School of Agriculture, Sanjeev Agrawal Global Educational (SAGE) University Bhopal (Madhya Pradesh) to assess the impact of zinc (zn) and ZSB (Zinc Solubilizing Bacteria) solubilizers on the performance of field pea crops. The results of the experiment revealed that the application of 100% of the recommended dose of fertilizer in combination with zinc led to the highest values in various growth and yield parameters.

Specifically, the treatment of 100% Recommended dose fertilizer + Zinc statistically ( $p < 0.05$ ) compared to mean values demonstrated the maximum plant height, dry matter accumulation (in grams), branches per plant, number of pods per plant, number of seeds per pod and seed yield (1060 kg/ha). Additionally, it resulted in a straw of 2130 kg/ha., a biological yield of 4243 kg/ha, a net return of 1639 Rs/ha, and a benefit-to-cost (B: C) ratio of 1.12. It is worth noting that the

\*Corresponding author: E-mail: rahulaske@gmail.com;

performance achieved with the application of 100% ZSB was statistically on par with the treatment involving 100% Recommended Dose Fertilizer + Zinc. Overall, all of the treatments applied in the experiment had a significant positive impact on the growth, yield, and economic aspects of field pea cultivation in comparison to the control group.

**Keywords:** Field pea; Zinc; ZSB (Zinc solubilizing bacteria); growth; yield.

## 1. INTRODUCTION

Field pea (*Pisum sativum*), a primary winter grain legume crop, primarily thrives in cooler temperate regions. This versatile crop serves various purposes, including providing food, animal feed, and vegetables. It ranks as the third most popular rabi crop globally, following chickpea and arhar, making it a significant pulse crop.

In order to enhance both the production and productivity of field peas, there's a pressing need for improved nutrient management practices. Field pea has a positive response to the application of fertilizers. Nevertheless, the constituent use of chemical fertilizer in modern agriculture practices has adverse consequences for the population of beneficial native soil microorganisms and overall soil health. An alternative to chemical fertilizer is the use of bio-fertilizers, which are organic products containing living cells. Bio-fertilizer plays a crucial role in converting nutrients from unavailable forms to available ones, as indicated by Itelima et al and Ikeh et al. [1-9].

These organic solutions encompass microorganisms and growth hormones which boost crop productivity through the process of biological nitrogen fixation, zinc solubilization, and the regulation of growth factors essential for plants' growth and development.

Biofertilizer is more sustainable in managing soil fertility [10]. In this context, the combination of nutrient management and bio-fertilizers is emerging as a commercially viable and environmentally responsible approach to fertilization, as suggested by Kaur and Purewal, [2] and Ikeh et al. [11].

## 2. MATERIALS AND METHODS

The current investigation was conducted in small experimental plots at SAGE University, Bhopal (Madhya Pradesh). The study aimed to assess the impact of zinc (Zn) and zinc solubilizers on the growth and development of field peas, with a focus on the selected variety, Prakash IPFD 1-1.

This particular variety is known for its large seeds and resistance to powdery mildew. Its distinct yellow ovoid grains are noteworthy for their exceptionally high protein content. The sowing of field peas took place on November 23, 2022, in the rabi season of 2022-2023.

The experimental design followed a randomized block format (RBD), with six different treatments. Each plot had an area of 6 square meters. The experimentation was carried out at the Instructional Cum Research Farm, School of Agriculture, SAGE University, Bhopal, situated in the central region of India, characterized by a latitude of 23.2599° N and a longitude of 77.4126° E. Bhopal (often referred to as the city of lakes) is nestled in the lowlands near the Vindhya mountain range on the Malwa plateau.

In four of the treatments, the recommended doses of nitrogen, phosphorus, and potassium (16.6g at the basal dose) were applied, with ZSB (Zinc Solubilizing Bacteria) being a common component in all treatments. The sowing date was November 23, 2022, and manual harvesting was conducted on March 24, 2023.

Data collections were taken from each randomly selected plot at 30, 60, and 90, days after planting (DAS) and at harvesting. As well as during the harvest of the crop. Throughout the growth period of the field pea, various observations were systematically collected from each plot, with proper tagging for future reference. The observations encompassed key growth and yield attributes of the field pea, including plant height (in centimetres), the number of branches per plant, the number of pods per plant, the number of seeds per pod, and seed yield. For plant height, number of branches per plant, number of pods per plant, and number of seeds per pod, measurements were taken from five randomly selected plants, and the mean value was computed. The weight of 1,000 seeds was recorded in grams. The seed, straw, and biological yield were measured in kilograms per hectare, and the harvest index was calculated. Net returns and the benefit-to-

cost (B:C) ratio were computed to evaluate the economic aspects of the field experiment.

### 3. RESULTS AND DISCUSSION

Plant height observations were made at 60 and 90 days after sowing (DAS). The results indicated that plant height was at its minimum up to 30 DAS but subsequently increased during the 60 DAS mark and at harvest, as the crop aged. Plant height varied between 65.20 cm and 84.34 cm at harvest. The treatment of the recommended dose fertilizer + 100% Zn (9 gm.), denoted as T2, significantly exhibited the highest plant height, reaching 84.34 cm at harvest, surpassing all other treatments.

Regarding dry matter accumulation (g/plant) at harvest, the application of the recommended dose of fertilizer + 100% Zn (19.68 gm.) resulted in significantly higher values compared to the other treatments (Table 1). The data (Table 2) demonstrates that the application of Zn and ZSB (Zinc Solubilizing Bacteria) levels significantly increased the number of branches per plant, pods per plant, and seeds per pod. The treatment of the recommended dose of fertilizer + 100% Zn led to the maximum branches per plant (2.46), pods per plant (16.61), and seeds per pod (6.85) among all the treatments.

The results showed that the application of the recommended dose of fertilizer (RDF) + 100%

Zn (T2) significantly increased seed, straw, and biological yield compared to the control. The maximum seed yield (1060 kg/ha), straw yield (2130 kg/ha), and biological yield (4243 kg/ha) were obtained with the application of the recommended dose fertilizer + 100% Zn (Table 2). The treatment of the recommended dose fertilizer + 100% Zn was statistically superior, especially when used in conjunction with Zinc Solubilizing Bacteria (ZSB) at 100%, possibly due to its positive impact on the number of pods per plant, seeds per pod, seed, straw, and biological yield. Economically, the data (Table 2) revealed that among the Zn and ZSB Solubilizer fertilizer treatments, the recommended dose of fertilizer + 100% Zinc resulted in the highest and statistically significant net return (1639 Rs/ha) and benefit-to-cost (B: C) ratio (1.12). Similar findings were in line with those reported by Venkatrao et al. [8] and Kuniya et al. [4]. The beneficial effects of zinc-solubilizing bacteria on yield attributes were also consistent with the findings of Kothyari et al. [3] and Raut [5]. Notably, test weight did not show significant differences concerning various Zn fertilizers and their solubilizer levels. Economic analysis (Table 2) indicated that, among the Zn fertilizer treatments, the recommended dose of fertilizer + 100% Zinc yielded the highest and statistically significant net return (1639 Rs/ha) and B: C ratio (1.12), which aligns with the results reported by Serawat et al. [6] and Singh et al. [7].




Fig 1. Graphical representation of the treatments

**Table 1. Impact of Zinc solubilizing microorganisms on plan growth of field pea**

Treatments	At harvesting time plant Height in cm.	Dry matter Accumulation (gram per plant) at harvest	Number of branches per plant at harvest	Number of pods per plant at harvest	Seed per pods at harvest	Test weight (1000 seed)(g)
<b>Zn and ZSB Levels</b>						
control (T <sub>1</sub> )	65.20	13.12	1.04	9.94	4.42	175.36
Recommended Dose Fertilizer + Zn (100%) (T <sub>2</sub> )	84.34	19.68	2.46	16.61	6.85	189.54
Zinc solubilizing bacteria (ZSB) (100%) (T <sub>3</sub> )	80.53	19.61	2.15	15.53	6.24	185.62
Recommended dose fertilizer + Zn +ZSB (75%) (T <sub>4</sub> )	75.20	17.82	1.78	13.78	5.78	182.41
Recommended dose fertilizer + Zinc +ZSB (50%)	71.48	15.7	1.70	12.63	5.23	179.45
Recommended dose fertilizer + Zinc +ZSB (25%) (T <sub>6</sub> )	69.91	14.9	1.25	11.5	4.5	176.69
<b>SE(m)</b>	1.52	0.46	0.07	0.31	0.18	3.84
<b>CD at 5%</b>	4.65	1.34	0.23	0.95	0.57	NS

**Table 2. Impact of zinc solubilising microorganism on seed yield, straw yield, biological yield and Harvest index**

Treatments	Seed yield (kg. per hectare)	Straw Yield (kg. per hectare)	Biological yield (kg. per hectare)	Harvest index (%)	Net return (Rs./ha)	B:C ratio
<b>Zn and ZSB Levels</b>						
Control (T <sub>1</sub> )	250	560	1005	30.06	820	0.52
Recommended Dose fertilizer + Zn (100%) (T <sub>2</sub> )	1060	2130	4243	32.26	1639	1.12
Zinc solubilizing bacteria (ZSB) (100%) (T <sub>3</sub> )	830	1655	3298	32.09	1460	1.03
Recommended dose fertilizer + Zn +ZSB (75%) (T <sub>4</sub> )	700	1390	2680	31.45	1234	1.01
Recommended dose fertilizer + Zn +ZSB (50%) (T <sub>5</sub> )	480	940	1860	31.15	1140	0.95
Recommended dose fertilizer + Zn +ZSB (25%) (T <sub>6</sub> )	380	735	1438	30.62	960	0.76
<b>SE(m)</b>	20.4	40.7	95.3	0.29	161.34	0.02
<b>CD at 5%</b>	61.1	120.3	285.4	0.86	484.05	0.07

#### 4. CONCLUSION

In conclusion, the study conducted at SAGE University, Bhopal, focusing on the impact of zinc (Zn) and Zinc Solubilising Bacteria (ZSB) on the growth and yield attributing character of field pea, revealed the data of plant height, observed at 60 and 90 days after sowing (DAS), exhibited an interesting growth pattern, minimum height was recorded at 30 DAS, followed by a steady and gradually increase up to 60 DAS and at the harvest stage. The treatment involving the recommended dose of fertilizer + 100% Zn (9 gm.) outshone all others, reaching a significant plant height of 84.34 cm at harvest.

Furthermore, dry matter accumulation (g/plant) at harvest was significantly higher when the recommended dose of fertilizer was combined with 100% Zn (19.68 gm.), outperforming the other treatments. The application of Zn and ZSB also had a substantial impact on the number of branches per plant, pods per plant, and seeds per pod. Once again, the treatment of the recommended dose of fertilizer + 100% Zn excelled, resulting in the highest number of branches per plant (2.46), pods per plant (16.61), and seeds per pod (6.85).

Importantly, the application of the recommended dose of fertilizer (RDF) combined with 100% Zn (T2) significantly increased seed, straw, and biological yields in comparison to the control. Maximum seed yield (1060 kg/ha), straw yield (2130 kg/ha), and biological yield (4243 kg/ha) were achieved with this treatment. The positive impact of combining the recommended dose fertilizer with 100% Zn was further enhanced when used in conjunction with Zinc Solubilizing Bacteria (ZSB) at 100%, contributing to the impressive results across parameters.

Economically, the study demonstrated that the treatment involving the recommended dose of fertilizer + 100% Zn yielded the highest net return (1639 Rs/ha) and a significant benefit-to-cost (B:C) ratio of 1.12 among the Zn and ZSB Solubilizer fertilizer treatments. These findings align with prior research, reinforcing the economic and agricultural advantages of this approach.

Notably, test weight did not show significant variations among the different Zn fertilizers and solubilizer levels. In summary, the study underscores the effectiveness of the recommended dose of fertilizer in combination

with 100% Zn, particularly when supplemented with ZSB, in enhancing the growth, yield, and economic viability of field pea cultivation, affirming the potential for sustainable and economically sound agricultural practices.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Itelima JU, Bang WJ, Onyimba IA, Egbere OJ. A review: Biofertilizer; A key player in enhancing soil fertility and crop productivity. *Journal of Microbiology and Biotechnology Report*. 2018;2(1):22-28.
2. Kaur P, Purewal SS. Biofertilizers and their role in sustainable agriculture. *Biofertilizers for Sustainable Agriculture and Environment Soil Biology*. 2019;55:285-300.
3. Kothyari HS, Yadav LK, Jat R, Gurjar PC. Influence of Biofertilizers on Plant Growth and Seed Yield of Pea (*Pisumsativum* [L.]). *International Journal Current Microbiology and Applied Science*. 2017;6(11):1810-1817.
4. Kuniya N, Chaudhary N, Patel S. Effect of sulphur and zinc application on growth, yield attributes, yield and quality of summer clusterbean (*Cyamopsistetragonoloba* [L.]) in light textured soil. *International Journal of Chemical Studies*. 2018;6(1):1529-1532.
5. Raut AD, Durgude AG, Kadlag AD. Effect of zinc solubilizing bacteria on zinc use efficiency and yield of summer groundnut grown in Entisol. *International Journal of Chemical Studies*. 2019;7(1):1710-1713.
6. Serawat A, Sharma Y, Serawat M, Kapoor A, Jakhar RK. Effect of Phosphorus and Sulphur on Growth attributes and Yield of mung bean (*Vignaradiata* [L.]). *International Journal of Current Microbiology and Applied Sciences*. 2018;7(8):2674-2683.
7. Singh N, Joshi E, Sasode DS, Sikarwar RS, Rawat GS. Liquid biofertilizer and inorganic nutrients effect on physiological, quality parameters and productivity of kharif groundnut (*Arachishypogaea* [L.]) *International journal Current Microbiology and Applied Science*. 2018;7(09):729-735.

8. Venkatarao CV, Naga SR, Yadav BL, Koli DK, Rao IJ. Effect of phosphorus and biofertilizers on growth and yield of mungbean [*Vignaradiata (L.) Wilczek*]. International Journal of Current Microbiology and Applied Sciences. 2017;6(7):3992-3997.
9. Keh AO, Okocha IO, Umekwe PN, Amanze AN, Ikeh CE. Effect of foliar application of cow dung extract on growth and yield of waterleaf (*Talinum triangulare Jacq.*) in an ultisol. Journal of Current Opinion in Crop Science. 2023;4(3):103-111.
10. Ikeh AO, Orji JO, Sampson JO, HU. Akata, OR. Effect of oil palm bunch refuse ash in sustainable production of Egusi-Melon (*Colocynthis citrullus*) in an Ultisol. Asian Journal Agricultural and Horticultural Research. 2023;10 (3)138-148.  
DOI: 10.9734/AJAHR/2023/v10i324
11. Ikeh A, Ndaeyo O, NU Ikeh CE. Effects of integrated fertilization on soil sustainability and cassava (*Manihot esculenta Crantz*) yield in an ultisol, Journal of Current Opinion in Crop Science. 2023;4(2):89-102.

---

© 2023 Somkuwar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/108043>