



Condylar Height: Age and Sex Determination Using Orthopantogram in Forensics

L. Casilda Sushanthi ^a, Abirami Arthanari ^{b*} and Pratibha Ramani ^{a^o}

^a Department of Oral Pathology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-77, India.

^b Department of Forensic Odontology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-77, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i64B35439

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83003>

Original Research Article

Received 08 November 2021

Accepted 28 December 2021

Published 30 December 2021

ABSTRACT

Background: Since time immemorial, Condyles has been the area of interest for anthropologists, and the condyle size in men is higher than that of women. The condyle measurement is usually studied in two planes, i.e. anteroposterior and mediolateral, which are more gender-alliant. Different conventional X-rays are used for condylar imaging whose panoramic views of both condyles are provided by the OPG. For the creation of biological profiles during mass disasters the skull has remained useful. However, the technical procedures must in most cases be based on fragmented skull bones, when the whole skull is not available.

Materials and Methods: The study samples were retrieved from DIAS of Saveetha Dental College and Hospitals between the retrospective months from November 2020 to February 2021. The study sample consisted of 300 OPG's (150 males and 150 females), which were taken using Genoray Papaya machine, of age ranging from 10 to 24.9 years in which mandibular condylar height was measured using PlanmecaRomexis Software.

Results: The standard deviation for evaluation of condylar height for female is ± 4.38 while for males it is ± 4.91 . The standard error mean for condylar height for females is 0.35808 and for males is 0.40136 with non-significant p value of 0.893.

[≡]Senior Lecturer;

^oProfessor and Head;

*Corresponding author: E-mail: abiramia.sdc@saveetha.com;

Conclusion: By analysing the condylar mandibular height, the results of this study could improve human identification. For validation and comparison purposes, further research may be required in order to expand other associated parameters into a broader sample size.

Keywords: Condylar height; forensics; age estimation; sex estimation; orthopantogram.

1. INTRODUCTION

The mandibular condyle is a major growth site for the mandible and plays a vital role in its development. Endochondral ossification, or the sequential formation and degradation of the formed cartilaginous blastema serving as templates for bone formation, is just how the mandibular condyle develops. Appositional growth occurs when the perichondrium's chondrogenic cells emerge, proliferate, synthesise, and lay down new cartilage at the structure's edges. Whether condyles develop by apposition or by appositional combined with interstitial mechanisms is a point of contention. This is vital for bearing the compressive stress that the condyle receives during mandibular movement and development [1,2].

For the creation of biological profiles during mass disasters the skull has remained useful. However, the technical procedures must in most cases be based on fragmented skull bones, when the whole skull is not available. In these cases, the largest and most dimorphic part of the skull, the mandible plays an important role in determining the genitality of the individual remains. Age and gender assessment are one of medical law officers' important duties recently when crimes of a diverse nature are on the rise. In particular, mandibular condyle, ramus or angles are most sexually dimorphic, since they are locations associated with the largest dimorphous morphological channel in general [3,4].

The occlusal status and age of participants are considered to have anatomic changes in the mandible in adults. With age, the mandibular bone undergoes remodelling. Routine radiographs of the jaws are widely used panoramic radiographs. Since the mandible is asymmetrical, the condylar and ramus processes can be used to calculate vertical differences on both sides because they have a bilateral view and are suitable for vertical mandible measurements. Because of the nonlinear variance, various depths can be calculated without fear of the horizontal measurements becoming invalid [5,6]. The aim of the present study is to formulate new regression formula to

determine age and sex in South Indian Population.

2. MATERIALS AND METHODS

The study samples were retrieved from DIAS of Saveetha Dental College and Hospitals between the retrospective months from November 2020 to February 2021. The study sample consisting of 300 OPG's (150 males and 150 females) of age ranging from 10 to 24.9 years are tabulated in Table 1. The OPGs were taken using the Genoray Papaya machine.

Inclusion Criteria includes OPGs of patients within age group 10-24.9 while Exclusion Criteria includes OPGs of patients less than 10 years and above 24 years. Measurement of mandibular condylar height from OPGs were done using PlanmecaRomexis Software. ConH (Condylar Height) was measured for all age groups (Fig. 1 and Fig. 2). Statistical analysis was done using SPSS and regression formula was derived.

Table 1. Sample distribution among the various age groups

GROUP S	AGE(years)	MALE	FEMALE
1	10 to 10.9	10	10
2	11 to 11.9	10	10
3	12 to 12.9	10	10
4	13 to 13.9	10	10
5	14 to 14.9	10	10
6	15 to 15.9	10	10
7	16 to 16.9	10	10
8	17 to 17.9	10	10
9	18 to 18.9	10	10
10	19 to 19.9	10	10
11	20 to 20.9	10	10
12	21 to 21.9	10	10
13	22 to 22.9	10	10
14	23 to 23.9	10	10
15	24 to 24.9	10	10
TOTAL		150	150

3. RESULTS AND DISCUSSION

During endochondral ossification of long bones, several differentiation factors, growth factors and angiogenic mediators played a key role. Similarly, the growth of condyles must be regulated by a host of orchestrated influences of different growth factors and other endogenous regulatory factors in condyles [7,8,9,10].

The present study revealed that the mean of condylar height measurement was slightly higher in males compared to females. The standard deviation for evaluation of condylar height for female is ± 4.38 while for males it is ± 4.91 . The standard error mean for condylar height for females is 0.35808 and for males is 0.40136 with non-significant p value of 0.893 (Table 2). The standard error mean for age for females is 0.34469 and for males is 0.34528 with non-significant P value of 0.984 (Table 2). The insignificance in the statistical analysis could be due to the limited number of sample size in each age groups.

Population-specific standards need always to be developed for precise gender determination based on ethnicity. In many studies for various populations worldwide measurement standards were therefore developed. Skeletal remains have differential characteristics from one population to another because osteometric standards have been developed for the determination of gender [11,12,13]. Hence, in the present study Indian specific regression formula was formulated to calculate the age of an individual which will be of

ease in identifying/estimating age of an unknown individual, which is most important in forensic dentistry.

Since time immemorial, Condyles has been the area of interest for anthropologists, and the condyle size in men is higher than that of women. The condyle measurement is usually studied in two planes, i.e. anteroposterior and mediolateral, which are more gender-alliant. Different conventional X-rays are used for condylar imaging whose panoramic views of both condyles are provided by the OPG. The mandibular condyle is a growing centre, and has a different morphology as a functional joint unit. The age, gender, face type, functional load, occlusal forces, malocclusion type between the right and left sides occur as a normal variation in the morphology of the condylars [14,15,16].

Rajkumari et al. [17], states that the Condylar height, projected mandible height and coronoid height increase with age, and a slight reduction in the values occurs in the sixth decade.

Several studies have been performed by many researchers, specifically vertical, horizontal and angle measures for the applicability of mandibular metric analysis in forensic odontology. The minimum ramus width and Condylar height, which were significantly lower in sexual prediction that contradicted few other studies with the greatest sexual dimorphism of the minimum ramus breadth and Condylar height [17,18,19].

Table 2. Since p values are greater than 0.05 (P>.05) there is no significant difference between Age as well as Condylar Height

	Gender	N	Mean	Std. Deviation	Std. Error Mean	t-test/p value
Age	Females	150	17.1924	4.22154	.34469	0.984
	Males	150	17.1828	4.22879	.34528	
Condylar height	Females	150	29.3553	4.38559	.35808	0.893
	Males	150	29.4280	4.91560	.40136	

**Regression formula for identification of an unknown individual
 Females = 16.237 + 0.033 × CH; SD ±4.38 years
 Males = 17.097 + 0.03 × CH; SD ±4.91 years*



Fig. 1. OPG for condylar height measurement of female



Fig. 2. OPG for condylar height measurement of male

4. CONCLUSION

In conclusion, the mandible may be present in numerous morphological forms and measures. Different socio-demographic factors that could contribute to its final appearance may influence those factors. By analysing the condylar mandibular height, the results of this study could improve human identification. For validation and comparison purposes, further research may be required in order to expand other associated parameters into a broader sample size.

CONSENT AND ETHICAL APPROVAL

As per international standard or university standard guideline Patient's consent and ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Li QF, Rabie ABM. A new approach to control condylar growth by regulating angiogenesis. *Arch Oral Biol.* 2007;52(11): 1009–17.
- Yamunadevi A, Pratibha R, Rajmohan M, Ganapathy N, Porkodisudha J, Pavithrah D, et al. Molecular Insight into Odontogenesis in Hyperglycemic Environment: A Systematic Review. *J Pharm Bioallied Sci.* 2020;12(Suppl 1): S49–56.
- Behl AB, Grewal S, Bajaj K, Baweja PS, Kaur G, Kataria P. Mandibular ramus and gonial angle—Identification tool in age estimation and sex determination: A digital panoramic radiographic study in north indian population. *Journal of Indian Academy of Oral Medicine and Radiology.* 2020;32(1):31.
- Antony JVM, Ramani P, Ramasubramanian A, Sukumaran G. Particle size penetration rate and effects of smoke and smokeless tobacco products - An invitro analysis. *Heliyon.* 2021;7(3): e06455.
- Bhuyan R, Mohanty S, Bhuyan SK, Pati A, Priyadarshini S, Das P. Panoramic radiograph as a forensic aid in age and gender estimation: Preliminary retrospective study. *J Oral Maxillofac Pathol.* 2018;22(2):266.
- R H, Hannah R, Ramani P, Ramanathan A, Jancy MR, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology.* 2020;130: 306–12. Available:<http://dx.doi.org/10.1016/j.oooo.2020.06.021>
- Rabie ABM, Hägg U. Factors regulating mandibular condylar growth. *Am J Orthod Dentofacial Orthop.* 2002;122(4):401–9.
- Umashankar K, Abilasha, Hannah, Ramani P, Gheena. Knowledge and attitude about COVID-19 pathogenesis among oral pathologists in Chennai. *Int J Curr Res Rev.* 2020;12(19):143–51.
- Suvarna K, Abilasha R, Gheena S, Ramani P. Analysis of Prevalence of Oral Squamous Cell Carcinoma in Patients with History of Chronic Irritation of Oral Tissues-A Retrospective Study. *Indian J Forensic Med Toxicol* [Internet]. 2020; 14(4). Available:<http://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=09739122&AN=148410023&h=XYlwRUZ0p54Y6mZHESI7LVi228pkq%2FRhoPO4Jt4231oGPh6KuqA4bM6e9ZsIMs9HCBZY6ZF8tl2TIR3cgyExXQ%3D%3D&crl=c>
- Behera A, Hannah R, Ramasubramanian A, Ramani P. Association of the Depth of Invasion with Lymph Node Metastasis in Oral Squamous Cell Carcinoma Patients-A Retrospective Study. *Indian J Forensic Med Toxicol* [Internet]. 2020;14(4). Available:<http://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=09739122&AN=148410053&h=nnwHkKvd5DGQXazZu6t5p6l8D0PbUqebPTLzhmdSI6FyXTWsSzGTNdBhQxTZUG9F%2BD8YcgHrW%2FSaXuV%2BCg9vCA%3D%3D&crl=c>
- Alias A, Ibrahim A, Abu Bakar SN, SwarhibShafie M, Das S, Abdullah N, et al. Anthropometric analysis of mandible: an important step for sex determination. *Clin Ter.* 2018;169(5):e217–23.
- K M, Monica K, Vijayshree PJ, Gheena S, Ramani P, Abhilasha R, et al. *In silico* gene expression analysis of crucial cell cycle control gene *cdkn2a* and *cdkn2b* in head and neck squamous cell carcinoma [Internet]. *Annals of Tropical Medicine & Public Health.* 2020;23. Available:<http://dx.doi.org/10.36295/asro.2020.232323>
- Sinduja P, Ramani P, Gheena S, Ramasubramanian A. Expression of metallothionein in oral squamous cell carcinoma: A systematic review. *J Oral Maxillofac Pathol.* 2020;24(1):143–7.
- Dwivedy S, Chandra S, Srivastava A, Chandra S, Shrestha P, Thakur R. A novel approach toward mandibular condyle imaging and quantification through modified reverse panoramic radiography for determination of gender. *International Journal of Forensic Odontology.* 2017; 2(2):67.
- Ramani P, Krishnan RP, Karunakaran M, Muthusekhar MR. Odontogenic sarcoma: First report after new who nomenclature with systematic review. *J Oral Maxillofac Pathol.* 2020;24(1):157–63.
- Romañach M, Canedo N, Cortezzi E, Abrahão A, Cabral M, Agostini M. Clear

- cell variant of oral squamous cell carcinoma. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2014;118(6):e195.
17. Rajkumari S, Nikitha K, Monisha S, Nishagrade S, Thayumanavan B, Murali BA. Role of Orthopantamograph in Forensic Identification: A Retrospective Study among Chennai Population. *J Pharm Bioallied Sci.* 2019;11(Suppl 2): S393–6.
18. A study on the variability of drug responsiveness to anti inflammatory drugs - A pilot survey. *Int J Pharm Res [Internet].* 2020;12(02). Available:<http://www.ijpronline.com/ViewArticleDetail.aspx?ID=17202>
19. Thamilselvan S, Abilasha R, Ramani P, Gheena S, Hannah R. Evaluation of Accuracy between Habit History and Incidence of Oral Squamous Cell Carcinoma [Internet]. *International Journal of Current Research and Review.* 2020; 30–5. Available:<http://dx.doi.org/10.31782/ijcrr.2020.122503>
18. A study on the variability of drug responsiveness to anti inflammatory drugs - A pilot survey. *Int J Pharm Res [Internet].*

© 2021 Sushanthi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83003>