



A Cross-Sectional Investigation of the Public Health Consequences of Hepatitis B infection among Abattoir Workers in Port Harcourt Nigeria

Chidi L.C. Ndukwu ^{a*} and Jane U. Chinedu-Madu ^a

^a *Faculty of Medical Laboratory Science, Federal University Otuoke, Bayelsa State, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123805>

Original Research Article

Received: 22/07/2024

Accepted: 24/09/2024

Published: 27/09/2024

ABSTRACT

Introduction: Infection with Hepatitis B virus (HBV) is a public health threat accounting for an estimated nearly one million deaths annually largely due to cirrhosis and hepatocellular carcinoma. Abattoir workers were reported to be at high risk of contracting HBV infection. This study was conceived to ascertain the public health consequences of HBV infection among abattoir workers.

Materials and Methods: About 3 - 4 millilitres of venous blood samples were collected from each of 275 participants into pre-labelled EDTA vacutainer tubes and transported in a cold box to the Laboratory. They were analyzed with three different rapid diagnostic test strips and positive results were recorded only for specimens that were reactive to at least two test strips. Test of association between HBV and abattoirs, gender and age were conducted using Pearson's Chi-Square test of independence and Fisher exact test at significance level below 0.05

*Corresponding author: E-mail: chidisteve.cn@gmail.com;

Cite as: Ndukwu, Chidi L.C., and Jane U. Chinedu-Madu. 2024. "A Cross-Sectional Investigation of the Public Health Consequences of Hepatitis B Infection Among Abattoir Workers in Port Harcourt Nigeria". *Asian Journal of Research and Reports in Hepatology* 6 (1):37-44. <https://journalajrrhe.com/index.php/AJRRHE/article/view/39>.

Results: The total prevalence of HBV for all the abattoirs was 6.2%. Tereama was dominant with 13.9% positive results, followed by Iloabuchi (12.5%) Wazobia (10.7%), Eagle Island 8.3% Mile three 6.5% and Oil mill 4.7%. Zero prevalence was recorded in six abattoirs. A chi-square test of independence to evaluate the relationship between the independent and dependent variables showed no significant relationship between them, as the p value of .742 is not less than 0.05, therefore the null hypothesis which states that the variables are independent is not rejected.

Conclusion: This study revealed an intermediate prevalence of HBV among abattoir workers in the same range with the general population. The abattoir workers, it appears were not exposed to any more risks of contracting the infection than the general population in the study area. All the same the prevalence of 6.2% was unacceptable and efforts need to be intensified to ensure the eradication of the infection.

Keywords: Hepatitis B virus; public health; meat handlers; abattoirs; seroprevalence.

1. INTRODUCTION

Infection with Hepatitis B virus is an insidious global public health threat accounting for an estimated nearly 300 million persons currently living with the chronic condition, and about 30 million people being infected annually [1]. The estimated number of deaths due to hepatitis B infection is put at about one million persons with over eighty percent attributable to cirrhosis and hepatocellular carcinoma (HCC), making hepatitis B the 15th commonest cause of human mortality worldwide [2,3]. The high number of deaths associated to HBV is largely due to the large number of undiagnosed cases, with less than 10% of infected persons being diagnosed and less than 1%, receiving treatment. The disease burden of hepatitis B is heavy on the African continent with over 60 million infected persons hepatitis B, which amounts to over 60,000 deaths each year [3].

Nigeria is one of the African countries with high prevalence of HBV, defined as more than 8% with a vast majority of the infected persons being ignorant of their status and therefore not receiving treatment. It is thus not surprising that the country has one of the highest incidences of liver cancer caused by HBV, estimated at close to 5.1 cases per 100,000 person-years [4].

Hepatitis B virus is a small enveloped DNA virus. It is hepatotropic and belongs to the family *hepadnaviridae*; and has a narrow host range, infecting only humans and a few related primate species [5]. It is transmitted through both vertical and horizontal routes, majorly, mother to child, sexual intercourse and parenterally through blood and blood products [6]. Vertical transmission through mother-to-child at birth, is recognized as the prevalent mode of transmission of the virus in areas of endemicity,

like Sub-Saharan Africa and Asia-Pacific countries; and those who become infected at infancy are more likely to become chronic carriers if not treated on time. On the other hand, horizontal transmission at adult ages is more likely to result in self-limiting acute forms of the infection [5]. It is a resilient virus, which is highly stable at 37 °C and can survive on environmental surfaces for more than 22 days; it is detectable in blood and body fluids such as saliva, tears, sweat, semen, and vaginal secretions. In areas of low prevalence, injection drug use and high-risk sexual exposures are the commonest modes of transmission [6].

Hepatitis B virus infections may be acute or chronic forms. The post-infection incubation period of the acute infection ranges from one to four months. Symptoms in the acute stage may only be found in adult where about one third of infected persons manifest serum sickness type symptoms such as fever, arthralgias, and rash, which may occur in the prodromal period; followed by constitutional symptoms, anorexia, nausea, jaundice, and right upper quadrant discomfort. Liver function tests are marked with elevated serum alanine transaminase and mild elevation in bilirubin. The most notable complication of the acute infection acute liver failure [7].

The chronic infection is a resultant effect of interactions between the host and viral factors, and occurs when viral clearance in the acute stage is ineffective. It is characterized by persistence of serum HBsAg for at least 6 months [7]. The induced immune response following the infection of human hepatocytes leads to the elimination of infected cells especially in immunocompetent persons. In cases of immature or compromised immunity the formation of chronic disease or their reactivation

after remissions are the likely outcomes. In adults HBV infection results in acute infection in 90–95% of cases, while in the case of infants chronicity is the outcome in almost all cases [8].

Although the infection of humans by hepatitis B virus is reported to have been in existence for 2000–3000 years, the virus was only discovered until 1966, and in 1970 the virus particle was identified by Dane *et al.*, by the use of electron microscopy. Consequently, antiviral therapies and vaccines has been produced in the bid to combat the viral menace. In spite of the efforts and breakthrough in the treatment and prevention of the infection, the mortality and morbidity remain high [9].

Horizontal transmission of HBV through occupational, recreative and similar activities occasioning inadvertent contacts with human fluids particularly blood has received attention in recent times. Studies have considered slaughter house workers as high-risk groups for HBV infection, given that they are exposed to a number of risks such as knife cuts, close contacts with animal blood, exchange of knives in the course of their occupational activities [10,11]. It has also been reported that hepatitis virus has been detected in some animals like cattle, and the fact that a large proportion of pathogenic organisms are zoonotic organisms [11].

This study was thus conceived to ascertain the prevalence of hepatitis B virus among persons involved in slaughtering animals and cutting up of meat for sale to the public in public abattoirs in Port Harcourt with the intent of determining the public health consequences.

2. MATERIALS AND METHODS

2.1 Study Design

This is an observational, retrospective and cross-sectional research. It was commissioned by the state environmental health department in Port Harcourt as part of routine screening and monitoring of meat handlers within the abattoirs in the metropolis. The screening exercise was coordinated by the staff of the department and the executives of the various abattoirs' unions. The study was conducted between October 2014 and February 2015.

2.2 Study Site

The area for the study is Port Harcourt, the capital and major city of Rivers. It is located on

Latitude: 4°46'38" N Longitude: 7°00'48" E with an Elevation above sea level: 16m. Twelve abattoirs spread across the Port Harcourt metropolis were involved in the study. The abattoirs were located at Tereama, Mile 3, Eagle Island, Emenike, Wazobia, Iloabuchi, St Andrews, Nkpor, Rukpokwu, Shukura, Choba and Oil mill.

2.3 Subject Selection and Demographics

All the butchers and meat handlers including those involved in slaughtering, wholesales and retail trade of the mostly cow and goat meat were involved in the study. Semi-structured interviewer- administered questionnaire and an observation checklist were used to obtain demographic data such as age, sex, length of stay on the job. Only persons above the age eighteen and who have been on the job for at least one year were included in the study. A total of 275 blood samples were collected from persons who freely gave their informed consents.

2.4 Sample Collection

A total of 275 blood samples were collected from persons who freely gave their consents after the reasons and procedures have been well explained and their questions entertained to their apparent satisfaction. About 3 - 4 millilitres of venous blood samples were collected from each study participant into a pre-labelled EDTA vacutainer tubes and transported in a cold box to the Diagnostix and Scientifique Laboratory, a medical diagnostic and research laboratory offering services for public and private healthcare facilities in Port Harcourt for analysis. The laboratory analyses were conducted with three different rapid diagnostic test strips (names not recorded) and positive results were recorded only for specimens that tested reactive for at least two of the test strips.

2.5 Data Analysis

Data were cleaned using Excel spreadsheet 2016, and analyzed using IBM SPSS Statistics version 25. Descriptive and inferential statistics were employed in results presentation and interpretation. Associations between possible risk factors namely abattoirs, gender, age and hepatitis B infections were determined using Pearson's Chi-Square test of independence and Fisher exact test at significance level below 0.05.

3. RESULTS

The descriptive statistics indicate that 275 participants were involved in the study. The youngest was 20 years while the eldest was 67 years old. The mean age was 38.35 years with standard deviation of 9.752; the median and modal ages were 37 and 40 years respectively. All the data as analyzed were complete.

3.1 Prevalence of Hepatitis B infections among of Meat Handlers in Abattoirs in Port Harcourt Nigeria

The prevalence of hepatitis B infection among the abattoir workers as observed in this study was 6.2%. The highest prevalence was observed at the Tereama abattoir (13.9%), it contributed 13.1% of the overall number of samples and 5 (29.4%) of the positive results. It was followed by Iloabuchi (12.5 %), yielding 8.7% of the total number and 3 (17.7%) of the positive results. Wazobia (10.7 %), produced 10.8% of the overall Figure, 3 (17.7%) of the positive results. The Eagle Island (8.3%) accounted for 4.4% of the whole samples and one (5.9%) of the positive samples. Oil Mill (7.3%) the second abattoir recorded 15.6% of the entire samples and 2 (11.8) of positive samples; while the predominant abattoir, Mile 3 (6.5%) yielded 16.7% of the overall and 3 (17.7%) of the positive results.

A review of the records of 275 specimens tested for HBV in this study showed that 271 of them were males while 4 were females. The prevalence of HBV infection among the males was 5.9%, accounting for 16 (94.1%) of the total prevalence; the prevalence for the females was 25%, contributing 1 (5.9%) to the overall prevalence (Table 1).

3.2 Prevalence of Hepatitis B infections across Age Groups of Meat Handlers in Port Harcourt Nigeria

The 20-29 age group recorded the highest of prevalence of 8.7% among all the age groups; but accounted for 16.7% of the total 275 persons and 4 (23.5%) of the 17 positive results. The 30-39 group was the dominant group by contributing 40.7% of the overall 275, and recorded the second highest group prevalence of 7.1%, contributed 8 (47.1%) of the positive results. The second predominant group was 40-49 producing 27.3% of the overall patients, had a prevalence of 5.3% and accounted for 4 (23.5%) of the 17 positive results. The 50-59 age group had a prevalence of 3.2%, produced one (5.9%) of the

positive results and yielded 11.3%% of the overall number. Zero prevalence was only recorded with the 60-69 age group which contributed the least (4%) to the overall Figure. (Table 1).

3.3 Statistical Analysis

Chi-square test of independence and Fisher's exact test was performed to evaluate the relationship between the independent variables (abattoirs, age and gender) and results of HBsAg tests (dependent variable). The relationship between these variables were found not be significant, given that the p values were not less than 0.05, we therefore failed to reject the null hypothesis which states that the variables are independent. In other words, there was no sufficient evidence to conclude that a significant association exists between the variable and the test results obtained for the hepatitis B screening.

4. DISCUSSION

This study has been able to establish the prevalence of hepatitis B infection among meat handlers in twelve abattoirs across Port Harcourt metropolis in Nigeria. A number of previous studies had reported high prevalence of viral hepatitis, including Hepatitis B virus among meat handlers at abattoirs. A national survey on seroprevalence of hepatitis B put the national prevalence at 12. 2% [12] Other studies across the country has given the prevalence as 9.5% [4] ,12.6% and 15.2% [13], 2.23% [14], and13.1% [15]. These differences were shown not to be statistically significant and so does not pose much public health threat.

The prevalence of 6.2% found for HBV infection in this study in lower than most values reported elsewhere across the country. It is much lower than 13.7% reported for abattoirs in Port Harcourt [16], but higher than an earlier 4.3% also in Port Harcourt among pregnant women [12] however, it is in alignment with a prevalence of 5.9% [17] and 6.49% [18] reported in recent nationwide studies of HBV prevalence in pregnant women and 5.1% [19] also among persons attending a Primary Health Centre in Port Harcourt. It is apparent that the prevalence of HBV varies widely in time and space. This could be gleaned from the prevalence across the abattoirs which ranged from zero in 50% of the abattoirs to 13.9% in one of the abattoirs.

Table 1. Prevalence of Hepatitis B infections among Meat Handlers across Abattoirs in Port Harcourt Nigeria

Characteristics	Number Tested	Number Positive
Abattoirs		
Choba	18 (6.6)	0 (0.0)
Eagle Island	12 (4.4)	1 (8.3)
Emenike	21 (7.6)	0 (0.0)
Iloabuchi	24 (8.7)	3 (12.5)
Mile 3	46 (16.7)	3 (6.5)
Nkpor	12 (4.4)	0 (0.0)
Rukpokwu	16 (5.8)	0 (0.0)
Shukura	12 (4.4)	0 (0.0)
St Andrews	7 (2.6)	0 (0.0)
Tereama	36 (13.1)	5 (13.9)
Wazobia	28 (10.8)	3 (10.7)
Total	275 (100)	17 (6.2)
Gender		
Male	273 (99.3)	16 (5.9)
Female	4 (0.07)	1 (25)
Total	275 (100)	17 (6.2)
Age Groups		
20-29	46 (16.7)	4 (8.7)
30-39	112 (40.7)	8 (7.1)
40-49	75 (27.3)	4 (5.3)
50-59	31 (11.3)	1 (3.2)
60-69	11 (4.0)	0 (0.0)
Total	275 (100)	17(6.2)

The result from this study which is an intermediate prevalence does not differ much from the results for other populations in Port Harcourt such as pregnant women [12]. It is however lower than those obtained for abattoir workers and livestock merchants with a reported prevalence of 10.9% [20]. in Ibadan, and 8.0% reported for donkey butchers in Ebonyi state [21]. The hyper prevalence in those areas may align with the prevailing population prevalence, while it may be said that abattoir workers in Port Harcourt have imbibed safety practices in their jobs, especially given there was a decline from a previous reported prevalence; and the fact that during the course of the screening and preceding ones, there were enlightenment and awareness campaigns on safety practices in the abattoirs.

Whereas several reports had focused on the prevalence of HBV among the general population and some high-risk groups across the country, there appear to have been not much reports on the prevalence of the virus in the abattoirs were found. However, the factors that contribute to put abattoir workers in high risk as it pertains to HBV infection have been studied by a number of researchers. Personnel involved in meat processing for instance are known to be exposed to a lot of biological and physicochemical agents which may result in infectious and noninfectious diseases [10].

Some of the factors that put abattoir personnel at risk of HBV include knife cuts, sharing of knives, cut from bones, injury by animals, close, long contacts with animals and animal products among others [18,21]. The risks associated working with animals is heightened by the fact that vast majority of emerging infections are zoonotic which makes animals potential carriers of unrecognized pathogens.

It is remarkable that over eighty percent of the participants and over ninety percent of those with positive results are in the age brackets below fifty years. The above sixty age bracket recorded no positive results, while the 50 -59 group had only one positive case. Though the chi square test of independence did not reveal any significant association between HBV results and age groups, it can be seen that the most active age brackets are in the below fifty groups and any efforts at combating the HBV infection should be focused on the younger age brackets. The wide disparity among the genders as seen in this study is not surprising, given that abattoir workers are predominantly males.

The limitations of the study are mainly those faced in retrospective studies, given that the samples have been analyzed before the commencement of the study. The sociodemographic data are limited to those contained in the laboratory records, and there is no personal interaction with the participants. Some information that may enrich the study are therefore not captured. Future studies may be designed to overcome these limitations [22].

5. CONCLUSION

This study has been able to ascertain and present the prevalence of HBV among abattoir workers in Port Harcourt. The intermediate prevalence level found in this study are within the same range as the reported among the general population in the metropolis; thus, the abattoir workers appear not to be exposed to any higher levels of risks to the infection than those faced by the general population. All the same, the prevalence of 6.2% observed in this study is not an acceptable one. Efforts should be intensified to eradicate HBV infection through mass vaccination and creation of awareness about safety practices, to safeguard the public from the dangers associated with hepatitis B infection.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Asandem DA, Segbefia SP, Kusi KA, Bonney JHK. Hepatitis B Virus Infection: A Mini Review. *Viruses*. 2024;16(5):724. DOI: 10.3390/v16050724. PMID: 38793606; PMCID: PMC11125943

2. MacLachlan JH, Cowie BC. Hepatitis B virus epidemiology. *Cold Spring Harb Perspect Med.* 2015;5(5):a021410. DOI:10.1101/cshperspect.a021410 PMID: PMC4448582
3. Sandem DA, Segbefia SP, Kusi KA, Bonney JHK. Hepatitis B Virus Infection: A Mini Review. *Viruses.* 2024;16(5):724. DOI: 10.3390/v16050724. PMID: 38793606; PMID: PMC11125943.
4. Ajuwon BI, Yujuico I, Roper K. *et al.* Hepatitis B virus infection in Nigeria: a systematic review and meta-analysis of data published between 2010 and 2019. *BMC Infect Dis.* 2021;21:1120. Available:<https://doi.org/10.1186/s12879-021-06800-6>
5. Chuang YC, Tsai KN, Ou JJ. Pathogenicity and virulence of Hepatitis B virus. *Virulence.* 2022;13(1):258-296. DOI: 10.1080/21505594.2022.2028483 PMID: 35100095; PMID: PMC8812780.
6. Sabeena S, Ravishankar N. Horizontal Modes of Transmission of Hepatitis B Virus (HBV): A Systematic Review and Meta-Analysis. *Iran J Public Health.* 2022;51(10):2181-2193. DOI: 10.18502/ijph.v51i10.10977. PMID: 36415805; PMID: PMC9647610.
7. Campos-Valdez M, Monroy-Ramírez HC, Armendáriz-Borunda J, Sánchez-Orozco LV. Molecular Mechanisms during Hepatitis B Infection and the Effects of the Virus Variability. *Viruses.* 2021;13(6):1167. DOI: 10.3390/v13061167. PMID: 34207116; PMID: PMC8235420.
8. Ogunnaike M, Das S, Raut SS, Sultana A, Nayan MU, Ganesan M, Edagwa BJ, Osna NA, Poluektova LY. Chronic Hepatitis B Infection: New Approaches towards Cure. *Biomolecules.* 2023;13(8):1208. DOI: 10.3390/biom13081208. PMID: 37627273; PMID: PMC10452112.
9. Sekiba K, Otsuka M, Ohno M, Yamagami M, Kishikawa T, Suzuki T, Ishibashi R, Seimiya T, Tanaka E, Koike K. Hepatitis B virus pathogenesis: Fresh insights into hepatitis B virus RNA. *World J Gastroenterol.* 2018;24(21):2261-2268. DOI: 10.3748/wjg.v24.i21.2261. PMID: 29881235; PMID: PMC5989240.
10. Johnson OE, Etokidem AJ. Occupational Hazards and Health Problems among Butchers in Uyo, Nigeria. *Niger Med J.* 2019;60(3):106-112. DOI: 10.4103/nmj.NMJ_57_19. PMID: 31543560; PMID: PMC6737801.
11. Tariq H, Kamal MU, Makker J, Azam S, Pirzada UA, Mehak V, Kumar K, Patel H. Hepatitis in slaughterhouse workers. *World J Hepatol.* 2019;11(1):37-49. DOI: 10.4254/wjh.v11.i1.37. PMID: 30705717; PMID: PMC6354121.
12. Olayinka AT, Oyemakinde A, Balogun MS, Ajudua A, Nguku P, Aderinola M, Egwuenu-Oladejo A, Ajisegiri SW, Sha'aibu S, Musa BO, Gidado S, Nasidi A. Seroprevalence of Hepatitis B Infection in Nigeria: A National Survey. *Am J Trop Med Hyg.* 2016;95(4):902-907. DOI: 10.4269/ajtmh.15-0874. Epub 2016 Aug 15. PMID: 27527630; PMID: PMC5062798.
13. Chikwendu A, Unikutelle HL, Olumide AT. Hepatitis B and C virus prevalence among patients and healthcare workers' prevention practices towards the viruses in a secondary healthcare facility in Northern Nigeria. *Pan Afr Med J.* 2023;46:82. DOI: 10.11604/pamj.2023.46.82.40530. PMID: 38314232; PMID: PMC10837276.
14. Ojerinde OA, Ojo SKS, Udewena UL, Oladeji SJ. A cross-sectional study on the prevalence of HIV and hepatitis B virus co-infection among students of a tertiary institution in Ekiti State, Southwest Nigeria. *Pan Afr Med J.* 2023;44:7. DOI: 10.11604/pamj.2023.44.7.31416. PMID: 36818031; PMID: PMC9935660.
15. Fasola FA, Fowotade AA, Faneye AO, Adeleke A. Prevalence of hepatitis B virus core antibodies among blood donors in Nigeria: Implications for blood safety. *Afr J Lab Med.* 2022;11(1):1434. DOI: 10.4102/ajlm.v11i1.1434. PMID: 35937767; PMID: PMC9350478.
16. Ibanga ES, Adikema NA, Adebayo O, Adegoke Margaret Amala S, Emejuru AK. Prevalence of Hepatitis B Virus among Butchers in Port Harcourt Metropolis, Elixir Physio. & Anatomy. 2015;79:30651-30654
17. Federal Ministry of Health. Nigeria HIV/AIDS Indicator and Impact Survey (NAIIS) 2018: Final report. Abuja; 2019.
18. Olakunde BO, Adeyinka DA, Olakunde OA, Uthman OA, Bada FO, Nartey YA, Obiri-Yeboah D, Paintsil E, Ezeanolue EE. A systematic review and meta-analysis of the prevalence of hepatitis B virus infection among pregnant women in Nigeria. *PLoS One.* 2021;16(10):e0259218. DOI: 10.1371/journal.pone.0259218. PMID: 34714888; PMID: PMC8555786.

19. Erasmus MA, Wokem GN. Evaluation of Hepatitis-B and Its Markers Among Some Attendees of Two Health Facilities In Port Harcourt, Rivers State, Nigeria, Nigerian Journal of Parasitology. 2021;42(2). DOI: 10.4314/njpar.v42i2.27
20. Opayele AV, Arege OT, Faneye AO, Olaleye DO, Odaibo GN. Prevalence of HIV, HBV and HCV among livestock merchants and slaughterhouse workers in Ibadan, Nigeria. Afr Health Sci. 2024;24(1):16-24. DOI: 10.4314/ahs.v24i1.4. PMID: 38962339; PMCID: PMC11217837
21. Elom P, Unah A, Onwasigwe C. Comparative study of knowledge, risk factors, and prevalence of Hepatitis B virus Infection Among Donkey Butchers and Herders in Ebonyi state, Nigeria. African Journal of Gastroenterology and Hepatology. 2024;7(1):64-97. DOI: 10.21608/ajgh.2024.245381.1045
22. Ndukwu CLC. Microbial communities and antimicrobial resistance patterns in aerobic bacteria associated with the *Vaginal microbiota*: A retrospective study in port harcourt, Nigeria. Asian Journal of Research in Infectious Diseases. 2024;15(1):39–48. Available: <https://doi.org/10.9734/ajrid/2024/v15i1324>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/123805>