



Neck Circumference, a Screening Tool for Obesity in Newly Pubescent Children-A Case Study

M. Ponmalar^{1*}, S. Rajini¹ and V. Tamilsevi¹

¹Department of Community Medicine, Sri Lakshmi Narayana Institute of Medical Sciences,
Puducherry, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i44A32645

Editor(s):

(1) Prof. Juan Carlos Troiano, University of Buenos Aires, Argentina.

Reviewers:

(1) Ana Cristina García Ulloa, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico.

(2) Sanjeev Kumar, L. N. Mithila University, India.

Complete Peer review History: <https://www.sdiarticle4.com/review-history/73608>

Original Research Article

Received 08 July 2021
Accepted 18 September 2021
Published 21 September 2021

ABSTRACT

“Childhood Obesity and its co-morbidities is considered as a prime health hitch that lay up consideration among the public health society. Obesity and above the over weight has bagged the fifth position in the global risk for mortality. The obesity shows a very early onset in life which has laid the utmost importance in the prevention of it among the children. Neck circumference is being considered as the marker of upper body subcutaneous tissue distribution.

Aim: The main objective is to evaluate the association between neck circumference and obesity among the children.

Methodology: This is a cross sectional study conducted between the periods of august 2019 - September 2019, among 150 male + 129 female students, aged 11 to 18 years. Anthropometric markers of obesity- body mass index (BMI), waist circumference (WC), Hip circumference (HC), waist hip ratio (WHR), neck circumference (NC) were collected. Pearson's correlation coefficient was used to compute the neck circumference with other obesity indices.

Results: The mean BMI was 27.70 ± 1.27 kg/m² and 27.72 ± 2.12 kg/m², mean neck circumference was 35.42 ± 2.26 cm and 32.85 ± 1.92 in above the over weight /obesity boys + girls respectively. The hip circumference in boys was found to be greater than girls, which is higher in above the over weight /obesity students with $P < 0.001$. Neck circumference shows a strong positive correlation with other anthropometric measures BMI, WC, HC in boys and girls ($P < 0.001$).

Conclusion: The present study showed that the weight increase in school children and need to aware the parents.

Keywords: BMI; NC; above the over weight; obesity; WC; HC; children.

1. INTRODUCTION

Above the over weight is defined as body mass index between 25 and 29.9 kilogram per meter square and obesity is defined as BMI of 30 kilogram per meter square or higher. Obesity is a pathological condition in which excess body fat accumulated, leading adverse effects on health and life expectancy [1]. It is a chronic disorder with complex interaction between genetic and environmental factors. It characterized by high cholesterol, fatty acid levels; imbalance in metabolic energy; insulin desensitization; lethargy, gallstones; high blood pressure; shortness of breath; emotional and social problems; and excessive adipose mass accumulation with hyperplasia and hypertrophy [2]. Pathological obesity is associated with several secondary commodities like heart disease, type 2 diabetes, breathing difficulties during sleep, cancer and osteoarthritis.

It is most commonly caused by a combination of excessive dietary calories, lack of physical activity, and genetic susceptibility. Evidence to support this view is that some obesity people eat little yet gain weight due to slow metabolic rate. The primary treatment for obesity are dieting and physical exercise. To addition to this, or in case of failure, anti-obesity drugs may be taken to reduce appetite or impede fat absorption [3]. In serious cases, surgery is carryout or an intragastric balloon is placed to lower stomach volume and/or bowel length, precede to earlier satiation and make less ability to absorb nutrients from food [4].

Obesity can be associated with several risk factors such as cardiovascular and metabolic disturbances. There are numerous methods for assessing above the over weight and obesity. Some techniques are applicable at primary care facilities, such as measurements of weight, height, hip circumference, calculations of waist-hip ratio and BMI. It is important to develop a quick method of for the assessment of obesity in primary care clinic. The purpose of the study to determine whether neck circumference is useful alone for assessing obesity [5].

The restraint of BMI as a risk evaluation tool is also conceded, and there is continuing interest in alternative or complementary indices linking body adiposity and disease risk. For example, some studies suggest that abdominal circumference is better correlated than BMI with the risk of type 2 diabetes [6]. Although it is well established that visceral adiposity plays a central role in the metabolic disorders associated with obesity, the dearth of a pragratis method to assess visceral fat in routine examinations precludes its use as a screening tool for the public [7-9]. Forkhead box O (FOXO) transcription factors are the main factors that contribute the direct and indirect pathways for glucose homeostasis. The recent studies showed that it actively hyperglycemia in chronic insulin resistance in obesity patients [10]. Developing simple and authentic methods to assess body fat compartments should be an important prime concern of obesity research.

1.1 Objective

Above the over weight in children is described by using body mass index along with BMI, waist-hip ratio, mid-upper arm circumference, waist thigh ratio indices of body fatness is needed to determine obesity.

The aim of the study to determine whether neck circumference is useful alone for assessing obesity, Children who were to 11 to 18 yrs and undergoing schooling where the subject of the study, trained CRRJ collected the anthropometric data from school-going children. We calculated and correlated the neck circumference and other indexes of obesity and we analyzed the optimal neck circumference cutoff for identifying children with high BMI.

2. METHODOLOGY

A cross-sectional study was conducted during the period of August 2019- September 2019 in the private schools, of Pondicherry. Participants were of age 11-18 years comprising both males and females as the separate groups. Considering approximately 50 private schools in Puducherry , 9 schools were chosen by systematic random sampling method. 35 samples from each school

were selected using simple random sampling method using the lottery method with the help of attendance registry maintained in the schools to arrive at the sampling frame. A total of 280 children, among which 150 were males and 129 were females. A pretested questionnaire was used as a study stool for the data collection. The validity of the tool was assessed by consultation with expert opinion. Anthropometry measurements were taken by Doctors through proper modes.

Neck circumference was measured using a plastic tape it was measured in a standing position while the child's head was held erect with eyes looking forward, shoulders in a relaxed position, Waist circumference measurements were measured using tape. Measured at the top of the hip bone, then brought the tape and measured all the way around the body, level with the belly button. Made sure it wasn't too tight and that it's straight, even at the back. asked the children not to hold their breath while measuring, measured the number on the tape right after they exhale. Hip circumference was measured at the maximum circumference of the Gluteal region.

Participants were also examined regarding height, weight, and BMI.

Height (cm): Standing height was measured using an audiometer and height rod. It was measured in cm; the participants were made to stand in an erect position.

Weight (kg): Standard weight monitoring device was used that was placed on a hard flat surface and checked for zero balance.

BMI: BMI was calculated by dividing weight in kg with the square of height in the meter as per the formula kg/M^2 .

The Data was entered in excel sheets and analysed using SPSS software version 21 [11].

3. RESULTS

In a total of 279 children, Table 1 shows mean anthropometric measurement values in boys. The mean weight, height, BMI, WC, HC, WHR were 71.54 ± 9.83 kg, 161.92 ± 7.53 cm, 27.70 ± 1.27 kg/m^2 , 91.07 ± 5.29 cm, 104.28 ± 4.63 cm, 0.87 ± 0.07 cm respectively in obesity / above the over weight boys. Table 2 shows mean anthropometric measurement values in Girls. The mean weight, height, BMI, WC, HC, WHR were 57.54 ± 14.77 kg, 150.87 ± 5.6 cm, 27.72 ± 2.12 kg/m^2 , 87.75 ± 4.57 cm, 97.58 ± 4.75 cm, 0.89 ± 0.054 cm respectively in obesity / above the over weight girls. All the anthropometric measures are significantly found to be higher in above the over weight/obesity children compare to their normal peers and higher in boys compared to girls.

The study showed that mean anthropometric measurement values in boys. The mean weight, height, BMI, WC, HC, WHR were 71.54 ± 9.83 kg, 161.92 ± 7.53 cm, 27.70 ± 1.27 kg/m^2 , 91.07 ± 5.29 cm, 104.28 ± 4.63 cm, 0.87 ± 0.07 cm respectively in obesity / above the over weight boys (Table 1).

The Table 2 shows mean anthropometric measurement values in girls. The mean weight, height, MI, WC, HC, WHR were 57.54 ± 14.77 kg, 150.87 ± 5.6 cm, 27.72 ± 2.12 kg/m^2 , 87.75 ± 4.57 cm, 97.58 ± 4.75 cm, 0.89 ± 0.054 cm respectively in obesity / above the over weight girls.

Table 1. Comparison of anthropometric measurements between controls and above the over weight / obesity in girls

Girl (mean)	Controls (normal / underweight)	Above the over weight-obesity	P-value
Age (years)	13.96 ± 2.08	15.16 ± 2.08	0.079
Weight in kg	42.75 ± 9.59	57.54 ± 14.77	0.005
Height in cm	152.91 ± 8.87	150.87 ± 5.6	0.275
BMI (kg/m^2)	18.10 ± 3.15	27.72 ± 2.12	<0.001
Hip circumference in cm	80.98 ± 9.29	97.58 ± 4.75	<0.001
Waist circumference In cm	70 ± 10.61	87.75 ± 4.57	<0.001
Waist hip ratio	0.85 ± 0.075	0.89 ± 0.054	0.028
Neck circumference in cm	29.79 ± 2.31	32.85 ± 1.927	<0.001

Table 2. Pearsons correlation (r) between the neck circumference and other anthropometric measurements

	NC-BMI (r)	NC-WC (r)	BMI – WHR (r)	P value
Boys	0.491	0.453	0.496	<0.01
Girls	0.708	0.694	0.300	<0.01

4. DISCUSSION

The present study conducted on children aged 11-17years had shown a notable relation between neck circumference and anthropometric measurement of obesity in both boys and girls.

Childhood obesity has become a paramount worry. Many studies had manifest expand unfavourable health issue of childhood obesity both short term besides long term consequences. Controlling the epidemic of childhood obesity, awareness prior precaution and treatment of childhood obesity are important priorities that need adequate diagnostic measures. Both patients and health practitioner must have tools with low cost, quick and easy to use for screening and monitoring [12-13].

The first person to suggest that different body morphology and type of fat distribution which is associated with health risk of obesity is Vague J.Hatipoglu et al also suggested neck circumference can be used for screening children with above the over weight and obesity and correlated positively between neck circumference and BMI, waist circumference. Nafiu et al conducted a study on 1102 children aged between 6-18yrs had shown that neck circumference correlated with age, BMI and waist circumference in both boys and girls and neck circumference cut off value were given according to age, optimal neck circumference cutoff indicative of high BMI in boys ranged from 28.5-39 cm, in girls 32.7-34.6 cm respectively. Similar results were also found in other studies by Roberta et al and Mozaffer et al. The present study results which conducted on 279 students aged between 11 to 17 years has shown that neck circumference correlated with BMI , weight , hip circumference ,waist circumference in both boys and girls .Optimal NC cut off values indicative of high BMI in boys ranged from 33.16-37.68 cm, in girls 30.92-34.77 cm respectively. Neck circumference is reliable and easy to measure compare to other indexes of adiposity.

5. CONCLUSION

The present study findings in children are consistent with other studies, therefore we

conclude that above the over weight/obesity children could be screened through neck circumference, through which we could identify the pre obesity children and carry out the needed management since there in increased level of obesity related health conditions among the children and the adolescents.

CONSENT AND ETHICAL APPROVAL

As per international standard or university standard guideline Patient’s consent and ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Available:https://watermark.silverchair.com/mxm012.pdf?token=AQECAHi208BE49Ooan9kkhW_Ercy7Dm3ZL_9Cf3qfKAc485ysgAAAkMwggI_BgkqhkiG9w0BBwagggIwMIICLAIBADCCAIUGCSqGSIb3DQEHATAeBgIghkgBZQMEAS4wEQQMstPXLIPfLMnxdtVAgEQgIIB9trW0cr1azeGMQIdVPsovYEJLE9W5w6SWXmzQG62GF-1PZIRBrbr-MnxCfiTu7-lv1eM_aSmkt8V3V37jGRWITl6NjZQ7S9UZSaHvv5jzGhRe406PdWhZzCk4QntZtObmIay6uncl6ZQzOQ2Ze9HSSjzhrXj-loT4VP8J2YzrQG
2. Gupta P, Tyagi S, Mukhija M, Saini AS, Goyal R, Sharma PL. Obesity: An introduction and evaluation. Journal of Advanced Pharmacy Education & Research. 2011;2:125-37.
3. Hatipoglu N, Mazicioglu MM, Kurtoglu S, Kendirci M. Neck circumference: an additional tool of screening above the over weight and obesity in childhood. European Journal of Pediatrics. 2010;169(6):733-9.
4. Naif OO, Burke C, Lee J, Voepel-Lewis T, Malviya S, Tremper KK. Neck circumference as a screening measure for identifying children with high body mass index. Pediatrics. 2010;126(2):e306-10.

5. Ben-Noun L, Sohar E, Laor A. Neck circumference as a simple screening measure for identifying above the over weight and obesity patients. *Obesity Research*. 2001;9(8):470-7.
6. Hingorjo MR, Qureshi MA, Mehdi A. Neck circumference as a useful marker of obesity: a comparison with body mass index and waist circumference. *JPMA- Journal of the Pakistan Medical Association*. 2012;62(1):36.
7. LaBerge RC, Vaccani JP, Gow RM, Gaboury I, Hoey L, Katz SL. Inter-and intra-rater reliability of neck circumference measurements in children. *Pediatric pulmonology*. 2009;44(1):64-9.
8. Yan Q, Sun D, Li X, Zheng Q, Li L, Gu C, Feng B. Neck circumference is a valuable tool for identifying metabolic syndrome and obesity in Chinese elder subjects: a community-based study. *Diabetes / Metabolism Research and Reviews*. 2014; 30(1):69-76.
9. Taheri M, Kajbaf TZ, Taheri MR, Aminzadeh M. Neck circumference as a useful marker for screening above the over weight and obesity in children and adolescents. *Oman Medical Journal*. 2016;31(3):170.
10. Whitlock, Anna E. Garcia, Jamarie Sostre-Colón, Matthew Gavin, Niels D. Martin, Joseph A. Baur, Carrie A. Sims, and Paul M. Titchenell. Loss of FOXO transcription factors in the liver mitigates stress-induced hyperglycemia. *Molecular Metabolism*. 2021;51:101246.
11. Assyov Y, Gateva A, Tsakova A, Kamenov Z. A comparison of the clinical usefulness of neck circumference and waist circumference in individuals with severe obesity. *Endocrine Research*. 2017;42(1): 6-14
12. Asif M, Aslam M, Wyszzyńska J, Altaf S, Ahmad S. Diagnostic performance of neck circumference and cut-off values for identifying overweight and obese pakistani children: a receiver operating characteristic analysis. *Journal of Clinical Research in Pediatric Endocrinology*. 2020;12(4): 366.
13. Valencia-Sosa, Evelyn, Clío Chávez-Palencia, Juan R. Vallarta-Robledo, Enrique Romero-Velarde, Alfredo Larrosa-Haro, Edgar Manuel Vásquez-Garibay, and César Octavio Ramos-García. Percentile reference values for the neck circumference of Mexican children. *Children*. 2021;8(5):407.

© 2021 Ponmalar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle4.com/review-history/73608>