



## **Health Benefits, Therapeutic and Pharmacological Properties of Moringa- A Review**

**N. A. Tamilselvi<sup>1\*</sup> and T. Arumugam<sup>1</sup>**

<sup>1</sup>*Department of Vegetable Crops, Horticultural College and Research Institute,  
Tamil Nadu Agricultural University, Coimbatore-641 003, India.*

### **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2018/37179

#### Editor(s):

(1) Othman Ghribi, Department of Biomedical Sciences, School of Medicine and Health Sciences, University of North Dakota, USA.

#### Reviewers:

(1) Voravuth Somsak, Walailak University, Thailand.

(2) Helen Saada, Egypt.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/37179>

**Review Article**

**Received 03 October 2017**  
**Accepted 09 January 2018**  
**Published 23 January 2019**

### **ABSTRACT**

Moringa yields at least four different edible parts viz., pods, leaves, flowers and seeds. Moringa leaves are the excellent source of protein,  $\beta$ -carotene, vitamins, A, B, C and E, riboflavin, nicotinic acid, folic acid, pyridoxine, amino acids, minerals and various phenolic compounds, phytochemicals and omega 3 and 6 fatty acids. The leaves of moringa are rich in palmitic and linolenic acids whereas the seeds are predominated by oleic acid. The leaves are used to combat malnutrition, especially among infants and nursing mothers in many developing countries, particularly in India, Pakistan, Philippines, Hawaii and many parts of Africa. Pregnant women and lactating mothers use moringa leaf powder to enhance their child or children's nourishment especially in underdeveloped countries where women are suffering from malnutrition. These natural plant phenolics present in moringa leaves could be a good source of antioxidants and antimicrobials for food and pharmaceutical industries. Perhaps using the multi-mix approach of food product development more food products could be developed especially for programs on malnutrition.

**Keywords:** *Moringa; Moringa oleifera Lam.; pharmacological properties; therapeutic uses.*

## 1. INTRODUCTION

*Moringa oleifera* Lam is the most widely cultivated multipurpose tree species of a monogeneric family, Moringaceae in which immature fruits, fresh leaves and flowers are used for the culinary purpose [1,2]. The plant is also known as Drumstick, Sahjan or Sohanjana in India [3] which has different vernacular names pertaining to each region and these were listed in Table 1. All plant parts of this tree have a remarkable range of functional, medicinal and nutraceutical properties [4,5]. In India and other countries, the tree is valued mainly for the tender pods which used as a vegetable [6], and it is used medicinally in Guinea, Madagascar, and Burma [7,8,9]. It is referred as a "multipurpose tree" "miracle tree" or a "wonder tree" [10,11,12] because of its several nutritional, pharmacological [11,13,14] and industrial applications [10,12,15,16]. The Moringa seeds found to exhibit natural coagulants/flocculants properties, which have potential to clear turbidity in drinking water and sludge in sewage respectively [17,18,19,20].

## 2. HEALTH BENEFITS

With four times the  $\beta$ -carotene of carrot, moringa has a unique potential for programs dealing with avitaminosis or hypovitaminosis syndrome known as vitamin A deficiency that causes 70 percent of childhood blindness. Presence of various types of antioxidant compounds makes this plant leaves a valuable source of natural antioxidants [21,22] and a good source of nutraceuticals and functional components as well [23]. Consumption of diet supplemented with moringa leaves could protect against diseases induced by oxidative stress. Many moringa nutritional supplements exist in the market including moringa dry leaf powder, capsules, nutrient shake and health booster. The moringa seed contains high-quality edible oil (up to 40% by weight). In Haiti, the oil has been used as general culinary and salad oil. It resembles olive oil in its fatty acid composition [24]. Hence, moringa plant is of great potential that could be cultivated as an economically profitable crop to contribute to poverty alleviation [25].

## 3. CULINARY USES

Moringa is consumed in diverse culinary preparations [26]. Almost all parts of the plant used for taste, flavour or as vegetables [27]. In

South India, pod are used to prepare a variety of sambars, curries with dals, fried curries and also used to add flavour to cutlets etc. In West Bengal and Bangladesh, it is used in a variety of dishes by mixing with coconut, poppy seeds or mustard and boiled until the moringa pods are semi-soft and consumed directly without any extra processing or cooking. In Maharashtra, the pods are used in sweet and sour curries called "Aamatee". Tender moringa leaves are finely chopped and used as a garnish for vegetable dishes, dals, sambar and salads [28]. The fresh succulent leaves are harvested daily for soups, sauces, or salads [25].

## 4. THERAPEUTIC AND PHARMACOLOGICAL PROPERTIES OF MORINGA

Besides rich nutritional value, moringa also has curative and prophylactic properties [29]. Almost all the parts viz., root, bark, gum, leaf, pods, flowers, seed and seed oil have been used for various ailments. Rajangam et al. [30] stated that the plant possesses rich pharmacological properties like cardia and circulatory stimulants, anti-tumour, anti-pyretic, anti-inflammatory, anti-spasmodic, diuretic, anti-hypertensive, anti-diabetic, hepato-protective, anti-bacterial and anti-fungal properties. Enormous research and development programmes were conducted on pharmaceutical and therapeutic properties of moringa and were summarized hereunder.

### 4.1 Anti-Microbial Activity

Caceres et al. [13] studied the antimicrobial activities of moringa leaves, roots, bark and seeds against bacteria, yeast, dermatophytes and helminths by a disk-diffusion method. The fresh leaf juice and aqueous extracts from seeds inhibit the growth of *Pseudomonas aeruginosa* and *Staphylococcus aureus*. They inferred that no activity was demonstrated against other pathogenic viz., Gram-positive, Gram-negative bacteria and *Candida albicans*. Juice from the stem bark showed an antibacterial effect against *S. aureus* [31].

Dahot [32] investigated the antimicrobial activity of three fractions of moringa leaf extract against *Escherichia coli*, *Klebsiella aerogenes*, *K. pneumoniae*, *S. aureus* and *Basillus subtilis* and observed that all the three fractions showed strong inhibitory activity against *E. coli*, *S. aureus* and *B. subtilis*. But clear zone of inhibition was noted against *K. aerogenes* and fraction 2

**Table 1. Vernacular names of morning pertaining to different region**

<b>Latin</b>	<b>: Moringa oleifera</b>
Sanskrit	: Danshamula, Shobhanjana, SigruShobhanjan, Sobhanjana, sigruh
Arabian	: Rawag
French	: Moringe à graineailée, Morungue
Spanish	: Ángela, Ben, Moringa
Portuguese	: Moringa, Moringueiro
Chinese	: La ken
English	: Drumstick tree, Horseradish tree, Radish tree, Ben oil tree, Mother's Best Friend, West Indian ben.
Tamil	: Morunga, Murungai, Murunkak-kai.
Telugu	: Mulaga, Munaga, Tella-Munaga, Sajana,
Kannada	: Guggala, mochaka, nugge, moxing
Malayalam	: Sigru, Moringa, Muringa, Murinna, Morunna
Punjabi	: Sainjna, Soanjna
Unani	: Sahajan
Ayurvedic	: Akshiva, Haritashaaka, Raktaka, Tikshnagandhaa
Hindi/ Orissa	: Munga ara, Shajmah, Shajna, Segra, Mungna, sahjan, saijna, sanjna, Soanjana, Soajna, Sohajna.
Gujarati	: Midho-saragavo, Saragavo, Saragvo, Suragavo, segto, seyla.
Bengali	: Munga ara, Sajina, Sajna, Sujana
Kanarese	: Nugga egipa, Nugge, Noogay, Nuggi Mara
Kol	: Mulgia, Munga ara, Mungna
Kumao – Himalayanregion	: Sunara
Konkani/Goa	: Moosing, Mosing
Marathi	: Sujna, Shevga, Shivga, Achajhada, shevgi
Modesia/W. Bengal	: Mangnai
Monghye/Punjab	: Sejana
Oriya	: Munigha, Sajina, Munika, Sojina, Sojaba
Punjabi	: Sanjna, Senjna, soanjna
Rajasthan	: LalSahinjano
Sindhi	: Swanjera
Teling	: Morunga, Morungai
Urdu	: Sahajna
Central provinces	: Mulaka, Saihan
Western region	: Sundan

[Source: Ram and Mehrotra, [117]; Roloff et al., [118]; Paliwal et al.[28]; Mishra et al., [106]; [www.moringanews.org/documents/VERNACULAR.doc](http://www.moringanews.org/documents/VERNACULAR.doc)][119]

showed a significant zone of inhibition against *Aspergillus niger*. Similarly, Amer et al. [33], Renitta et al. [34], Peixoto et al. [35] and Mbikay [36] stated that aqueous and ethanol moringa leaf extract could be a potential source for treatment against certain bacterial infection.

Since bacteria in water are attached to solid particles, treatment of water with moringa powder can remove bacteria up to 90 to 99% [37, 38, 39,40]. Additional treatment of water by boiling or adding chlorine is needed to render it completely safe to drink. Similarly, Shekhar et al. [41] tested the effect crude ethanol extract of moringa seed against *E. coli*, *Salmonella typhii*, *Vibrio cholera*,

*Shigella dysentriae* and *Pseudomonas aeruginosa* in drinking water and inferred that moringa seed extracts had antibacterial activity against *E. coli*. Arama et al. [42] also tested the moringa seed extract for antibacterial activity against *Escherichia coli* (ATCC 25922), *S. typhii* and *V. cholerae* (ref. Romel Cary Blair Lot. 452610) and indicated that *V. cholerae* was the most tolerant bacteria species to moringa extract as compared to *E. coli* and *S. typhii*.

Alam et al. [43] investigated antibacterial activity of moringa leaf extracts against four Gram-negative bacteria (*Shigella shinga*, *Pseudomonas aeruginosa*, *Shigella sonnei* and

*Pseudomonas* spp.) and six gram-positive bacteria (*Staphylococcus aureus*, *Bacillus cereus*, *Streptococcus-B- haemolytica*, *Bacillus subtilis*, *Sarcina lutea* and *Bacillus megaterium*) and inferred that leaf extract exhibited inhibitory effect against all the tested Gram-negative bacteria and Gram-positive bacteria except in *S. aureus* and *S. haemolytica*. Nantachit [44]; Doughari et al. [45] and Prashith et al. [46] reported similar anti-bacterial activity of moringa against certain bacteria. Nwosu and Okafor [47]; Nikkon et al. [48]; Chen et al. [49]; Jamil et al. [50] and Prashith et al. [46] reported antifungal activity of moringa leaf extract against seven pathogenic fungi using the broth dilution and agar plate methods.

#### 4.2 Anti-Inflammatory Activity

Medhi et al. [51]; Ndiaye et al. [52] and Sashidhara et al. [53] evaluated anti-inflammatory activity of methanol and aqueous extract of moringa root bark at a dose of 750 mg/kg and observed significant inhibition of oedema development at 1, 3 and 5 hours after treatment. Mahajan et al. [54,55] investigated anti-inflammatory activity from the ethanol extract of moringa seeds in toluene diisocyanate (TDI as antigen) induced asthma in Wistar rats and guinea pigs. An anti-fibrotic study conducted by Hamza [56] indicated that moringa seed extract possessed anti-inflammatory properties against CCL4 induced liver damage [57] and fibrosis. Paliwal et al. [28] stated that moringa seed oil has been used for various ailments in indigenous medicine of South Asia, including the treatment of inflammation and infectious diseases along with cardiovascular, gastrointestinal, haematological and hepatorenal disorders. Moringa leaves are therefore a potential source of natural antioxidants which were related to anti-inflammatory activity [13,14,48, 36].

#### 4.3 Anti-Oxidant Activity

Ashok Kumar and Pari [58] investigated antioxidant potential of moringa on hepatic marker enzymes, lipid peroxidation and antioxidants. The result of this study revealed that moringa extract and silymarin significantly decreased hepatic marker enzymes and lipid peroxidation with a simultaneous increase in the level of antioxidants. Bajpai et al. [59] tested the antioxidant activity of moringa leaves and inferred that kaempferol content is mainly responsible for this antioxidant property.

Siddhuraju and Becker [60] reported the antioxidant and free radical scavenging property of water, aqueous methanol and ethanol extracts of freeze-dried moringa leaves. The major bioactive compounds of phenolics were flavonoid groups such as quercetin and kaempferol. Various types of antioxidant compounds present in leaves and roots of moringa make this plant as a valuable source of natural antioxidants [21,22, 27,61,62,63,64,65,66] and a good source of nutraceuticals and functional components as well [15].

#### 4.4 Anti-Cancer Activity

Moringa has long been recognised by folk medicine practitioners as it has anti-tumour properties [67] and cancer prevention potential [68]. Murakami et al. [69] investigated the antitumor activity of moringa leaves and inferred that thiocarbamate (TC) and isothiocyanate (ITC) related compounds present in leaves are responsible for the anti-tumour activity. Similarly, Aruna and Sivaramakrishnan [70]; Guevara et al. [71]; Bharali et al. [72]; Costa et al. [73]; Parvathy and Umamaheshwari [74]; and Ahmad et al. [75] also reported the anticancer activity of ethanol extract of moringa seeds and leaves.

#### 4.5 Hepato-Protective Activity

Aqueous and alcohol extract of moringa flowers and roots possess hepatoprotective activity [76] which may be due to the presence of quercetin, a well-known flavonoid with hepatoprotective activity. Mazumder et al. [77] investigated haematological along with hepato-renal functions of methanol extract of moringa root in mice and concluded that high dose at daily treatment and moderate and high dose at weekly treatment with crude extract (CE) of moringa root increased WBC count and decreased clotting time significantly. Pari and Kumar [78] evaluated the hepatoprotective effect of ethanol extract of moringa leaves on liver damage induced by anti-tubercular drugs such as isoniazid (INH), rifampicin (RMP), and pyrazinamide (PZA) in rats and observed oral administration of the extract showed a significant protective action against hepatic disorders.

The hepatoprotective action of moringa seeds against Diclofenac (DIC) induced hepatic toxicity in male albino rats was investigated and the results of this study revealed that treatment with herbal extracts for 30 days before DIC treatment significantly reduces the indices of hepatotoxicity

induced by DIC [79]. Concomitant to this finding, Pal et al. [80]; Fakurazi et al. [81]; Hamza [56] and Paliwal et al. [28] also stated similar results.

#### 4.6 Cardiovascular Activity

The widespread combination of diuretic along with lipid and blood pressure lowering constituents makes this plant highly useful against cardiovascular disorders. According to Dahot [32], moringa leaf juice is known to have a stabilising effect on blood pressure. Faizi et al. [82,83 ad 84] isolated thiocarbamate and isothiocyanate glycosides from ethanol extracts of moringa leaves and inferred that these compounds are responsible for promising hypotensive activity. Gilani et al. [85] isolated four pure compounds viz., niazinin A, niazinin B, niazimicin and niazinin which showed a blood pressure lowering effect. Ghasi et al. [86] and Mehta et al. [31] investigated the hypocholesterolemic effect of crude leaf extract of moringa and observed significantly reduced serum cholesterol level but serum albumin level was increased by 15.22%.

Ara et al. [87] investigated the comparative effects of ethanol extracts of moringa leaves on serum cholesterol level, serum triglyceride level, blood glucose level, heart weight and body weight of adrenaline induced rats. The results of this study revealed that moringa leaves extract made significant changes in each cardiovascular parameter. Limaye et al. [88]; Mazumder et al. [77]; Nikkon et al. [48]; Ndong et al. [89]; Chumark et al. [27]; Nandave et al. [90]; Paliwal et al.[28]; Popoola and Obembe [91] and Abe and Ohtani [92] also reported the cardioprotective and hypotensive activity of ethanol extract of moringa leaves.

#### 4.7 Anti-Ulcer Activity

Pal et al. [80] reported that the moringa seed powder and leaves extracts have anti-ulcer and anti-gastritis activity. Moringa also has antibiotic activities against *Helicobacter pylori* which cause gastritis, gastric and duodenal ulcers [93]. Debnath and Guha [94] also reported the anti-ulcer effect of aqueous extract of moringa leaves on adult Holtzman albino rats of either sex using ondansetron as standard drug.

#### 4.8 Analgesic, Antipyretic and Wound Healing Activity

Medhi et al. [51] and Rao et al. [95] investigated the methanol extract of moringa root bark in mice

using acetic acid induced writhing method for analgesic activity. Rathi et al. [96] evaluated the wound healing property of aqueous extract of moringa leaves in male Swiss albino mice. Significant increase in wound closure rate, skin breaking strength, granuloma breaking strength, hydroxyproline content, granuloma dry weight and decrease in scar area was observed. Hukkeri et al. [97] investigated the antipyretic and wound healing activity of ethanol and ethyl acetate extracts of moringa leaves. The ethanol and ethyl acetate extracts of seeds showed significant antipyretic activity in rats; whereas ethyl acetate extract of dried leaves showed significant wound healing activity (10% extracts in the form of ointment) on excision, incision and dead space (granuloma) wound models in rats.

#### 4.9 Anti-Diabetic Activity

Suzuki et al. [98] studied the anti-diabetic effect of moringa leaves on glucose tolerance in Goto-Kakizaki and Wistar rats. Moringa leaf extract significantly decreased the blood glucose in Wistar rats. Jaiswal et al. [99] reported anti-diabetic activity of aqueous extract of moringa leaves on glycemic control, haemoglobin, total protein, urine sugar, urine protein and body weight. Ezeamuzie et al. [100] and Siddhuraju and Becker [60] indicated the anti-diabetic property of moringa. Hypo-cholesterolemic and hypoglycemic properties of moringa leaves were also reported by Siddiqui and Khan [101]; Ghasi et al. [86] and Dangi et al. [102].

#### 4.10 Diuretic and Antiurolithiatic Activity

Morton [103] and Caceres et al. [13] reported diuretic activity of hot water infusions of flowers, leaves, roots, seeds and stalks or bark of moringa. The diuretic components present in these plant parts play a complementary role in lowering blood pressure. Karadi et al. [104,105] reported antiurolithiatic property of aqueous and alcohol extract of moringa root bark and inferred that both the extracts significantly lowered the urinary excretion and kidney retention levels of oxalate, calcium and phosphate. Moreover, elevated serum levels of urea nitrogen, creatinine and uric acid were significantly reduced by these extracts.

#### 4.11 Other Diverse Pharmaceutical Activities

Moringa has been reported to exhibit other diverse beneficial activities. The plant has also

been used for the treatment of ascites, rheumatism [22], venomous bites [106] and CNS depressant [100,107,108]. The seed extract has been reported to be administered nasally to control the diseases like rhinitis and the dried seeds used successfully as an 'anti-allergic' agent by the Ayurvedic practitioners [109,110]. Additionally, the leaves have been reported for its radio-protective [111,112] and anthelmintic activity [113,114]. Moringa roots have been reported to possess anti-spasmodic activities [13,115,102] which helps for the management of gastrointestinal motility disorders. Aqueous leaf extract of moringa regulate "thyroid hormone" and can be used to treat hyperthyroidism [80,116].

## 5. CONCLUSION

Therefore, from this review it can be concluded that leaf, flower and pods of moringa could be very important for food, medicine and pharmaceutical industries. The moringa leaf powder become a vital ingredient of malnutrition programs.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

It is not applicable.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Dsouza J, Kulkarni AR. Comparative studies on nutritive values of tender foliage of seedlings and mature plants of *Moringa oleifera* Lam. J Econ Taxon Bot. 1993;17:479-485.
2. Oduro WI, Ellis O, Owusu D. Nutritional potential of two leafy vegetables: *Moringa oleifera* and *Ipomoea batatas* leaves. Sci Res Essays. 2008;3(2):57-60.
3. EISOhaimy SA, Hamad GM, Mohamed SE, Amar MH, Rashad RAH. Biochemical and functional properties of *Moringa oleifera* leaves and their potential as a functional food. Global Advanced Res J Agric Sci. 2015;4(4):188-199.
4. Jeng AS. Jatropha and Moringa: Two tropical multipurpose trees with potentials. Bioforsk. 2006;1(3):1-4.
5. Singh Y, Jale R, Prasad KK, Sharma RK, Prasad K. *Moringa oleifera*: A miracle tree. Proceedings, International Seminar on Renewable Energy for Institutions and Communities in Urban and Rural Settings, Manav Institute, Jevra, India. 2012;73-81.
6. Muthuswamy S. The culture of drumstick (*Moringa pterygosperma* (Gaertn.) syn. *Moringa oleifera* Lam in South Indian Hort. 1954;2:18-21.
7. Basu BD, Kirtikar KR. Indian medicinal plants. Second edn, Bishen Singh Mahendra Pal Singh, Dehradun. 1980;1:676-683.
8. Vaidyaratnam PSV. Indian medicinal plants-A compendium of 500 Species. Orient Longman Ltd., Madras. 1994;4:59-64.
9. Nadkarni AK. In: Indian Materia Medica, Popular Prakashan Pvt. Ltd., Bombay, 3rd Edition. 2002;1:811-816.
10. Jahn SAA. The traditional domestication of a multipurpose tree *Moringa stenopetala* (Bak.f.) Cuf. In the Ethiopian Rift Valley. AMBIO. 1991;20:244-247.
11. Fuglie L. The miracle tree: The multiple attributes of moringa. CTA Publication, Wageningen, The Netherlands. In combating malnutrition with moringa. Gopalan C., B.V. Rama Sastri & S.C. 2001;117-136.
12. National Research Council (NRC). Lost crops of Africa: Vegetables. 2006;2. Available:<http://www.nap.edu/catalog/11763.html>
13. Caceres A, Saravia A, Zabala L, Leon E. Pharmacologic properties of *Moringa oleifera*. 2: Screening for antispasmodic, anti-inflammatory and diuretic activity. J Ethnopharmacol. 1992;36:233-237.
14. Caceres A, Cabrera O, Morales O, Mollinedo P, Mendia, P. Pharmacological properties of *Moringa oleifera*. 1: Preliminary screening for antimicrobial activity. J Ethnopharmacol. 1991;33:213-216.
15. Makkar H, Becker K. Nutrients and anti-quality factors in different morphological parts of the *Moringa oleifera* tree. J Agri Sci. 1997;128:311-322.
16. Foidl N, Makkar HPS Becker K. The potential of *Moringa oleifera* for agricultural and industrial uses. In: The miracle tree: The multiple attributes of moringa; Fuglie,

- L.J., Ed.; Church World Service: Dakar, Senegal. 2001;45–76.
17. Gassenschmidt U, Klans D, Jarry D, Taucher B, Neebdergall H. Isolation and characterization of a flocculating protein from *Moringa oleifera* Lam. Biochem. Biophys. Acta. 1995;1243:477-481.
  18. Jahn SAA. Effectiveness of traditional flocculants as primary coagulants and coagulant aids for the treatment of tropical raw water with more than a thousand fold fluctuation in turbidity. Water Supp (2 Special Subject). 1984;6:8-10.
  19. Muyibi SA, Evanson LM. Optimizing physical parameters affecting coagulation of turbid waters with *Moringa oleifera* seeds. Water Res. 1995;29:2689-2695.
  20. Ndabigengesere A, Narasiah KS. Quality of water treated by coagulation using *Moringa oleifera* seeds. Water Res. 1998;32(3):781-791.
  21. Price M L. The moringa tree. ECHO Tech Note. North Fort Myers, USA. 2007.
  22. Anwar F, Latif S, Ashraf M, Gilani AH. *Moringa oleifera*: A food plant with multiple medicinal uses, Phytotherapy Res. 2007;21:17-25.
  23. Makkar HPS, Becker K. Nutritional value and anti-nutritional components of whole and ethanol extracted *Moringa oleifera* leaves. Animal Feed Sci Tech. 1996;63:211-228.
  24. Abdulkarim SM, Long K, Lai OM, Muhammad SKS, Ghazal HM. Some physio-chemical properties of *Moringa oleifera* seed oil extracted using solvent and aqueous enzymatic methods. Food Chem. 2005;93:253–263.
  25. Fatima AGH, Muna AI. *Moringa oleifera*: Nature is most nutritious and multi-purpose tree. Int J Sci Res. 2013;3(4):1-5.
  26. Iqbal S, Bhangar MI. Effect of season and production location on antioxidant activity of *Moringa oleifera* leaves grown in Pakistan. J Food Comp Anal. 2006;19:544-551.
  27. Chumark P, Khunawat P, Sanvarinda Y, Phornchirasilp S, Morales PN, Phivthongngam L, Ratanachamnong P, Srisawat S, Pongrapeeporn K S. The *in vitro* and *ex vivo* antioxidant properties, hypo-lipidaemic and anti-atherosclerotic activities of the water extract of *Moringa oleifera* Lam. leaves. J Ethnopharmacol. 2008;116:439-446.
  28. Paliwal R, Sharma V, Pracheta. A review on horse radish tree (*Moringa oleifera*): A multipurpose tree with high economic and commercial importance. Asian J Biotechnol. 2011;1-12.
  29. Ngugi GW. Country Reports: Kenya. Bambara groundnut *Vigna subterranea* (L.) Verdc. Promoting the conservation and use of underutilized and neglected crops 9. In: Heller J, egemann F, Mushonga J. Eds: Proceedings of the workshop on Conservation and Improvement of Bambara Groundnut (*Vigna subterranea* (L.) Verdc.). Harare, Zimbabwe; 1997.
  30. Rajangam J, Azahakia Manavalan RS, Thangaraj T, Vijayakumar A, Muthukrishan N. Status of production utilisation of moringa in Southern India, Division of Vegetable Sciences, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Tamil Nadu, India, this reference also used in “Le Moringa, *Moringa oleifera*, Un Arbre Tropical Aux Multiples Usages”, Phyto Magazine le magazine de Phytomania; 2001. Available:[http://www.phytomania.com/phyto/Moringa\\_oleifera.htm](http://www.phytomania.com/phyto/Moringa_oleifera.htm).
  31. Mehta LK, Balaraman R, Amin AH, Bafna PA, Gulati OD. Effect of fruits of *Moringa oleifera* on the lipid profile of normal and hypercholesterolaemic rabbits. J Ethnopharmacol. 2003;86:191–195.
  32. Dahot MU. Vitamin contents of flower and seeds of *Moringa oleifera* Lam. J Islamic Academy Sci. 1998;11(1):27-32.
  33. Amer J, Raheela J, Muhammad S, Muhammad A. Microscopic evaluation of the antimicrobial activity of seeds extracts of *Moringa oleifera*. Pak J Bot. 2008;40(4):1349-1358.
  34. Renitta RE, Nepolean P, Anitha J. Isolation, analysis and identification of phytochemicals of antimicrobial activity of *Moringa oleifera* Lam. Current Biotica. 2009;3(1):33-39.
  35. Peixoto JR, Silva GC, Costa RA. *In vitro* antibacterial effect of aqueous and ethanolic moringa leaf extracts. Asian Pac. J Trop Med. 2011;4:201-204.
  36. Mbikay M. Therapeutic potential of *Moringa oleifera* leaves in chronichyperglycemia and dyslipidemia: A review. Front Pharmacol. 2012;3:1-12.
  37. Schwarz D. Water clarification using *Moringa oleifera*. Technical Information W1e, Gate Information Service, Eschborn, Germany; 2000. Available:<http://www.gtz.de/gate/gateid.afp> (Accessed on 27th February, 2011)

38. Oloduro AO, Aderiye BI. Efficacy of *Moringa oleifera* seed extract on the microflora of surface and underground water. *J. Plant Sci.* 2007;2:453-458.
39. Amagloh FK, Benang A. Effectiveness of *Moringa oleifera* seed as coagulant for water purification. *Afr J Agric Res.* 2009;4:119-123.
40. Bukar A, Uba A, Oyeyi TI. Antimicrobial profile of *Moringa oleifera* Lam. extracts against some food-borne microorganisms. *Bayero J Pure Applied Sci.* 2010;3(1):43-48.
41. Shekhar C, Shukla R, Kumar A, Dubey, NK. Laboratory of herbal pesticides, Centre of advance study on botany, Banaras Hindu University, Varanasi, India. *Eur J Clin Microbiol Infectious Dis.* 2000;6:23-28.
42. Arama PF, Wagai SO, Ogur JA, Walter AO, Owido SO, Mahagayu CM. Harvesting surface rainwater – purification using *Moringa oleifera* seed extracts and aluminum sulfate. *J Agric Ext Rural Dev.* 2011;3(6):102-112.
43. Alam MF, Rahman MM, Sheikh MMI, Sharmin SA, Islam MS, Rahman A, Mizanur M, Rahman. Antibacterial activity of leaf juice and extracts of *Moringa oleifera* Lam. against some human pathogenic bacteria. *CMU J.* 2009;8(2):219-228.
44. Nantachit K. Antibacterial activity of the capsules of *Moringa oleifera* Lamk. Moringaceae. *CMU J.* 2006;5(3):365-368.
45. Doughari JH, Pukuma MS, De N. Antibacterial effects of *Balanites aegyptiaca* L. Drel. and *Moringa oleifera* Lam. on *Salmonella typhi*. *Afr J Biotechnol.* 2007;6(19):2212-2215.
46. Prashith KTR, Mallikarjun N, Swathi D, Nayana KV, Aiyar MB, Rohini TR. Antibacterial and antifungal efficacy of steam distillate of *Moringa oleifera* Lam. *Pharm Sci Res.* 2010;2(1):34-37.
47. Nwosu MO, Okafor JI. Preliminary studies on the antifungal activities of some medicinal plants against basidiobolus and some other pathogenic fungi. *Mycoses.* 1995;38(5-6):191-195.
48. Nikkon F, Saud A, Rahman MH, Haque ME. *In vitro* antimicrobial activity of the compound isolated from chloroform extract of *Moringa oleifera* Lam. *Pak J Biol Sci.* 2003;6(22):1888-1890.
49. Chen HM, Chuang PH, Lee CW, Chou JY, Murugan M, Shieh BJ. Anti-fungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. *Bioresour Technol.* 2007;98:232-236.
50. Jamil A, Raheela J, Muhammad S, Muhammad A. Microscopic evaluation of the antimicrobial activity of seed extracts of *Moringa oleifera*. *Pak J Bot.* 2008;40(4):1349-1358.
51. Medhi B, Khanikor HN, Lahon LC, Mohan P, Barua CC. Analgesic, anti-inflammatory and local anaesthetic activity of *Moringa* in laboratory animals. *Int J Pharmacogn.* 1996;34(3):207-212.
52. Ndiaye M, Dieye AM, Mariko F, Tall A, Diallo AS, Faye B. Contribution to the study of the anti-inflammatory activity of *Moringa oleifera* (Moringaceae). *Dakar Med.* 2002;47(2):210-212.
53. Sashidhara KV, Rosaiah JN, Tyagi E, Shukla R, Raghubir R, Rajendran SM. Rare dipeptide and urea derivatives from roots of *Moringa oleifera* as potential anti-inflammatory and antinociceptive agents. *Eur J Med Chem.* 2009;44(1):432-436.
54. Mahajan SG, Banerjee A, Chauhan BF, Padh H, Nivsarkar M, Mehta AA. Inhibitory effect of the n-butanol fraction of *Moringa oleifera* Lam. seeds on ovalbumin-induced airway inflammation in a guinea pig model of asthma. *Int J Toxicol.* 2009;28(6):519-527.
55. Mahajan SG, Mali RG, Mehta AA. Effect of *Moringa oleifera* Lam. seed extract on toluene diisocyanate-induced immune-mediated inflammatory responses in rats. *J Immunotoxicol.* 2007a;4(2):85-96.
56. Hamza AA. Ameliorative effects of *Moringa oleifera* Lam seed extract on liver fibrosis in rats. *Food Chem Toxicol.* 2010;48:345-55.
57. Rao KS, Mishra SH. Anti-inflammatory and anti-hepatotoxic activities of the rat of *Moringa pterygosperma* Geaerten. *Indian J Pharma Sci.* 1998;60:12-16.
58. Ashok Kumar N, Pari L. Antioxidant action of *Moringa oleifera* Lam. (drumstick) against antitubercular drugs induced lipid peroxidation in rats. *J Med Food.* 2003;6(3):255-259.
59. Bajpai M, Pande A, Tewari SK, Prakash D. Phenolic contents and antioxidant activity of some food and medicinal plants. *Int J Food Sci Nutr.* 2005;56(4):287-291.
60. Siddhuraju P, Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agro-climatic origins of drumstick



- tree (*Moringa oleifera* Lam.) leaves. J Agric Food Chem. 2003;51(8):2144-2155.
61. Sreelatha S, Padma PR. Antioxidant activity and total phenolic content of *Moringa oleifera* leaves in two stages of maturity. Plant Food Hum Nutr. 2009;64:303-311.
  62. Diallo AK, Gadegbeku TE, Mobio S, Moukha A, Agbonon K, Aklikokou EE, Creppy Gbeassor M. Protective effect of *Moringa oleifera* Lam and *Lannea kerstingii* extracts against Cadmium and Ethanol-induced lipid peroxidation. J. Pharmacol. Toxicol. 2009;4:160-166.
  63. Sultana B, Anwar F, Ashraf M. Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. Molecules. 2009;14(6): 2167-2180.
  64. Singh BN, Singh BR, Singh RL, Prakash D, Dhakarey R, Upadhyay G, Singh HB. Oxidative DNA damage protective activity, antioxidant and anti-quorum sensing potentials of *Moringa oleifera*. Food Chem Toxicol. 2009;1109-1116.
  65. Verma AR, Vijayakumar M, Mathela CS, Rao CV. *In vitro* and *in vivo* antioxidant properties of different fractions of *Moringa oleifera* leaves. Food and Chem Toxicol. 2009;47:2196-2201.
  66. Atawodi SE, Atawodi JC, Idakwo GA, Pfundstein B, Haubner R, Wurtele G, Bartsch H, Owen RW. Evaluation of the polyphenol content and antioxidant properties of methanol extracts of the leaves, stem, and root barks of *Moringa oleifera* Lam. J Med Food. 2010;13:710-716.
  67. Hartwell JL. Plants used against cancer: A survey. Lloydia. 1971;30-34.
  68. Fahey JW, Kostova ATD, Talalay P. The "Prochaska" microtiter plate bioassay for inducers of NQO1. *Methods in Enzymology*. (Eds) H.Sies and L. Packer, Elsevier Science, San Diego, CA. 2004;382:243-258.
  69. Murakami Y, Kitazono S, Jiwajinda K, Koshimizu, Ohigashi H. Niaziminin, a thiocarbamate from the leaves of *Moringa oleifera*, holds a strict structural requirement for inhibition of tumour promoter induced Epstein-Barr virus activation. Planta Med. 1998;64(4):319-323.
  70. Aruna K, Sivaramkrishnan VM. Plant products as protective agents against cancer. Indian J Exp Biol. 1990;28:1008-1011.
  71. Guevara AP, Vargas C, Sakurai H, Fujiwara Y, Hashimoto K, Maoka T, Kozuka M, Ito Y, Tokuda H, Nishino H. An anti-tumour promoter from *Moringa oleifera* Lam. Mutat Res. 1999; 440(2):181-188.
  72. Bharali R, Tabassum JM, Azad RH. Chemomodulatory effect of *Moringa oleifera* Lam, on hepatic carcinogen metabolizing enzymes, antioxidant parameters and skin papillomagenesis in mice. Asian Pac J Cancer Prev. 2003;4(2):131-139.
  73. Costa LLV, Khan MT, Ather A, Wilke DV, Jimenez PC, Pessoa C, De Moraes ME, De Moraes M O. Studies of the anticancer potential of plants used in Bangladeshi folk medicine. J Ethnopharmacol. 2005;99(1): 21-30.
  74. Parvathy MVS, Umamaheshwari A. Cytotoxic effect of *Moringa oleifera* leaf extracts on human multiple myeloma cell lines. Trends in Med Res. 2007;2(1):44-50.
  75. Ahmad FAR, Muhammad DI, Saie BK. Health benefits of *Moringa oleifera*. Asian Pac J Cancer Prev. 2014;15(20):8571-8576.
  76. Ruckmani K, Kavimani S, Anandan R, Jaykar B. Effect of *Moringa oleifera* Lam. On paracetamol-induced hepatotoxicity. Indian J Pharm Sci. 1998;60:33-35.
  77. Mazumder UK, Gupta M, Chakrabarty S, Pal DK. Evaluation of hematological and hepatorenal functions of methanolic extract of *Moringa oleifera* Lam. Root treated mice. Indian J Exp Biol. 1999;37:612-614.
  78. Pari L, Kumar NA. Hepato protective activity of *Moringa oleifera* on anti-tubercular drug induced liver damage in rats. J Med Food. 2002;5(3):171-177.
  79. Hamza AA. *Curcuma longa*, *Glycyrrhiza glabra* and *Moringa oleifera* ameliorate diclofenac-induced hepatotoxicity in rats. American J Pharmacol Toxicol. 2007;2(2): 80-88.
  80. Pal SK, Mukherjee PK, Saha BP. Studies on the antiulcer activity of *Moringa oleifera* leaf extract on gastric ulcer models in rats. Phytother Res. 1995a;9:463-465.
  81. Fakurazi S, Hairuszah I, Nanthini U. *Moringa Oleifera* Lam prevents acetaminophen induced liver injury through restoration of glutathione level. Food Chem Toxicol. 2008;46(8):2611-2615.

82. Faizi S, Siddiqui BS, Saleem R, Siddiqui S, Aftab K, Gilani AH. Fully acetylated carbamate and hypotensive thiocarbamate glycosides from *Moringa oleifera*. *Phytochem.* 1995;38:957-963.
83. Faizi S, Siddiqui BS, Saleem R, Siddiqui S, Aftab K, Gilani AH. Hypotensive constituents from the pods of *Moringa oleifera*. *Planta Med.* 1998;64(3):225-228.
84. Faizi S, Siddiqui BS, Saleem R, Siddiqui S, Aftab K, Gilani, and AH. Isolation and structure elucidation of new nitrile and mustard oil glucosides from *Moringa oleifera* and their effect on blood pressure, *J Nat Prod.* 1994;57(9):1256-1261.
85. Gilani AH, Aftab K, Suria A. Pharmacological studies on hypotensive and spasmodic activities of pure compounds from *Moringa oleifera*. *Phytother Res.* 1994;8:87-91.
86. Ghasi S, Nwobodo E, Ofili JO. Hypocholesterolemic effects of crude extract of leaf of *Moringa oleifera* Lam in high-fat diet fed Wistar rats. *J Ethnopharmacol.* 2000;69:21-25.
87. Ara N, Rashid M, Amran MS. Comparison of *Moringa oleifera* leaves extract with ethanol on serum triglyceride, serum cholesterol, blood glucose, heart weight, body weight in adrenaline induced rats. *Saudi J Bio Sci.* 2008;15(2):253-258.
88. Limaye DA, Nimbkar AY, Jain R, Ahmed M. Cardiovascular effects of the aqueous extract of *Moringa pterygosperma*. *Phytother Res.* 1995;9:37-40.
89. Ndong M, Uehara S, Katsumata, Sato S, Suzuki K. Preventive effects of *Moringa Oleifera* (Lam) on hyperlipidemia and hepatocyte ultra structural changes in iron deficient rats. *Biosci Biotechnol Biochem.* 2007;71:1826-1833.
90. Nandave M, Ojha SK, Joshi S, Kumari S, Arya DS. *Moringa oleifera* leaf extract prevents isoproterenol-induced myocardial damage in rats: evidence for an antioxidant, antiperoxidative and cardioprotective intervention. *J Med Food.* 2009;12(1):47-55.
91. Popoola JO, Obembe OO. Local knowledge, use pattern and geographical distribution of *Moringa oleifera* Lam. (Moringaceae) in Nigeria. *J Ethnopharmacol.* 2013;150:682-691.
92. Abe R, Ohtani K. An ethnobotanical study of medicinal plants and traditional therapies on Batan Island, the Philippines. *J. Ethnopharmacol.* 2013;145:554-565.
93. Harray X, Fahey J, Scholtus I, Lozniewski A. Evaluation of antimicrobial effects of several isothiocyanates on *Helicobacter pylori*. *Planta med.* 2005;71:326-330.
94. Debnath S, Guha D. Role of *Moringa oleifera* on enterochromaffin cell count and serotonin content of experimental ulcer model. *Indian J Exp Biol.* 2007;45:726-731.
95. Rao CV, Ojha SK, Mehrotra S. Analgesic effect of *Moringa oleifera* leaf extract on rats. In: *Proceedings of the Second World Congress on Biotechnological Developments of Herbal Medicine*, Lucknow, India. 2003;42.
96. Rathi BS, Bodhankar SL, Baheti AM. Evaluation of aqueous leaves extract of *Moringa oleifera* L. for wound healing in albino rats. *Indian J Exp Biol.* 2006; 44(11):898-901.
97. Hukkeri V I, Nagathan CV, Karadi RV, Patil BS. Antipyretic and wound healing activities of *Moringa oleifera* Lam. in rats. *Indian J Pharm Sci.* 2006;68:124-126.
98. Suzuki K, Ndong M, Uehara M, Katsumata S. Preventive Effects of *Moringa oleifera* (Lam) on hyperlipidemia and hepatocyte ultrastructural changes in iron deficient rats. *J Clin Biochem Nutr.* 2007;40:229-233.
99. Jaiswal D, Kumar Rai P, Kumar A, Mehta S, Watal G. Effect of *Moringa oleifera* Lam. leaves aqueous extract therapy on hyperglycemic rats. *J Ethnopharmacol.* 2009;123(3):392-396.
100. Ezeamuzie IC, Ambakederemo AW, Shode FO, Ekwebelm SC. Anti-inflammatory effect of *Moringa oleifera* root extract. *Int J Pharmacol.* 1996;34:207-212.
101. Siddiqui S, Khan MI. Pharmacological Study of *Moringa pterygosperma*. Central Laboratories. *Pak Council Sci Ind Res.* 1968;268-272.
102. Dangi SY, Jolly CI Narayanan S. Anti-hypertensive activity of the total alkaloids from the leaves of *Moringa oleifera*. *Pharm. Biol.* 2002;40:144-148.
103. Morton J F. The horse radish tree, *Moringa pterygosperma* (Moringaceae). A boon to arid lands. *Econ Bot.* 1991;45:318-333.
104. Karadi RV, Palkar MB, Gaviraj EN, Gadge NB, Mannur VS, Alagawadi KR. Anti-uro lithiatic property of *Moringa oleifera* root bark. *Pharmaceut Biol.* 2008;46(12):861-865.
105. Karadi RV, Gadge NB, Alagawadi KR, Savadi RV. Effect of *Moringa oleifera* Lam.

- root-wood on ethylene glycol induced urolithiasis in rats. *J Ethnopharmacol.* 2006;105:306–311.
106. Mishra D, Gupta R, Pant S, Kushwah K, Satish HT, Flora SJS. Co- administration of monoisoamyl dimercaptosuccinic acid and *Moringa oleifera* seed powder protects arsenic induced oxidative stress and metal distribution in mice. *Toxicol Mech Meth.* 2009;19:169-182.
107. Ray K, Hazra R, Guha. D. Central inhibitory effect of *Moringa oleifera* root extract: possible role of neurotransmitters. *Indian J Exp Biol.* 2003;41(11):1279-1284.
108. Ganguly R, Guha D. Alteration of brain monoamines and EEG wave pattern in rat model of Alzheimer's disease and protection by *Moringa oleifera*. *Indian J Med Res.* 2008;128(6):744-751.
109. Chopra RN, Chopra IC, Handa KL, Kapur ID. Eds: Chopra's indigenous drugs of India. UN Dhur and Sons, Calcutta, India. 1938;110.
110. Mahajan SG, Mali RG, Mehta AA. Protective effect of ethanolic extract of seeds of *Moringa oleifera* Lam. against inflammation associated with development of arthritis in rats. *J Immunotoxicol.* 2007b;4(1):39-47.
111. Rao VA, Devi PU, Kamath R. In vivo radioprotective effect of *Moringa oleifera* leaves. *Indian J Exp Biol.* 2001;39:858–863.
112. Arora R, Gupta D, Chawla R, Sagar R, Sharma A, Kumar R, Prasad J, Singh S, Samanta N, Sharma RK. Radioprotection by plant products: present status and future prospects. *Phytother Res.* 2005;19: 1–22.
113. Bhattacharya SB, Das AK, Banerji N. Chemical investigations on the gum exudates from Sonja (*Moringa oleifera*). *Carbohydr Res.* 1982;102:253-262.
114. Rastogi T, Bhutda V, Moon K, Aswar PB, Khadabadi SS. Comparative studies on anthelmintic activity of *Moringa oleifera* and *Vitex negundo*. *Asian J Research Chem.* 2009;2(2):181-182.
115. Gilani AH, Aftab K, Shaheen F. Antispasmodic activity of active principle from *Moringa oleifera*. In natural drugs and the digestive tract, Capasso F, Mascolo N (eds). EMSI: Rome. 1992;60–63.
116. Tahiliani P, Kar A. Role of *Moringa oleifera* leaf extract in the regulation of thyroid hormone status in adult male and female rats. *Pharmacol Res.* 2000;41:319–323.
117. Ram P, Mehrotra BN. Compendium of Indian medicinal plants. Vol. I, Central Drug Research Institute Lucknow and National Institute of Science communication, New Delhi. 2001;280.
118. Roloff A, Weisgerber H, Lang U, Stimm B. Enzyklopädie der Holzge wächse, Handbuch und Atlas der Dendrologie; WILEY-VCH: Weinheim, Germany; 2009.
119. Available:www.moringanews.org/document s/VERNACULAR.doc

© 2018 Tamilselvi and Arumugam; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/37179>