



Assessment of Genetic Divergence in White Finger Millet [*Eleusine coracana* (L.) Gaertn.] Genotypes for Yield and Yield Contributing Traits

Basavaraj^a, L. N. Yogeesh^{b++*} and T. M. Ningaraju^{c#}

^a College of Agriculture Raichur, University of Agricultural Sciences, Raichur, India.

^b Department of Genetics and Plant Breeding, College of Agriculture, Shimoga, Karnataka, India.

^c Department of Plant Biotechnology, College of Agriculture, GKVK, UAS, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i82247

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102632>

Original Research Article

Received: 20/04/2023

Accepted: 23/06/2023

Published: 24/06/2023

ABSTRACT

The present investigation is carried out to study the genetic divergence among 55 finger millet genotypes for fourteen quantitative characters using Mahalanobis D^2 statistics during *kharif*, 2020. D^2 statistics indicated that the genotypes studied were genetically diverse. The 55 genotypes of finger millet were grouped into 6 clusters irrespective of geographical diversity, indicating no parallelism between geographic and genetic diversity. Clusters I had highest number of 50 genotypes and remaining all clusters had solitary. A wider genetic diversity was observed for the different traits studied among the genotypes as evidenced by the formation of six clusters. Out of fourteen quantitative traits studied grain yield per plant contributed majorly towards divergence with the value of 15.65% followed by productive tillers per plant contributes (12%), harvest index (10%) and other traits contribute minorly for divergence.

⁺⁺ Assistant Professor (GPB);

[#] Assistant Professor;

*Corresponding author: E-mail: yogeeshhari@gmail.com, yogeeshhari@gmail.com;

Keywords: Genetic divergence; D^2 statistics; white finger millet.

1. INTRODUCTION

"Finger millet (*Eleusine coracana* L. Gaertn) is cultivated as a subsistence farming crop in Asian and African continents. Global area under finger millet is 4-4.5 million hectares with the production of 5 million tonnes" [1]. "Finger millet seeds are consumed in variety of forms such as unleavened bread (roti), mudde, thin or thick porridge, fermented porridge and also used in brewing. Demand for finger millet grains has picked up in the urban areas and baking industry in the recent times due to high fibre and other health benefits associated with its consumption. However, the dark colour of grains has been the major hindrance for its acceptability in baking and food industry. Among both brown and white grain types, white grain types are preferred because of high protein, low fibre, low tannins and higher consumer acceptability" [2]. "However, the yield potentiality of all white grain types is significantly lower than the brown types Ragi is commonly called as "Nutritious millet" as the grain is nutritionally superior to many cereals providing fair amount of proteins, minerals, calcium and vitamins. It contains almost all the nutrients like protein (9.2%), carbohydrates (76.32%) and fat (1.29%). It is very rich in minerals (2.70%) such as calcium (452 mg/1000g), iron (3.90 mg/100g) and ash (3.90 %) which are the core ingredients of normal human diet. The protein of finger millet has been reported to possess a fairly high biological value, which is needed for the maintenance of nitrogen equilibrium of the body. It has crude fiber content (3-4%) to supply energy for a long time after consumption and thus whole day sustenance, high cholesterol formation and intestinal cancer. Hence, people suffering from diabetics are advised to take finger millet and other small millets instead of rice" [3].

Genetic improvement through conventional breeding approaches depends mainly on the availability of the diverse germplasm and the amount of genetic variability present in the population. A method suggested by Mahalanobis [4] known as "Mahalanobis D^2 statistics" is "a powerful tool for quantifying the divergence between two populations. Therefore, the present study was undertaken to assess the nature and magnitude of genetic divergence for yield and its component in finger millet and also to identify divergent parents from distantly related clusters for suitable hybridization through genetic divergence analysis".

2. MATERIALS AND METHODS

The present investigation was consisted of evaluation of 55 white finger millet accessions for qualitative morphological descriptors at the Agricultural Research Station, Hagari, Ballari, Karnataka. The experiment was laid out in Randomized complete block design with one check variety (KMR 340) of white finger millet were obtained from the Indian Institute of Millets Research (IIMR), Hyderabad. Each entry was represented by two rows of 3-meter length. Present experiment consists of 11 blocks and one check which was randomly repeated in two replications. In each block, each genotype was sown in a single row with spacing of 30 cm between the rows and 10 cm between the plants within the rows. All recommended agronomical cultural practices were carried out to raise a good crop.

Observations were recorded based on five randomly selected plants in each genotype in each replication for fourteen important morphological characters viz., days to 50 per cent flowering, days to maturity, plant height (cm), flag leaf blade length (cm), peduncle length (cm), productive tillers per plant, number of fingers per ear, finger width (cm), finger length (cm), ear head length (cm), harvest index (%), 1000 seed weight (g) and grain yield per plant (g). The mean data of these five plants were utilized for the statistical analysis. The genetic divergence was computed using Mahalanobis (1936) D^2 statistics among all the fifty five genotypes. Based on genetic distance, all the genotypes were grouped in different clusters [5].

3. RESULTS AND DISCUSSION

D^2 statistics, a concept developed by Mahalanobis [6] is an important tool to plant breeder to classify the genotypes into different groups based on genetic divergence between them. The basic idea behind formation of clusters is to get the intra and inter-cluster distances. The serves as an index for selection of parents with diverse origin.

Clustering of genotypes following the Tocher's method as described by Rao [5]. The fifty-five genotypes of finger millet were grouped into six different clusters (Table 1). Clusters I had the highest number of 50 genotypes, clusters II, III, IV, V and cluster VI were solitary clusters. The formation of distinct solitary clusters may be due to the fact

that geographic barriers hindering gene flow, intensive natural and human selection for diverse and adaptable gene complexes must be responsible for this genetic diversity [7]. The mutual relationship between clusters is represented by considering average intra and inter-cluster Mahalanobis's D^2 distances in Table 2. The present observations were concurring with the reports of earlier workers like Patel et al. [8], Vara Prasad and Shivani [9], Ponsiva et al. [4].

Among the six clusters formed Cluster I (21.21) has the highest intra cluster distance however, all other 5 clusters had the value of 0.00 with no intra cluster Mahalanobis's D^2 distance due to solitary clusters.

Cluster I is found to have nearest value with cluster II (30.45), cluster III (31.22), cluster IV (34.54), cluster V (44.6) and cluster VI (47.41). Cluster II is found to have nearest value with cluster I (30.45), cluster IV (34.66), cluster VI (57.88), cluster III (62.66) and cluster V (91.51). Cluster III is found to have nearest distance value with cluster V (20.98), cluster I (31.22), cluster IV (32.2), cluster VI (62.44) and cluster II (62.66).

Cluster IV is found to have nearest distance value with cluster III (32.2), cluster I (34.54), cluster II (34.66), cluster V (45.85) and cluster VI (82.58). Cluster V is found to have nearest distance value with cluster III (20.98), cluster I (44.6), cluster IV (45.85), cluster II (91.51), cluster VI (99.09). Cluster VI is found to have

value of nearest distance with cluster I (47.41), cluster II (57.88), cluster III (62.44), cluster IV (82.58) and V (99.09). The present results were in confirmative with the report of earlier workers like Negi et al. [10], Geethanjali and Jagadeeswar [11] and Kaluthantri and Dasanayak [12].

Cluster group means for 14 characters are presented Table 3. Cluster I having 50 genotypes showed highest cluster mean for test weight. Cluster II having 1 genotype, exhibited highest cluster mean for harvest index and days to 50 per cent flowering. The monogenotypic cluster III resulting highest cluster mean for plant height, finger length, fingers per ear head, flag leaf blade length and ear head length. The monogenotypic cluster IV not recorded higher cluster mean for any of the characters. Cluster V, which was represented by 1 entry possessed highest cluster mean for productive tillers per plant, flag leaf blade length and grain yield per plant. Cluster IV had 1 genotype, exhibited highest cluster means for days to maturity, peduncle length and finger width. On the basis of above results it is evident that cluster III had maximum cluster means for most of desirable characters viz., plant height, finger length, fingers per ear head, flag leaf blade length and ear head length. Therefore, genotypes improvement of a large number of seed yield and yield contributing characters, simultaneously. Earlier worker Bedis et al., [13] and Sahu et al., [14] also reported wide variability among clusters for yield and most of the yield contributing characters.

Table 1. Clustering pattern of 55 genotypes of finger millet on the basis of genetic divergence

Clusters	No. of genotypes	Genotypes
I	50	E 319, E 325, IC0473986, IC0473924, IC0474226, E331, IC0474231, GPU67, IC0473978, IC0473970, IC0621996, IC0473987, IC0473948, IC0473957, IC0474219, OUAT 2, IC0473874, IC0473959, IC0473996, IC0474228, KMR 340, IC0474183, IC0473971, IC0474232, IC0474233, IC0473993, IC0473984, IC0473980, VL 352, IC0621993, ER 97, IC0474038, IC0474215, IC0473947, IC0473950, IC04022775, IC0474356, IC0473983, PR202, IC0474045, IC0474037, IC0474227, IC0474222, IC0473992, IC0474206, IC0475624, IC0473990, IC0474207, IC0065595, IC0474433.
II	1	IC0473972
III	1	IC0474065
IV	1	IC0473989
V	1	IC0474226
VI	1	IC0474044

Table 2. Intra and inter cluster distance D² values among 55 genotypes of finger millet

Clusters	Cluster distances					
	I	II	III	IV	V	VI
I	21.21	30.45	31.22	34.54	44.6	47.41
II	30.45	0	62.66	34.66	91.51	57.88
III	31.22	62.66	0	32.2	20.98	62.44
IV	34.54	34.66	32.2	0	45.85	82.58
V	44.6	91.51	20.98	45.85	0	99.09
VI	47.41	57.88	62.44	82.58	99.09	0

Table 3. Intra cluster group means for various components of 55 finger millet genotypes

Clusters	N	DFF	DM	PH	PTP	FL	FPE	1000SW	HI	FLBL	FLBW	PL	EHL	FW	GYP	Overall Score	Rank
I	50	72.03 (4)	104.72 (4)	65.5 (5)	2.26 (3)	7.23 (4)	6.71 (4)	3.23 (1)	29.39 (5)	35.67 (4)	0.79 (4)	16.21 (3)	9.95 (3)	0.78 (2)	21.17 (3)	49	3
II	1	79 (1)	102 (5)	67.2 (3)	2 (5)	5.9 (6)	6.5 (5)	3.03 (4)	32.1 (1)	33.8 (5)	0.67 (6)	16.4 (2)	9.8 (4)	0.72 (5)	17.7 (6)	58	2
III	1	74.5 (3)	112 (2)	74.9 (1)	2.6 (2)	9.5 (1)	9 (1)	3.2 (2)	29.6 (4)	36.5 (3)	0.87 (1)	15.2 (4)	10.7 (1)	0.75 (4)	22.7 (2)	31	6
IV	1	60 (6)	108 (3)	67.15 (4)	2.1 (4)	9 (3)	6.5 (6)	2.62 (6)	31.5 (2)	33.5 (6)	0.71 (5)	12.9 (6)	9.2 (5)	0.68 (6)	20.2 (4)	66	1
V	1	76.5 (2)	99 (6)	55.7 (6)	2.9 (1)	9.1 (2)	7.6 (2)	2.98 (5)	24.4 (6)	39.6 (1)	0.82 (3)	15 (5)	9.1 (6)	0.77 (3)	25.9 (1)	49	4
VI	1	65 (5)	118.5 (1)	71.4 (2)	2 (6)	7.2 (5)	7.6 (3)	3.19 (3)	31.3 (3)	36.8 (2)	0.85 (2)	16.7 (1)	10.3 (2)	1.59 (1)	18.9 (5)	41	5

Table 4. Contribution of different plant growth and grain yield characters to total divergence in 55 finger millet genotypes

Sl. No	Source	No. of first rank	Contribution %	Cumulative
1	Days to 50% flowering	45	3	3
2	Days to maturity	45	3	6
3	Plant height (cm)	30	2	8
4	Productive tillers per plant	179	12	20
5	Finger length (cm)	104	7	27
6	Fingers per ear head	144	9.63	36.63
7	1000 seed weight (g)	119	8	44.63
8	Harvest index (%)	149	10	54.63
9	Flag leaf blade length (cm)	75	5	59.63
10	Flag leaf blade width (cm)	60	4	63.63
11	Peduncle length (cm)	104	7	70.63
12	Ear head length (cm)	85	5.72	76.35
13	Finger width (cm)	119	8	84.35
14	Grain yield/plant (g)	234	15.65	100

3.1 Contribution of Individual Traits towards Divergence

The per cent contribution of all the studied fourteen traits towards divergence is computed in Table 4. The knowledge of how the character's influence divergence is the pivotal aspect to a plant breeder. The total contribution of various traits towards genetic divergence follows, grain yield (15.67%) showed maximum contribution towards genetic diversity followed by the other characters viz., productive tillers per plant (12%), harvest index (10%), fingers per ear head (9.63%), 1000 seed weight (8%), finger width (8%), finger length (7%), peduncle length (7%), ear head length (5.72%), flag leaf blade length (5%), flag leaf blade width (4%), days to 50 per cent flowering (3%), days to maturity (3%) and plant height (2%) contributed towards genetic divergence. The present findings were on par with the report of earlier workers like Shinde et al. [15], Jadhav et al. [16] for plant height, Ulaganathan and Nirmala kumari [17] and Babu et al. [18] for days to maturity in ragi, Brunda et al. [19] for days to maturity, Gangurde et al. [20] for grain yield per plant towards the maximum contribution to the divergence.

4. CONCLUSION

There is acute need of improve of yield in Finger millet as it is one of the main subsistence cereal crop in Africa and India. Finding out the variation for the trait, among the different genotypes is most important and use of different statistical tools like Mahalanobis D^2 is also impartment for selecting proper genotypes. In addition to yield trait the yield contributing traits like productive tillers per plant, harvest index, fingers per ear head, finger width, finger length, peduncle length, ear head length, flag leaf blade length and flag leaf blade width also contribute the yield. The results of the current study showed that ample genetic diversity existed among finger millet genotypes to be used in future breeding program through selection and hybridization.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Global strategy for the ex situ conservation of Finger millet and its wildrelatives; 2012.

Available: <https://www.croptrust.org/sites/default/files/documents/files/Finger%20Millet-final.pdf>.

2. Sharathbabu KS, Shanthakumar G, Salimath PM. Genetic variability and character association studies in white Ragi (*Eleusine coracana* L. Gaertn). Kar. J Agric Sci. 2008;21(4):572-5.
3. Malleshi NG, Hadimani NA. Nutritional and technological characteristics of small millets and preparation of valueadded products from them. In: Advances in Small millet Praceeding of Second International Small Millet workshop, Bulawayo, Zimbabwe; 1993.
4. Ponsiva ST, Sabarinathan S, Kirubhakaran R, Senthilkumar N, Anandan, Thirugnanakumar S. Mahalanobis D^2 and Gauch's AMMI analyses in medium duration rice genotypes grown over six seasons under coastal eco-system. PI. Archives. 2019;19(1):986-94.
5. Rao CR. Advance statistical methods in biometric research. New York: Wiley & Sons. 1952;390.
6. Mahalanobis PC. On the generalized distances in statistics. Proc. Nat. Inst., India. 1936;11(1):49-55.
7. Arunachalam V, Ram J. Geographical diversity in relation to genetic divergence in cultivated sorghum. Indian J Genet Plant Breed. 1967;27:369-80.
8. Patel SN, Patil HE, Popat RC. Genetic diversity study in finger millet (*Eleusine coracana* L.) Genotypes: A multivariate analysis approach. Int J Pure App Biosci. 2017;5(5):183-9.
9. Varaprasad BV, Shivani D. Genotype clustering of maize (*Zea mays* L.) germplasm using mahalanobis d statistic. J Glob Biosci. 2017;6:4776-83.
10. Negi S, Kumar V, Bhatt A. Genetic diversity among finger millet [*Eleusine coracana* (L.) Gaertn] genotypes for yield and its contributing traits. Int J Curr Microbiol Appl Sci. 2017;6(8): 3332-7.
11. Geethanjali S, Jagadeeswar M. Genetic diversity and variability in foxtail millet germplasm based on morphological traits. Electron J Plant Breed. 2016;7(2) :303-16.
12. Kaluthanthri DVS, Dasanayaka PN. Assessment of genetic diversity of some finger millet (*Eleusine coracana* (L.) Gaertn.) accessions using morphological markers. J Trop. 2016;6(2):25-35.

13. Bedis MR, Patil HS, Patil VS, Jangale GD. Genetic divergence in finger millet (*Eleusine coracana* G.). Natl J Plant Imp. 2007;9(1):58-9.
14. Sahu S, Pradhan K. Genetic divergence in finger millet (*Eleusine coracana* (L.) Gaertn). Environ Eco. 2012;30(2):291-4.
15. Shinde SR, Desai SV, Pawar RM. Genetic diversity pattern in finger millet (*Eleusine coracana* (L.) Gaertn). Electro J Plant Breed. 2013;4(3):1242-5.
16. Jadhav R, Babu DR, Lal AM, Rao VS. Assessment of genetic divergence in finger millet [*Eleusine coracana* (L.) Gaertn.] for yield and yield contributing traits. Int J Food Ferment Technol. 2014;4(2):113-20.
17. Ulaganathan V, Nirmalakumari A. Finger millet germplasm characterization and evaluation using Mahalobinis D² analysis. SABRAO J Breed Genet. 2015;47(2):79-88.
18. Babu KB, Sood S, Agrawal PK, Chandrashekara C, Arun K, Anil K. Molecular and phenotypic characterization of 149 finger millet accessions using microsatellite and agro-morphological markers. J Natl Acad Sci. 2017;87(4): 1217-28.
19. Brunda SM, Kamatar MY, Naveen KKL, Hundekar R, Sowmya HH. Evaluation of foxtail millet (*Setaria italica*) genotypes for grain yield and biophysical traits. J Bio Sci. 2015;4(5):2142-9.
20. Gangurde SS, Moharil MP, Jadhav PV, Wandhare MS, Gawai D, Dikshit N et al. Research article genetic divergence studies in foxtail millet (*Setaria italica* L.). Electron J Plant Breed. 2016;7(3): 514-9.

© 2023 Basavaraj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/102632>