



# Influence of Seaweed Sap and Organic Manures on Growth and Yield of Baby Corn

Amit Raj<sup>a++\*</sup>, Biswarup Mehera<sup>b#</sup> and Prateek Kumar<sup>a†</sup>

<sup>a</sup> Department of Agronomy, Naini Agricultural institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211007, Uttar Pradesh, India.

<sup>b</sup> Naini Agricultural Institute (NAI), Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211007, Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i102652

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104761>

Original Research Article

Received: 07/06/2023

Accepted: 11/08/2023

Published: 16/08/2023

## ABSTRACT

A field experiment was carried out during *Rabi* 2022 season at Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh, to study the response of Seaweed Sap and Organic Manures on Growth and Yield of Baby Corn. The treatments consists of three level of Seaweed sap (5%, 7.5% and 10%), Organic manures (FYM, Poultry manure, Vermicompost) and control. The experiment was laid out in Randomized Block Design (RBD) with ten treatments and replicated thrice. The soil of the experimental field was sandy loam with pH (8.0), EC (0.56 ds/m), Organic Carbon (0.62%), Available N (225 kg/ha), Available P (38.2 kg/ha), and Available K (240.7 kg/ha). The results showed that application of Seaweed sap (10%) + Vermicompost – 8.5t/ha recorded

<sup>++</sup> M. Sc. Scholar;

<sup>#</sup> Dean;

<sup>†</sup> Ph. D Scholar;

\*Corresponding author: E-mail: amitraj4639@gmail.com;

significantly higher plant height (167.58 cm), maximum dry weight (98.48 g), No. of cobs/plant (2.47), cob length (21.40 cm), Cob weight with husk (49.84 g/cob), Cob yield (11.03 t/ha) and stover yield (35.33 t/ha). Maximum gross returns (281216.67 INR/ha), net returns (216726.67 INR/ha) and Benefit Cost (B.C) ratio (3.36).

**Keywords:** *Baby corn; seaweed sap; growth; yield attributes; economics; FYM vermicompost; poultry manure.*

## 1. INTRODUCTION

“Maize (*Zea mays* L.) is the third most important cereal crop next to rice and wheat. Across the globe it is famous as queen of cereals since it possesses higher genetic yield potential among other cereal crops like rice, wheat, oat, millets etc. Its significance lies in the way that it is not only utilized for human consumption and animal feed but also it is utilized by the industries for the production of corn oil, corn starch etc. Baby corn is not a genetically dwarf maize as the name suggests it is the immature ear of normal maize. Baby corn ear are soft and consumed as vegetable by human being” [1]. “A recent trend is of growing maize for vegetable purpose, which is commonly known as ‘baby corn’. It is a small young cob or ear or the female inflorescence before pollination or fertilization. The important attributes relevant to baby corn are early maturity, synchronized ear emergence and small palatable yellow kernels” [2].

“FYM seems to act directly by increasing crop yield by acceleration of respiratory process or by cell permeability or by hormonal growth action. Under organic management, nutrients release and crop demand synchrony is very much required; hence, a thorough understanding of nutrients release pattern from organic sources is essential to avoid nutrients stress. FYM is a store house of nutrient, which contain all essential plant nutrients. It is beneficial as apply fertilizer like Zn and K in combination with FYM” [3].

Poultry manure is a readily available fertilizer that works well as a substitute for chemical-based fertilizers. Exchangeable cations increase after manure application, and the amount of nitrogen in the soil rises by nearly 53% (from 0.09 to 0.14%) Boateng et al. [4]. In agriculture, using poultry manure is primarily done to provide crops with nutrients and to alter the soil organically Warnen et al. [5]. Therefore, the goal of this study was to ascertain how applying poultry manure at various rates will affect the soil's

availability of phosphorus and the growth of the maize plant.

“Vermicomposting is the process of producing compost by utilizing earthworms to turn the organic waste into high-quality compost that consists mainly of worm cast in addition to decayed organic matter” [6].

Sea weeds are marine algae, saltwater dwelling, and simple organisms that fall into the rather outdated general category of “plants”. Most of them are the red (6000 species), brown (2000 species) or green (1200 species). A wide range of beneficial effects have been reported from the use of liquid seaweed extracts [7], including increased crop yields, resistance of plants to frost, increased uptake of inorganic constituents from the soil, more resistance to stress conditions and reduction in storage losses of fruit. “Sea weeds have proved effective in enhancing yield, pest and frost resistance in vegetables, fruits, flowers, cereals and pulses. Seaweed extracts had beneficial effect on seed germination and plant growth” [8]. The seaweeds are known to add major plant nutrients viz., N, P, K, Ca, Mg, S, and different micronutrients which are required for crop growth and development.

## 2. MATERIALS AND METHODS

This experiment was carried out during the *Rabi* season of 2022 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The crop research farm is situated at 25° 39' 42" N latitude, 81° 67' 56" E longitude and at an altitude of 98 m above mean sea level. The experiment was laid out in Randomized Block Design consisting of ten treatments with  $T_1$  Seaweed sap (5%) + FYM -15t/ha,  $T_2$  - Seaweed sap (5%) + Poultry manure – 3.5t/ha,  $T_3$  – Seaweed sap (5%) + Vermicompost - 8.5t/ha,  $T_4$  - Seaweed sap (7.5%) + FYM -15t/ha,  $T_5$  - Seaweed sap (7.5%) + Poultry manure – 3.5t/ha,  $T_6$  - Seaweed sap (7.5%) + Vermicompost – 8.5t/ha,  $T_7$  - Seaweed sap (10%) + FYM -15t/ha,

$T_8$  - Seaweed sap (10%) + Poultry manure – 3.5t/ha,  $T_9$  - Seaweed sap (10%) + Vermicompost – 8.5t/ha,  $T_{10}$  - Control (RDF). The soil of experimental field was sandy loam with pH (8.0), Organic carbon (0.62%), available N (225 kg/ha), available P (38.2 kg/ha), and available K (240.7 kg/ha). The crop was sown on 17<sup>th</sup> December 2022 with seed rate of 20 kg/ha. Seeds are sown at a spacing of 45 cm × 20 cm. Data recorded on different aspects of crop, viz., growth parameters and yield attributes were subjected to statistical analysis by “Analysis of Variance Technique” [9] and economic data were analyzed by mathematical method.

### 3. RESULTS AND DISCUSSION

#### A. Growth parameters

**Plant height (cm):** The significantly higher plant height (167.58 cm) was observed in treatment 9 with the application of Seaweed sap (10%) + Vermicompost – 8.5t/. However, treatment 4 with application of Seaweed sap (7.5%) + FYM - 15t/ha, treatment 5 with application of Seaweed sap (7.5%) + Poultry manure – 3.5t/ha, treatment 6 with application of Seaweed sap (7.5%) + Vermicompost – 8.5t/ha, treatment 7 with application of Seaweed sap (10%) + FYM -15t/ha and treatment 8 with application of Seaweed sap (10%) + Poultry manure – 3.5t/ha was statistically at par with treatment 9.

“Statistical analysis did not reveal any significant difference in the plant height of sweet corn as influenced by vermicompost. Thus, in this study, it is obvious that the highest rate of vermicompost at 5 t /ha produced the highest plant height (172 cm), although statistically insignificant” [10].

The study of Joshi et al., [11] revealed that vermicompost improved the plants height due to humic acid and growth promoting bacteria.

**Plant Dry Weight (g/plant):** The significantly higher dry weight (98.48 g) was observed in treatment 9 with the application of Seaweed sap (10%) + Vermicompost – 8.5t/. However, treatment 6 with application of Seaweed sap (7.5%) + Vermicompost – 8.5t/ha, treatment 7 with application of Seaweed sap (10%) + FYM - 15t/ha and treatment 8 with application of Seaweed sap (10%) + Poultry manure – 3.5t/ha was statistically at par with treatment 9.

Dadarwal et al. [12] results showed that maximum dry weight accumulation were recorded under 75% NPK from 2.25 tons vermicompost /ha. Similar result is also achieved by (Sharma et. al/2014). Use of organic fertilizers such as vermicompost has a positive effect on crop growth, yield.

#### B. Yield Attributes

**Number of cobs/plant:** The results revealed that there was significant difference between the treatments and maximum number of cob /plant (2.47/plant) was observed in treatment 9 with application of Seaweed sap (10%) + Vermicompost - 8.5t/ha which was significantly superior to all the treatments. Similar findings are reported by Naveen et al. [13].

**Cob length (cm):** The results revealed that there was significant difference between the treatments and maximum Cob length (21.40 cm) was observed in treatment 9 with application of Seaweed sap (10%) + Vermicompost - 8.5t/ha which was statically at par to all the treatment.

Younas et al. [14] revealed that “the application of vermicompost proved the most effective improving growth and yield attributes maize crop varieties. The highest values cob length were recorded with the application of vermicompost whereas, control treatment recorded the lowest values of these traits”.

**Cob weight (g):** The results revealed that there was significant difference between the treatments. Maximum Cob weight (49.84 g) was observed in treatment 9 with application of Seaweed sap (10%) + Vermicompost - 8.5t/ha which was statically at par to all the treatment.

“Individual cob weight in vermicompost received plots could be due to better interception, absorption and utilization of radiation energy leading to higher photosynthetic rate and finally more accumulation. The overall improvement reflected into better source- sink relationship, which in turn enhanced the yield and yield attributes” [15].

**Cob yield (t/ha):** The results revealed that there was significant difference between the treatments and maximum cob yield (11.03 t/ha) was observed in treatment 9 with application of Seaweed sap (10%) + Vermicompost - 8.5t/ha. However, treatment 6 with application of Seaweed sap (7.5%) + Vermicompost - 8.5t/ha

**Table 1. Influence of seaweed sap and organic manures on growth parameters and yield of baby corn**

S. No.	Treatment combinations	At 80 DAS	
		Plant height (cm)	Dry weight (g/plant)
1.	Seaweed sap (5%) +FYM -15t/ha	144.39	66.84
2.	Seaweed sap (5%) + Poultry manure – 3.5t/ha	148.78	69.83
3.	Seaweed sap (5%) + Vermicompost - 8.5t/ha	150.93	72.88
4.	Seaweed sap (7.5%) +FYM -15t/ha	156.31	78.04
5.	Seaweed sap (7.5%) + Poultry manure – 3.5t/ha	154.45	83.00
6.	Seaweed sap (7.5%) + Vermicompost - 8.5t/ha	156.83	85.58
7.	Seaweed sap (10%) +FYM -15t/ha	161.46	89.37
8.	Seaweed sap (10%) + Poultry manure – 3.5t/ha	163.11	93.41
9.	Seaweed sap (10%) + Vermicompost - 8.5t/ha	167.58	98.48
10.	Control (RDF 120:60:40 kg/ha)	147.20	64
	F-test	S	S
	SEm(±)	4.53	4.56
	CD (p=0.05)	13.46	13.55

**Table 2. Influence of seaweed sap and organic manures on yield attributes and yield of baby corn**

S. No.	Treatment combinations	No. of cobs/plant	Cob length (cm)	Cob weight (g/cob)	Cob yield (t/ha)	Stover yield (t/ha)
1.	Seaweed sap (5%) +FYM -15t/ha	1.67	18.93	44.16	7.50	28.33
2.	Seaweed sap (5%) + Poultry manure – 3.5t/ha	1.93	19.27	44.03	8.59	30.33
3.	Seaweed sap (5%) + Vermicompost - 8.5t/ha	2.00	18.73	45.04	10.10	32.66
4.	Seaweed sap (7.5%) +FYM -15t/ha	1.73	18.80	38.81	8.73	30.33
5.	Seaweed sap (7.5%) + Poultry manure – 3.5t/ha	2.07	18.87	40.83	9.44	32.66
6.	Seaweed sap (7.5%) + Vermicompost - 8.5t/ha	2.13	19.27	48.54	10.55	33.66
7.	Seaweed sap (10%) +FYM -15t/ha	2.00	18.47	40.28	7.97	29.00
8.	Seaweed sap (10%) + Poultry manure – 3.5t/ha	2.13	18.40	42.05	9.16	32.00
9.	Seaweed sap (10%) + Vermicompost - 8.5t/ha	2.47	21.40	49.84	11.03	35.33
10.	Control (RDF 120:60:40 kg/ha)	1.47	18.33	37.23	6.17	25.00
	F Test	S	S	S	S	S
	SEm (±)	0.07	0.30	0.18	0.16	0.59
	CD (P=0.05)	0.22	0.90	0.54	0.48	1.78

**Table 3. Effect of seaweed sap and organic manures on economics of baby corn**

<b>S. No.</b>	<b>Treatment combinations</b>	<b>Cost of Cultivation (INR/ha)</b>	<b>Gross Returns (INR/ha)</b>	<b>Net Returns (INR/ha)</b>	<b>B:C</b>
1.	Seaweed sap (5%) +FYM -15t/ha	66,240	1,91,750	1,25,510	1.89
2.	Seaweed sap (5%) + Poultry manure – 3.5t/ha	72,240	2,19,383.33	1,47,143.33	2.04
3.	Seaweed sap (5%) + Vermicompost - 8.5t/ha	63,990	2,57,483.33	1,93,493.33	3.02
4.	Seaweed sap (7.5%) +FYM -15t/ha	66,490	2,22,800	1,56,310	2.35
5.	Seaweed sap (7.5%) + Poultry manure – 3.5t/ha	72,490	2,40,983.33	1,68,493.33	2.32
6.	Seaweed sap (7.5%) + Vermicompost - 8.5t/ha	64,240	2,68,966.67	2,04,726.67	3.19
7.	Seaweed sap (10%) +FYM -15t/ha	66,740	2,03,683.33	1,36,943.33	2.05
8.	Seaweed sap (10%) + Poultry manure – 3.5t/ha	72,740	2,33,883.33	1,61,143.33	2.22
9.	Seaweed sap (10%) + Vermicompost - 8.5t/ha	64,490	2,81,216.67	2,16,726.67	3.36
10.	Control (RDF 120:60:40 kg/ha)	50,140	1,58,166.67	1,08,026.67	2.15

were found to be statistically at par with treatment 9.

Mohammadi et al. [16] reported that “the cob yield was significantly influenced by various levels of vermicompost. Significantly the highest cob yield (14.15t/ha) were recorded with the application of vermicompost at 4.00 t/ha followed by application of vermicompost at 2.00 t/ha. The increase in cob yield might be due to remarkable improvement in yield attributes. This might be due to better growth with higher nutrient supply by vermicompost and also provided nutrients for longer period and readily to use form with growth promoting substances which improve overall growth and reflected in yield”.

**Stover yield (t/ha):** The results revealed that there was maximum stover yield (35.33t/ha) observed in treatment 9 with application of Seaweed sap (10%) + Vermicompost - 8.5t/ha. Whereas, treatment 6 with application of Seaweed sap (7.5%) + Vermicompost - 8.5t/ha were found to be statistically at par with treatment 9.

This result was in accordance with the data recorded by Madhavi et al. [15]. Higher rate of nitrogen by organic manure had beneficial effect on physiological processes, plant metabolism, dry matter production, growth etc. there by leading to higher green fodder.

### C. Economics

**Maximum Gross Returns (INR/ha):** Maximum gross returns (2,81,216 INR/ha) was recorded in treatment T9: [Seaweed sap (10%) + Vermicompost - 8.5t/ha] whereas, the lowest value (1,58,166 INR/ha) was observed in treatment T10: Control plot [(RDF 120-60-40 kg/ha)].

**Net Returns (INR/ha):** Maximum net return (2,16,726 INR/ha) was recorded in treatment T9: [Seaweed sap (10%) + Vermicompost - 8.5t/ha] whereas, the lowest value (1,08,026 INR/ha) was observed in treatment T10: Control plot [(RDF 120-60-40 kg/ha)].

**Benefit Cost Ratio:** Maximum Benefit cost ratio (3.36) was recorded in treatment T9: [Seaweed sap (10%) + Vermicompost - 8.5t/ha], whereas the lowest value (1.89) was observed in treatment T1: [Seaweed sap (5%) +FYM - 15t/ha].

## 4. CONCLUSION

Based on the results, it is concluded that application of 10% seaweed sap and 8.5 t/ha of vermicompost has improved growth metrics, yield characteristics, and Benefit Cost (B:C) ratio.

## ACKNOWLEDGEMENT

The authors are thankful to Department of Agronomy, Naini Agricultural Institute, Prayagraj, Sam Higginbottom University of Agriculture Technology and sciences, (U.P) India for providing necessary facilities to undertaken the studies.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Jinjala VR, Virdia HM, Saravaiya NN, Raj AD. Effect of integrated nutrient management on baby corn (*Zea mays* L.). Agric. Sci. Digest. 2016;36(4):291-294.
2. Kumar S, Kalloo G. Attributes of maize genotype for baby corn production. Maize genetics News Letter. 1998;74.
3. Nawab K, Shah P, Arif M, Khan M, Ali K. Effect of cropping patterns, FYM, K and Zn of wheat growth and grain yield. Sarhad Journal of Agriculture. 2011;27(3):371-375.
4. Boateng S, Zickermann AK, Koraharens M. Effect of poultry manure on growth and yield of maize. West Africe J. App. Eco. 2006;9:1-11.
5. Warren JG, Phillips SB, Mullins GL, Keahey D, Peun CJ. Environmental and production consequences of using alumamended poultry litter as a nutrients source for corn. Journal Environmental Quality; 2006.
6. Devi J, Prakash M. Microbial Population Dynamics during Vermicomposting of three different substrates amended with cowdung. International Journal of Current Microbiology and Applied Sciences. 2015;4(2):1086- 1092.
7. Blunden G. Seaweed resources in Europe: Uses and potential. John Wiley & Sons, Chichester. 1991;65-81.
8. Thirumal Thangam R, Maria Victorial Rani S, Peter Marian M. Effect of seaweed liquid fertilizer on the growth and

- biochemical constituents of *Cyamopsis tetragonoloba* (L.) Taub. Seaweed Research and Utilisation. 2003;25(1&2): 99-103.
9. Gomez KA, Gomez AA. Three or more factor expects. In statistical Procedure for Agricultural Research. 1976;2:139-141.
  10. Villaver PJ. Response of sweet corn (*Zea mays* L. var.saccharata) to vermicompost and inorganic fertilizer application. International Journal of Humanities and Social Sciences. 2020;12(6):17-27.
  11. Joshi R, Singh J, Vig AP. Vermicompost as an effective organic fertilizer and biocontrol agent: Effect on growth, yield and quality of plants. Reviews in Environmental Scienceand Bio/Technology. 2015;14(1): 137–159.
  12. Dadarwal RS, Jain NK, Singh D. Integrated nutrient management in baby corn (*Zea mays* L.) Indian J. of Agic. Sci. 2009;79 (12):1023-1025.
  13. Naveen J, Das R. Yield performance of organic baby corn (*Zea mays* L.) as influenced by nutrient management and moisture conservation practices in sandy loam soils of Assam, Indian Journal of Agricultural Research. 2020;4(1):A-5292.
  14. Younas M, Zou H, Laraib T, Abbas W, Akhtar MW, Aslam MN. The influence of vermicomposting on photosynthetic activity and productivity of maize (*Zea mays* L.) crop under semi-arid climate, Journal. Pune. 2021;16(8):0256450.
  15. Madhavi BL, Reddy MS, Rao PC. Integrated nutrient management using poultry manure and fertilizers for maize, Ind. J. of Agron. 1995;40:1-4.
  16. Mohammadi NK, Patel KM. Influence of inorganic fertilizer, vermicompost and biofertilizer on yield & economic of sweet corn and nutrient status in soil, International Journal of Applied Research. 2017;3(5):183-186.

© 2023 Raj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/104761>