



Study of Correlation between Soil Chemical Properties, Zinc Fractions and Yield of Rice under Long Term Fertilization in Chromustert Soil

Chhaviraj Baghel ^{a++*}, Alok Tiwari ^{b#} and Vinay Bachkaiya ^{ct}

^a Department of Soil Science, CSK HPKV, Palampur, Himachal Pradesh- 176062, India.

^b College of Agriculture and Research Station, Saja, Bemetara, Chhattisgarh- 491993, India.

^c Department of Soil Science and Agricultural Chemistry, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh- 492012, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i224143

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109789>

Original Research Article

Received: 15/09/2023

Accepted: 21/11/2023

Published: 23/11/2023

ABSTRACT

A long-term experiment entitled "Study of correlation between soil chemical properties, zinc fractions and yield of rice under long term fertilization in Chromustert soil" was conducted at IGKV Raipur experimental farm during kharif 2019-20. This study comprised of ten treatments: control (no fertilizer), 50% NPK (sub-optimal dose), 100% NPK (optimal dose), 150% NPK (super optimal dose) and 100% NPK along with ZnSO₄ @ 10kg ha⁻¹, 100% NP, 100% N alone, 100% NPK along with farmyard manure @ 5t ha⁻¹, 50% NPK along with blue green algae @ 10 kg ha⁻¹ and 50% NPK along with green manure @ 40kg h⁻¹. The experiment was laid out at a randomized complete block

⁺⁺ Ph.D. Scholar;

[#] Dean;

[†] Scientist;

*Corresponding author: E-mail: chhaviraj.baghel007@gmail.com;

design which was replicated four times. The variations in Zn uptake by grain, straw and total (both grain and straw) varied from 17.29-129.73 g ha⁻¹, 27.19 to 196.54 g ha⁻¹ and 44.48 to 326.27 g ha⁻¹, respectively. Amongst various treatments 100% NPK+ ZnSO₄ had significantly higher values of Zn uptakes by grain, straw and total as compared to the control plot. The grain and straw yield of rice varied from 2430 to 6970 kg ha⁻¹ and 3110-8364 kg ha⁻¹, respectively with the application of various treatments of inorganic fertilizers and manures. From the statistical analysis it was observed that a correlation coefficient was computed between Zn fractions with respect to soil chemical properties and yield of rice that could possibly differ in imbalance and balance applications of varying doses of fertilizer. In this context, the significant correlation coefficient values (r) were obtained with all the parameters that showed organic carbon content in soil predominantly affected almost all individual fractions that ultimately maintain the pool of total Zn in soil for uptake by crop plants.

Keywords: Correlation; LTFE; uptake; yield; zinc fractions.

1. INTRODUCTION


Zinc (Zn) is the most important trace element for the growth of rice especially in submerged soil. Zinc deficiency are common in both tropical and temperate climates [1-3]. There is a critical need to enhance food production in developing (third-world) countries, where zinc shortages are most prevalent. Reducing zinc deficiency in the soil is necessary to meet crop productivity goals, but low Zn content in grains and straw is also low for humans and animals with inadequate Zn uptake. The bioavailability of zinc is associated with its transformation in soils and plants, which can be attributed to multiple mechanisms such as adsorption by clay surfaces, hydrous oxide minerals, organic matter, and other variables that impact crop uptake of zinc. To improve comprehension, successive fractionation systems can be used to separate total soil Zn into five mechanistic components [4]. These fractions consist of a water-soluble pool that is present in the soil solution, an exchangeable pool with ions bound to soil particles by electrical charges, an organically bound pool with ions adsorbed, chelated, or complexed with organic ligands, a pool of zinc non-exchangeably sorbed onto clay minerals and insoluble metallic oxides, and a pool of weathering primary minerals [5]. Understanding complex interactions between plant, soil, climate, management techniques, and their consequences on soil over extended time periods is a key goal of long-term fertilizer experimentation (LTFE). As a result, these studies have gained recognition as important examples of knowledge about sustainability, especially in the context of intensive farming [6]. The first goal is to determine and assess how input management affects the usage of chemical fertilizers, sustainable soil fertility, and subsequent crop productivity [7]. In post-harvest

soil, Zn applied in conjunction with organics (Farmyard manure or Vermicompost) greatly enhanced biomass yield and Zn uptake by Indian spinach and DTPA-extractable Zn. The combined application of zinc and organics (Farmyard manure or Vermicompost) had a good effect on the distribution of various zinc fractions [8]. It is crucial to accurately measure the various forms of zinc and their relationships to soil characteristics and crop Zn uptake in order to enhance farming profitability over the long run. Although the amount of zinc needed for crop development is significantly less than that of macronutrients, Zn deficiency in soil has typically been accounted for from diverse regions of the world. Therefore, the present investigation was carried out to "Study of correlation between soil chemical properties, zinc fractions and yield of rice under long term fertilization in Chromustert soil".

2. MATERIALS AND METHODS

2.1 Study Site Description

A long-term field experiment was conducted at the research farm of Indira Gandhi Krishi Vishwavidyalaya, Raipur during Kharif 2019-20. The site comes under the sub-humid zone, located at 22° 33' N to 21° 14' N latitude and 82° 6' E to 81° 38' E longitude with an altitude of 293m above Mean Sea Level which receives approximately 1200 – 1300 mm (51") of rainfall, mostly in the monsoon season from late June to early October. The soil of the study area classified under Vertisol order, having fine Montmorillonite clay also locally called as Kanhar and identified as Arang II series, with pH = 7.5, EC = 0.23 ds m⁻¹ and OC = 0.60%, available nitrogen 212.39, phosphorus 21.31 kg ha⁻¹, potassium 386.25 kg ha⁻¹ sulphur 25.98 kg ha⁻¹ and DTPA extractable zinc 1.2 mg kg⁻¹.



Map 1. Map showing study location

2.2 Experimental Details

The field experiment was carried out in randomized block design with ten treatments which were replicated four times. The ten treatments comprised of various inorganic and organic sources namely control (no fertilizer), 50% NPK (sub-optimal dose), 100% NPK (optimal dose), 150% NPK (super optimal dose) and 100% NPK along with ZnSO_4 @ 10 kg ha^{-1} , 100% NP, 100% N alone, 100% NPK along with farmyard manure @ 5 t ha^{-1} , 50% NPK along with blue green algae @ 10 kg ha^{-1} and 50% NPK along with green manure @ 40 kg ha^{-1} . The rice variety Rajeshwari R-1 was selected as a test crop and various organic sources applied in this experiment viz., FYM, green manure (*Sesbania aculeata*) and blue green algae. The recommended dose of NPK fertilizer for rice crop was 120:60:40 kg ha^{-1} , respectively. The N, P and K were applied through urea, SSP and MOP, respectively and nitrogen was applied in split doses. During the kharif season, sunhemp was grown as an in-situ green manure crop for 45 days. It was then chopped into small pieces (5-7 cm) and mixed into the soil using a power-operated rotavator during the puddling stage, prior to the transplantation of rice crops.

3. RESULTS AND DISCUSSION

1. Zinc uptake by rice crop

Nutrient intake was calculated by multiplying the nutrient content of the grain and grass by their respective yields. The total amount of grain and straw was considered to be total uptake of the rice crop. The data on uptake of zinc by rice crops is presented in Table 1. The variations in Zn uptake by grain, straw and total (both grain and straw) was varied from 17.29-129.73 g ha^{-1} , 27.19 to 196.54 g ha^{-1} and 44.48 to 326.27 g ha^{-1} , respectively. Amongst various treatments 100% NPK+ Zn had significantly higher values of Zn uptakes by grain, straw and total as compared to the control plot. On persual of data presented in table indicated that the highest Zn uptake by grain (129.73 g ha^{-1}), straw (196.54 g ha^{-1}) and total (326.27 g ha^{-1}) was recorded in 150% NPK, respectively. While, the lowest zinc uptake by grain (17.29 g ha^{-1}), straw (27.19 g ha^{-1}) and total (44.48 g ha^{-1}) was recorded in control plot. On the other hand Zn uptake by grain and straw in optimal dose of 100% NPK treatment was significantly lower than 100% NPK along with Zn and 100% NPK+FYM. The application of soil applied zinc (5 kg ha^{-1}) + one foliar spray

(0.5%) which lead to significant raise in Zn content of straw and seed and also total uptake. This may be because of that an increase in Zn concentration in the soil solution resulted in the absorption of nutrients from the soil solution and thus increased Zn content in the seeds and straw (Katyal et al. 1997). Since, total absorption is a function of the nutrient content of the seed and the grass and their proper yield, thus, leading to a significant increase in zinc absorption. These also be finding by [9]. The increase in Zn absorption is due to their additional mechanism can be attributed to changes in Zn availability in the root zone and their role in plant growth and development. More enzymatic and hormonal activity increase assimilates translocation in plant biomass which progress highest straw yield. Zn may have helped to stimulate vegetative growth due to the absorption of other essential nutrients influenced by the absorption of Zn. These findings are matched with Wijebandara and Iranie [10], Khan et al. [11] and Rajkumar et al. (2002). 62 Total dry weight and grain yield also increase by Zn under alternate wetting and drying regimes [12], reliable based on earlier research showing that applying Zn could considerably boost plant growth and grain output in both low and high-Zn status [13].

2. Effect of different inorganic and organic treatments on grain and straw yield of rice

This experiment was very important from the point of view in order to understand the complex interaction between soil, plants, climate, and management practice ultimately reflecting crop productivity. The data on grain and straw yield of rice is given in Table 2. The various treatments significantly influenced the grain and straw yield of rice.

The grain yield of rice varied from 2430 to 6970 kg ha⁻¹ and the straw yield of rice varied from 3110-8364 kg ha⁻¹ among different treatments of continuous application of chemical fertilizers and manures. The grain yield of rice was significantly influenced by the balanced application of nutrients alone as well as organic manures over the imbalanced application of nutrients and no fertilizer application. The highest grain yield (6970 kg ha⁻¹) of rice was obtained with the application of super optimal dose (150% NPK) which was significantly superior over all the treatments except 6890 (kg ha⁻¹) grain yield obtained in optimal dose (100% NPK + FYM), whereas, the lowest yield was recorded in control plot (2430 kg ha⁻¹). Similarly, zinc sulphate also increased the grain and straw yields in 100% NPK + ZnSO₄ over the sole application of nitrogen fertilizer and control. When compared to a control treatment on par with 150% NPK, the optimal dose of 100% NPK along with FYM demonstrated significantly greater yields. Only imbalance and/or sole application of nitrogenous fertilizer and control treatment was recorded lowest yields because of low fertility status of available nutrients.

3. Correlation between soil chemical properties, zinc fractions and yield of rice

From the statistical analysis it was observed that correlation coefficient was computed between Zn fractions with respect to soil chemical properties and yield of rice that could possibly be differ in imbalance and balance applications of varying doses of fertilizer. In this context, the significant correlation coefficient values (r) were obtained with all the parameters showed that organic carbon content in soil predominantly affected almost all individual fractions that ultimately

Table 1. Effect of various treatments on Zn uptake by rice

S. No.	Treatment	Zinc uptake (g ha ⁻¹)		
		Grain	Straw	Total
T ₁	Control	17.29	27.19	44.48
T ₂	50% NPK	48.87	61.59	110.46
T ₃	100% NPK	97.80	145.67	243.47
T ₄	150% NPK	129.73	196.54	326.27
T ₅	100% NPK + ZnSO ₄ @ 10kg ha ⁻¹	100.40	151.24	251.64
T ₆	100% NP	91.24	126.28	217.82
T ₇	100% N	31.20	44.06	75.26
T ₈	100% NPK + FYM @ 5 t ha ⁻¹	113.41	170.50	283.91
T ₉	50% NPK + BGA @ 10 kg ha ⁻¹	51.63	77.31	128.94
T ₁₀	50% NPK + Green Manure	62.43	90.20	152.63
	SEm (±)	2.74	5.95	8.31
	CD (α = 5% level of significance)	7.84	17.01	23.77

Table 2. Effect of fertilization on grain and straw yields of rice

S. No.	Treatment Name	Yield (kg ha ⁻¹)	
		Grain	Straw
T1	Control	2430	3110
T2	50% NPK	4580	5496
T3	100% NPK	6625	7950
T4	150% NPK	6970	8364
T5	100% NPK + ZnSO ₄ @ 10kg ha ⁻¹	6565	7878
T6	100% NP	6500	7800
T7	100% N	3535	4596
T8	100% NPK + FYM	6890	8268
T9	50% NPK + BGA	4600	5520
T10	50% NPK + GM	4920	5904
	SEm (±)	173	208
	CD (α = 5% level of significance)	496	594

maintain the pool of total Zn in soil for uptake by crop plants. This directly contributed to the available fraction in the root zone [14,15]. Moreover, the correlation study also revealed the direct contribution of Org-Zn fraction with available Zn [16,17]. The complexity of Zn availability in black soil [7] also suggested that there is a dynamic equilibrium of Org-Zn amongst the Zn fractions with the Zn availability in soil for plant utilization. Similar observations have been explained as well by Mishra *et al.* [18], Narwal *et al.* [14], and Obrador *et al.* [19].

The WSEX-Zn fraction tabulated in Table 3(a) and 3(b) showed a positive relationship with soil pH ($r=0.06$), positive as well as significant relationship with organic carbon ($r=0.66^*$), available Zn ($r=0.79^*$), grain ($r=0.72^*$), and straw yields ($r=0.73^*$) of rice. The WSEX-Zn fraction is of great importance in plant nutrition. Findings were also obtained by [20-22].

The Org-Zn fraction arranged in Table 3(a) and 3(b) demonstrated a positive and significant correlation with OC ($r=0.89^{**}$), available phosphorous ($r=0.82^{**}$), grain ($r=0.83^{**}$) and straw ($r=0.84^{**}$) yields, respectively. A positive and significant relationship with available nitrogen ($r=0.79^*$), available potassium ($r=0.61^*$) and available Zn ($r=0.64^*$) was also observed amongst different treatments. More exchange sites for the adsorption of Zn are provided by organic matter. Similar results were also observed by Prasad and Sakal [2]. The results are in conformity with that of Murthy *et al.* [23] and Mandal and Mandal [24].

The AMOX-Zn fraction in Tables 3(a) and 3(b) showed a negative relationship with soil pH ($r= -0.08$) a reduction in AMOX-Zn fraction with the

increase in pH can be explained by the natural reduction in oxide solubility and concentration as pH increases [25], and (Shiowatana *et al.* 2005). While a positive correlation with electrical conductivity ($r=0.16$) and a positive with a significant relationship with available Zn ($r=0.89^{**}$) was recorded amongst various treatments.

The CRYOX-Zn fraction tabulated in Tables 3(a) and 3(b) showed a negative relationship with soil pH ($r= -0.23$) and a positive and significant relationship with organic carbon ($r=0.48$) and available phosphorous ($r=0.56$). While a positive and significant relationship with available Zn ($r=0.86^{**}$) was observed with CRYOX-Zn. These results suggest that the use of organic fertilizers may play a major role in the availability of Zn in the soil system as a bound fraction [26].

The residual Zn fraction arranged in Table 3(a) and 3(b) showed a positive and significant relationship with OC ($r=0.81^{**}$), available nitrogen ($r=0.84^{**}$), available phosphorous ($r=0.86^{**}$), grain ($r=0.85^{**}$) and straw ($r=0.86^{**}$) yields, respectively and a positive and significant relationship with available potassium ($r=0.0.69^*$) was also observed. The increase in soil Zn availability with an increase in organic matter content has been demonstrated by the positive correlation between organic carbon and soil Zn availability [27]. The results were in conformity with the findings of Mishra *et al.* [18], Behra *et al.* [20] and Rupa *et al.* [28] who accounted that greater association between available Zn and sesquioxide explain the influence of oxides associated nutrient elements on the available pool of Zn.

Table 3(a). Effect of fertilization on correlation coefficient between soil chemical properties, yield of rice and Zn fractions (n = 40)

Parameters	Zn fractions at surface (0-15 cm)					
	WSEX-Zn	Org-Zn	AMOX-Zn	CRYOX-Zn	Residual Zn	Total Zn
Soil pH	0.06	0.31	-0.08	-0.23	0.23	0.20
EC	0.31	0.49	0.16	0.06	0.55	0.52
OC	0.66*	0.89**	0.59	0.48	0.81**	0.80**
Avail. N	0.71*	0.79*	0.54	0.42	0.84**	0.82**
Avail. P	0.75*	0.82**	0.64*	0.56	0.86**	0.85**
Avail. K	0.61*	0.61*	0.38	0.27	0.69*	0.67*
Avail. Zn	0.79*	0.64*	0.89**	0.86**	0.58	0.62*
Grain Y	0.72*	0.83**	0.63*	0.56	0.85**	0.84**
Straw Y	0.73*	0.84**	0.64*	0.56	0.86**	0.85**

*. Correlation is significant at the 0.05 level, **. Correlation is significant at the 0.01 level

Table 3(b). Effect of fertilization on correlation coefficient values between soil chemical properties, yield of rice and Zn fractions (n = 40)

Parameters	Zn fractions at sub soil (15-30 cm)					
	WSEX-Zn	Org-Zn	AMOX-Zn	CRYOX-Zn	Residual Zn	Total Zn
Soil pH	0.02	0.08	0.14	0.28	0.39	0.36
EC	0.33	0.33	0.41	0.49	0.52	0.51
OC	0.67*	0.68*	0.72*	0.78*	0.91**	0.90**
Avail. N	0.68*	0.73*	0.79*	0.85**	0.84**	0.84**
Avail. P	0.75*	0.76*	0.81**	0.76*	0.85**	0.85**
Avail. K	0.53	0.61*	0.64*	0.65*	0.62*	0.62*
Avail. Zn	0.91**	0.88**	0.85**	0.76*	0.73*	0.75*
Grain Y	0.74*	0.75*	0.79*	0.78*	0.85**	0.85**
Straw Y	0.74*	0.75*	0.80**	0.79*	0.86**	0.86**

*. Correlation is significant at the 0.05 level, **. Correlation is significant at the 0.01 level

The total Zn fraction in Table 3(a) and 3(b) showed a positive and significant relationship with organic carbon ($r=0.80^{**}$), available nitrogen ($r=0.82^{**}$), available phosphorous ($r=0.85^{**}$), grain ($r=0.84^{**}$) and straw ($r=0.85^{**}$) yields while a positive and significant relationship with available potassium ($r=0.67^*$) and available Zn ($r=0.62^*$) was also observed. This could probably be by reason of the fact that the OC directly involved in the availability of almost all fractions of Zn in soil system that directly contributed to pool of Zn [15,10] and prevent the conversion into a non-available form there by promoting the availability of Zn to a optimum level [22,29-39].

4. CONCLUSION

The yield and uptake of zinc by the rice crop are strongly influenced by the continuous application of organic manures and inorganic fertilizers. In comparison to the control plot, the 150% NPK (super optimum dose) treatment had significantly greater Zn uptake values by grain, straw, and overall. With the exception of grain yield attained in optimal dose of NPK with farm yard manure

(100% NPK + FYM), application of 150% NPK resulted in the highest grain yield of rice, which was significantly higher than all other treatments. Conversely, control had the lowest yield. Different zinc fractions showed a substantially positive correlation, and a long-term fertilizer experiment revealed that the residual Zn fraction alone greatly affected the production of rice.

ACKNOWLEDGEMENT

The authors are honored to acknowledge Dr. Alok Tiwari (Dean, College of Agriculture and Research Station, Saja, Bemetara, Chhattisgarh) for giving timely advice and technical support to completing this research by sponsoring the AICRP (All India Coordinated Research Project) on long term fertilizer experiment at Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Fageria NK, Dos Santos AB, Cobucci T. Zinc nutrition of lowland rice. *Communication in Soil Science and Plant Analysis*. 2011;42:1719–27.
2. Prasad R. Zinc in soils and in plant, human and animal nutrition. *Indian Journal of Fertility*. 2006;2:103–19.
3. Slaton NA, Normon RJ, Wilson CE. Effect of Zn source and application time on Zn uptake and grain yield of flood irrigated rice. *Agronomy Journal*. 2005;92: 272–78.
4. Jatav V. Vertical distribution of physico-chemical properties and zinc fractions of soil under different land use practices. M.Sc. (Ag.) thesis, Jawaharlal Nehru Krishi Vishwavidyalaya, Jabalpur
5. Alloway BJ. Zinc in soils and crop nutrition. Second edition published by IZA and IFA, Brussels, Belgium, Paris and France. 2008;59-74.
6. Singh VP, Pal B. Effect of Phosphorus and Zinc application on yield and nutrient composition of rice crop with water salinity. *Journal of Plant Development Sciences*. 2013;5(4):375-377.
7. Dwivedi AK and Dwivedi BS. Impact of long term fertilizer management for sustainable soil health and crop productivity: Issues and challenges. *JNKVV Res. J*. 2015;49(3):387-397.
8. Mourya, KK, Kumar S, Sharma VK, Yadav RK, Kumar V. Influence of integrated nutrient management on fractions of zinc and its uptake by Indian spinach (*Beta vulgaris* L.). *Communications in Soil Science and Plant Analysis*. 2023;1-8.
9. Suri D. Effect of chemical fertilizers, organics and lime on fractions of micronutrient cations under continuous maize-wheat cropping in an acid alfisol. Ph.D. (Ag.) thesis, Chaudhary Sarwan Kumar Himachal Pradesh Krishi Vishwavidyalaya, Palampur; 2023.
10. Wijebandara DMDI, Dasog GS, Patil PL and Hebbar M. Zinc fractions and their relationships with soil properties in paddy growing soils of northern dry and hill zones of Karnataka. *J. Indian Soc. Soil Sci*. 2011;59:141-147.
11. Khan P, Memon MY, Muhammad I, Depar N, Muhammad A, Memon MS, Shah JA. Determining the zinc requirements of rice genotype. *Sarhad Journal of Agriculture*. 2012;28(1).
12. Wang YY, Wei YY, Dong LX, Lu LL, Feng Y, Zhang J, Pan FS, Yang XE. Improved yield and Zn accumulation for rice grain by Zn fertilization and optimized water management. *Journal of Zhejiang University- SCIENCE B (Biomedicine & Biotechnology) ISSN*. 2014;1673-1581.
13. Wissuwa M, Ismail AM, Graham RD. Rice grain zinc concentrations as affected by genotype, native soil-zinc availability, and zinc fertilization. *Plant and Soil*. 2008;306:37-48.
14. Narwal RP, Kumar R, Antil RS. Long-term effect of farmyard manure and N on the distribution of zinc and copper fractions in soil under pearl millet-wheat cropping system. 19th World Congress of Soil Science, Soil Solutions for a Changing World, Brisbane, Australia; 2010.
15. Zhang YG, Zhang YY, Cai JP, Zhu P, Gao HJ and Jiang Y. Variation in available micronutrients in black soil after 30-year fertilization treatment *Plant Soil Environ*. 2014;60(9):387–393.
16. Dhiman S. Long-term effect of integrated nutrient supply on zinc dynamics in rice-wheat cropping system in an acid Alfisol of Himachal Pradesh. M.Sc. Thesis, CSK Palampur. 2007;147.
17. Dwivedi AK, Singh M, Kauraw DL, Wanjarri RH and Chauhan SS. Research bulletin on impact of fertilizer and manure use for three decades on crop productivity and sustainability and soil quality under Soybean-Wheat system on a Vertisol in Central India. IISS (ICAR), Bhopal. 2007;224-235.
18. Mishra P, Singh R, Srivastava PC, Bali R. Effect of continuous cropping and fertilization on zinc fractions and their contribution to plant uptake under rice-wheat system. *J.Indian Soc. Soil Sci*. 2009;57:167-171.
19. Obrador A, Alvares JM, Lopez-Valdivia LM, Gonzalez D, Movillo J and Rico MI. Relationship of soil properties with Mn and Zn distribution in acidic soils and their uptake by a barley crop. *Geoderma*. 2007;137:432-443.
20. Behera SK, Singh D, Dwivedi BS, Singh S, Kumar K and Rana DS. Distribution of fractions of zinc and their contribution towards availability and plant uptake of zinc under long-term maize (*Zea mays* L.) - wheat (*Triticum aestivum* L.) cropping on an inceptisol. *Australian Journal of Soil*; 2008.

21. Agbenin JO. Extractability and transformation of copper and zinc added to tropical savanna soil under long-term pasture. *Communications in Soil Science and Plant Analysis*. 2010;41:1016-1027
22. Kandali GG, Barua NG, Basumatary A, Medhi BK and Kurmi K. Zinc Fractions and their Availability to Wetland Rice in Acidic Soils of Assam. *J. Indian Soc. Soil Sci.* 2016;64:58-65
23. Murthy ASP. Zn fraction in wetland rice soils and their availability to rice. *Soil Science*. 1982;133(3):150-154
24. Mandal B, Haider BR and Mandal LN. 1986. Distribution of different forms of zinc in some rice growing soils. *J. Indian Soc. Soil Sci.*,34: 488-492
25. Ramzan, S., Bhat, M. A., Kirmani, N.A. and Rasool, R. 2014. Fractionation of zinc and their association with soil properties in soils of Kashmir Himalayas. *International Journal of Agricultural and Soil Science*, 2(8):132-142
26. El-Fouly MM, Fawzi AFA, Abou El-Nour EAA, Zeidan MS and Firgany AH. 2015. Impact of long-term intensive cropping under continuous tillage and unbalanced use of fertilizers on soil nutrient contents in a small holding village. *African Journal of Agricultural Research* 10(53):4850-4857
27. Shambhavi S. Dynamics of micronutrient cations in soil-plant system as influenced by long-term application of chemical fertilizers and amendments in an acid Alfisol. Ph.D. Thesis, CSK Krishi Vishvavidyalaya, Palampur. 2011;222.
28. Rupa TR, Srinivasa RC, Subba RA, Singh M. Effects of farmyard manure and phosphorous on zinc transformations and phyto availability in two Alfisols of India. *Bioresource Technology Oxford, U.K. Elsevier Science Limited*. 2003;87:279-288.
29. Verma G, Sharma RP, Sharma SP, Subehia SK and Shambhavi S. Changes in soil fertility status of maize-wheat system due to long-term use of chemical fertilizers and amendments in an alfisol. *Plant Soil Environ*. 2012;58(12):529–533.
30. Chaudhary SK, Singh NK. Effect of levels of nitrogen and zinc on grain yield and their uptake in transplanted rice. *Oryza*. 2007;44(1):44-47.
31. Jain NK, Dahama AK. Effect of phosphorus and zinc on yield, nutrient uptake and quantity of wheat (*Triticum aestivum*). *The Indian Journal of Agricultural Sciences*. 2007;77(5).
32. Jalali VK, Sharma MP. Status of available micronutrient cations in soils of mid-hill intermediate zone of Jammu region. *Indian J. Agril. Sci.* 2002;72(10):616 – 618.
33. Mandal LN, Mandal B. Zn fractions in soils in relation to zinc nutrition of lowland rice. *Soil Science*. 1986;142:141-148.
34. Safari M, Yasrebi J, Karimian N, Shan XQ. Evaluation of three sequential extraction methods for fractionation of zinc in calcareous and acidic soils. *Resources of Biology Science*. 2009;4:848-857.
35. Saini RK. Effect of zinc and iron on growth and yield of mothbean [*Vigna aconitifolia* (Jacq.) Marechal] in arid western Rajasthan. M.Sc. (Ag.) Thesis, Rajasthan Agricultural University, Bikaner; 2003.
36. Sakal R, Sinha RB, Singh AP, Bhogal NS. Evaluation of method and time of zinc application on rice. *Journal of Indian Society of Soil Science*. 1993;41:195-196.
37. Sharma BD, Aggarwal VK, Mukhopadhyay SS and Arora H. Micronutrient distribution and their association with soil properties in Entisols of Punjab. *Journal Agricultural Science*. 2002;72:334-340
38. Sharma, Kavita, Jain KK. Effect of zinc and thiourea on growth, yield and quality of clusterbean. *Annals of Agricultural Research*. 2004;25(1):169-171.
39. Verma S, Subehia SK. Zinc availability in an acid Alfisol as influenced by long-term cropping in a wet temperate zone of western Himalayas. *Agropedology*. 2005; 15:95-99.

© 2023 Baghel et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109789>