



Thyrotoxic Valvulopathy in a 7-year-Old Child: Case Report and Literature Review

**Ujunwa FA^a, Ohuche Ijeoma^{a*}, Eke CB^a, Chikani U^a
and Chinawa JM^a**

^a Department of Paediatrics, College of Medicine, University of Nigeria, Enugu Campus, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author CJM conceived and designed this study while authors UFA, OI, ECB and CU helped in critical revision of the article. All authors have read and approved the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CA/2024/v13i1396

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112389>

Case Study

Received: 04/12/2023
Accepted: 08/02/2024
Published: 13/02/2024

ABSTRACT

Introduction: Children with thyrotoxicosis may present rarely with valvulopathies. This disease is a syndrome complex of hyperdynamic status due to systemic manifestation of elevated circulating thyroid hormone levels.

Case Reports: O.C, a 7-year-old female child presented with abdominal pain that started seven days ago, passage of loose stool that started five days ago and weight loss that started five days ago. Salient findings at presentation were tachycardia, anterior neck swelling that moves with swallowing with a grade 3/6 pan-systolic murmur maximal at the apex. A diagnosis of hyperthyroidism at risk of thyrotoxicosis was made based on supported laboratory evidence. Echocardiography done showed severe mitral regurgitation. She is on Tabs Carbimazole and Propranolol. She has improved remarkably and has been on follow up.

*Corresponding author: E-mail: ijeomaohuche@gmail.com;

Conclusion: Hyperthyroidism affects multiple organ systems in the body and the heart is not spared. Most of the cardiac manifestations are functional and occur as a result of the hyperfunctioning gland. A high index of suspicion is required, as many of the patients present with only the consequences of hyperthyroidism.

Keywords: Valvopathy; echocardiography; murmurs; hyperthyroidism; child.

1. INTRODUCTION

Hyperthyroidism is a disease with multisystemic manifestations [1]. The heart is an organ which is highly susceptible to thyroid hormones and therefore cardiac manifestations are observed even in subclinical cases of hyperthyroidism [2]. Thyroid hormones have a significant impact on cardiac function and structure, with an attendant de novo cardiac condition or could uncover an underlying cardiac pathology [2]. The cardiovascular manifestations of hyperthyroidism include tachycardia, palpitations, bounding peripheral pulses, wide pulse pressure, hyperactive precordium, and murmurs [3]. The most common cardiac complications of hyperthyroidism are arrhythmias (mainly atrial fibrillation), valvulopathies, heart failure and hypertension [3]. From a US based study, the incidence of pediatric hyperthyroidism and thyrotoxicosis is about 0.44 cases per 1000 population for children aged 0-11 years and 0.59 cases per 1000 for those aged 12-17 years, therefore showing increasing incidence throughout childhood [4]. Despite the cardiovascular symptoms and effects being one of the major clinically significant features of thyrotoxicosis, there is paucity of data on the true incidence of thyrotoxic heart disease in children compared to adults. No data on the incidence or prevalence of thyrotoxic valvulopathy in children has been reported. This rarity necessitated the presentation of the case reports.

2. CASE PRESENTATION

O.C, a 7-year-old female presented with abdominal pain that started seven days ago, passage of loose stool that started five days ago and weight loss that started five days ago. She was then taken to a Teaching Hospital where some investigations and medications were given with persistence of symptoms necessitating her return to Nigeria. She was then referred to UNTH from a private hospital at Nsukka. She was first seen by the Gastroenterology team where she had a colonoscopy done. Following an incidental finding of a systolic murmur at the apex, she was

referred to the cardiology team for further evaluation. She is the first child in a monogamous family setting with 4 children. Other siblings are alive, with no history of similar complaints. Parents are of the lower socio-economic class. Salient findings at presentation were dyspnoea on exertion, easy tiredness, palpitation and sweating, tachycardia, anterior neck swelling that moves with swallowing, spherical in shape, not tender, no differential warmth and smooth to touch, wide pulse pressure, first and second heart sounds with a grade 3/6 pan-systolic murmur maximal at the apex.

A diagnosis of hyperthyroidism at risk of thyrotoxicosis was made. Complete blood count and serum electrolyte, urea and creatinine showed normal parameters. Thyroid function test (TFT) done showed elevated Free T3 and T4 of 36.05 and 61.34pmol/L respectively and low TSH of 0.010LmiU/L. Thyroid USS showed diffuse enlargement of the thyroid gland and isthmus with increased echogenicity and heterogeneous texture. Echocardiography done showed severe mitral regurgitation. She is currently being managed by the Paediatric Endocrinology team, and is on Tabs Carbimazole and Propranolol. Awaiting repeat of thyroid function tests and Echocardiography. She has improved remarkably and now on follow up.

3. DISCUSSION

Children with thyrotoxicosis may present rarely with valvulopathies as seen in our index case. Birrell [5] et al reported that children with undiagnosed thyrotoxicosis in the northern region of England have initially been referred to cardiologists with a heart murmur, gastroenterologists with diarrhoea and failure to thrive, as well as to psychiatric/psychology services because of challenging behaviour and school refusal. Griffith [6] et al also reported a 16-year-old girl with complaint of chest pains, palpitations and a new onset heart murmur with bilateral exophthalmos and thyromegaly.

There may be a link between race and mitral regurgitation in children with thyrotoxicosis. For instance, Lester [7] et al studied 18 hyperthyroid children (9 black and 9 white), six out of the 9 black children had findings of mitral regurgitation while none of the white children had MR. Our index patient is of a black race [7].

There may be some genetic link with thyrotoxic valvopathy. A study done in Canada by Stefani Doucette [8] et al of a 36-year woman who had been treated for thyrotoxicosis in pregnancy and was eventually delivered of a neonate who had features of thyrotoxicosis. Further examination revealed a systolic murmur at the lower sternal border and echocardiography finding of moderate to severe tricuspid valve regurgitation. We could not follow up the siblings and parents of our index patient for financial reasons [8].

The clinical features range from an asymptomatic presentation to overt signs of cardiac decompensation as seen in our patient. In a study by Ertek [9], the most frequent clinical features were palpitation, tachycardia, exercise intolerance, exertional dyspnoea and orthopnoea. Saxena et al reported Tachycardia, elevated blood pressure with wide pulse pressure, precordial apical systolic murmur. It could also be asymptomatic.

The investigations for diagnosing a hyperfunctioning thyroid gland; Echocardiography and ECG are important for the diagnosing CVS pathology in thyrotoxicosis. Echocardiography findings range from demonstrating regurgitant flow in the valves (especially mitral valve) to structural changes in the valve (prolapse) [2]. This was also reported in our index case ECG can show fibrillatory waves indicative of atrial fibrillation. In the index case, the mitral regurgitation could be from the functional and hyperdynamic effects of the thyrotoxicosis as there were no further lesion of the mitral valve and this is always reversible after treatment [10,11]. Treatment of thyrotoxic valvopathy include the use of diuretics and anti-failure medications and rarely valve replacement surgeries [1-3]. The prognosis is good especially in the paediatric population Most of the valvular manifestations are "functional". They resolve entirely or become less problematic with treatment of the hyperthyroidism Resolution is within weeks to months of starting the antithyroid medications [2-4]. The index case has improved remarkably and has been on follow up.

4. CONCLUSION

Hyperthyroidism affects multiple organ systems in the body and the heart is not spared. Most of the cardiac manifestations are functional and occur as a result of the hyperfunctioning gland. A high index of suspicion is required, as many of the patients present with only the consequences of hyperthyroidism.

FUNDING

This study was not funded by any organization. We bore all the expenses that accrued from in study.

ETHICAL APPROVAL

Not applicable.

CONSENT TO PARTICIPATE

Patients and parents or caregivers were duly informed in detail about the disease.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Elvan-Tüz A, Ayrancı İ, Ekemen-Keleş Y, Karakoyun İ, Çatlı G, Kara-Aksay A, Karadağ-Öncel E, Dündar BN, Yılmaz D. Are Thyroid Functions Affected in Multisystem Inflammatory Syndrome in Children? *J Clin Res Pediatr Endocrinol.* 2022;14(4):402-408. DOI: 10.4274/jcrpe.galenos.2022.2022-4-7. Epub 2022 Jun 30.
2. Nijith L, Ranjan R. Cardiovascular Manifestations in Hyperthyroidism: A Cross-Sectional Study in a Tertiary Care Hospital in South India. *Cureus.* 2022;14(5):e25232. DOI: 10.7759/cureus.25232. PMID: 35755496; PMCID: PMC9217679.
3. Pediatric Hyperthyroidism: Background, Pathophysiology ... Medscape Available: <https://emedicine.medscape.com/921707-overview>
4. Bhattad PB, Roumia M. Cardio-Thyrotoxicosis Syndrome: A Review of

- Thyrotoxic Cardiovascular Disease. *Cureus*. 2023;15(4):e37659. DOI: 10.7759/cureus.37659. PMID: 37200647; PMCID: PMC10188272.
5. Birrell G, Cheetham T. Juvenile thyrotoxicosis; Can we do better? *Archives of Disease in Childhood*. 2004;89:745-750.
 6. Griffith E, Nunlist E. Mitral valve prolapse in adolescent female with hyperthyroidism. *Prog Pediatr Cardiol*. 2020; 58:101264.
 7. Lester LA, Sodt PC, Rich BH, Anne W. Lucky, Nancy Hutcheon, René A. Arcilla. Cardiac abnormalities in children with hyperthyroidism. *Pediatr Cardiol*. 1982;2:215–223.
 8. Doucette S, Tierney A, Roggensack A, Yusuf K. Neonatal Thyrotoxicosis with Tricuspid Valve Regurgitation and Hydrops in a Preterm Infant born to a mother with Graves' Disease. *AJP Rep*. 2018;8(2): 85-e88. DOI: 10.1055/s-0038-1645879.
 9. Ertek S, Cicero AF. Hyperthyroidism and cardiovascular complications: A narrative review on the basis of pathophysiology. *Arch Med Sci*. 2013;9(5): 944-52.
 10. Fekri K, Michel CM, Tamilia M. Reversible, severe mitral regurgitation in thyrotoxic Graves' disease. *BMJ Case Rep*. 2021;14(2):e239626. DOI: 10.1136/bcr-2020-239626.
 11. Karashima S, Tsuda T, Kometani M et al. Severe mitral regurgitation as a result of rupture of mitral valve chordae tendineae in a patient with Graves disease. *J Endocr Soc*. 2018;2:1246–50. DOI: 10.1210/js.2018-00173

© 2024 Ujunwa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112389>