



Delineation of Soil Fertility Status and KVK Intervention to Improve the Productivity of Pepper in TSP Villages of Namakkal District under Soil Health Card Mission

S. Sathya ^{a++*}, K.Velmurugan ^{a#}, S. Paulpandi ^{a++}
and M.Daisy ^{at}

^a *Krishi Vigyan Kendra, Veterinary College and Research Institute Campus, TANUVAS, Namakkal, 637002, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i64622

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116291>

Original Research Article

Received: 22/02/2024
Accepted: 24/04/2024
Published: 29/04/2024

ABSTRACT

Soil is the most important resource needed for agriculture production. The unbalanced fertilizer application associated with more production by high yield varieties affects the soil quality vulnerably. Soils of tribal areas, having multi nutrient deficiencies due to intense cropping especially spices Pepper, coffee, cardamom and banana. Balanced application of plant nutrients is thus the only

⁺⁺Assistant Professors;

[#] Professor and Head;

^t Farm Manager;

^{*}Corresponding author: E-mail: sathyasac@gmail.com, sathyassac@gmail.com;

option to increase soil productivity for maintaining/enhancing the overall soil productivity. Delineation of soil nutrient status is an important practice to plan the nutrient scheduling for pepper crop. Hence 100 soil samples were collected from five adopted villages and tested its properties. Based on the analytical results, soil health cards issued along with fertilizer recommendation. Then 50 farmers were selected, trained all scientific package of practices through trainings. The field experiment was conducted in 50 farmer's field in an area of 20 ha. Pepper variety Panniyur 1 was the test crop. Application of soil test based integrated nutrient management practice increased the yield attributes and yield of pepper. An average of Rs. 74,220/ha net profit could be obtained by adopting all scientific package of practices in the farm holdings.

Keywords: Soil nutrients status; delineation; scientific intervention; pepper yield; BC ratio.

1. INTRODUCTION

Kollihills is a small mountain range located at the tail end of Eastern Ghats in the Namakkal of Tamil Nadu situated between 11.10' and 11.23' North Latitude and 78.17' and 78.28' East Longitude, the mountains are at 1000-1300 m height covering approximately 418.5 Km². There are 14 villages (Nadus/revenue villages) and 275 hamlets inhabited predominantly (>95 per cent) by scheduled tribes. The total population of Kollihills is 33888 living in 6480 households. Out of the 38678 people 36080 are Tribals and the rest have been declared as scheduled caste. The Malayali Malai Vazh Makkal is the major scheduled tribe's peoples living in this area.

The main occupation is agriculture and cattle rearing. Black Pepper, Hill Banana, Coffee, Pine apple, Tapioca, Paddy, Minor Millets like Foxtail Millet, Finger Millet and Little Millets, Vegetables are the Major crops cultivated in Kollihills in an area of 8745 ha under both irrigated and rainfed conditions. Recent studies said that soil fertility status getting deteriorated due to poor nutrient management practice adopted by the farmers thus led to stunted growth, flower drops, fruit shedding and susceptible to pest and disease attack which in turn resulted in yield loss.

To assess the soil fertility and suggest suitable management practices delineation of soil nutrient status is important. Farmers being practicing nutrients application either through organic way and unbalanced amount of inorganic fertilizer application might be strongly associated with multinutrient deficiency. Right applications of essential nutrients are critically needed to increase pepper production and quality by the way of increasing growth and quality traits. Pepper yield is decreased tremendously and crops are more susceptible for pest and disease attack. To keeping these points in view, fertility status study was planned and implemented in

adopted village of tribal areas. After testing, soil health card based fertilizer application promoted through integrated nutrient management. INM concept includes organic manures, biofertilizers application and inorganic fertilizers. Recently tribal farmers sensitized about biofertilizers application especially azospirillum, phosphobacteria, vesicular Arbuscular Microrrhiza and Arka Microbial Consortia. Microorganisms play a crucial role in mineralization and mobilization of nutrients in soil and facilitate the uptake by plants. Keeping all these points in view the present study was programmed.

2. MATERIALS AND METHODS

Gundurnadu panchayat was selected in Kollihills and soil samples were collected in Elangiyampatti, Nathukulipatti, A.Thannimathipatti, Semmedu and Thambapadipatti villages. Soil samples were collected, brought to the laboratory air dried, processed and analyzed for physico-chemical, chemical, physical and biological parameters after passing through 2 mm sieve. Soil pH and EC were determined in Soil: Water (1: 2.5ratio) extract by potentiometric and conductometric methods respectively [1]. Organic carbon was estimated by chromic acid wet digestion method [2]. Available N in soil was estimated by alkaline permanganate method [3], available P by Colorimetric method Olsen *et al.*, [4], available K by Neutral Normal Ammonium Acetate method [5], available S by Turbidimetric method [6].

After the soil testing, importance of soil health was sensitized among farmers and farm women through various group meeting, awareness programme, field visits, regular advisories and distribution soil health cards with suitable recommendations. Then 50 interested farmers were selected and FLD programme with INM practice was implemented in 2023-2024.

The treatment details as follows.

Treatments	Details
Check - Farmers Practice	<ul style="list-style-type: none"> • Cattle manure @ 6 kg/ vine applied once in 6 months • Neem cake @ 250 g/ vine and complex fertilizers @ 100 g/ vine applied twice once 6 months interval
Demo – INM concept	<ul style="list-style-type: none"> • FYM/compost @ 10 kg/ vine/year in two splits @ 6 months interval • Azospirillum @ 50 kg, phosphobacteria @ 50 kg, potash solubilizing bacteria @ 50 g, & VAM @ 200 g along with 50 % recommended dose of chemical fertilizers (according to soil health cards) applied in split doses twice in circular basin excavated in 1 feet away from the base of the pepper vine during month of May - June and September - October months. • Lime @ 500g / vine applied once in two years during May-June immediately after the onset of monsoon.

Among the various spices, pepper cultivated in larger scale for income generation and hence according to the soil fertility status nutrient management practices were suggested through integrated nutrient management concept in an area of 20 ha (50 farmers field) in randomized block design. Plot size was 0.5 acre (50 cents each) for check and 0.5 acre for INM demo. The plant population was maintained @ 1110 /ha.

After implementation of treatments, growth and yield attributes like length of spike, no. of spikes/vine, no. of berries/spike, pepper yield/vine (kg), pepper yield /ha (kg) were monitored through field visits and yield data & its attributes was collected and economics were worked out. Panniyur 1 pepper variety was taken as test crop for this experiment.

3. RESULTS AND DISCUSSION


3.1 Soil Reaction and Electrical Conductivity

Predominant soil type was red sandy loam in texture and non calcareous in nature. From this soil analysis, analysis report revealed that value of soil reaction (pH) was ranged from 6.67 to 7.27 (Table 1). Climatic condition and topography played a vital role in decreasing soil pH. None of the soil samples having alkaline pH like plain soil. The decrease in pH might be attributed to the availability of higher amount of organic carbon. Release of organic and inorganic acids during decomposition of organic matter present in the hilly soils also directly contributed to the reduction of soil pH. Kaleem Abbasi & Ghulam Rasool [7] also confirmed the above findings. Farmers in hilly areas doesn't have the tendency to dump more chemical fertilizers even right amount and leaching loss of salts by slopy nature of topography might be the reason behind in

non saline nature. Soil electrical conductivity was found to be normal in all villages. The value recorded was 0.049 - 0.098 dS m⁻¹. The organic carbon content varied from 0.54 to 0.79 %. The highest 0.79 % OC of the hill soil might be due to prevalence of low temperature and accumulation of organic material in the top soil compared to plain soil [8]. Plain soils in Namakkal district always reported OC found to be in low to medium category and farmers who applied right dose of organic manures had medium category. The same was confirmed by repeated analysis in the Soil and Water Testing.

3.2 Soil Available Nutrients

Available nitrogen content ranged from 198 to 302 kg ha⁻¹ in all villages and average N value was found to be low category 245.50 kg ha⁻¹. Available phosphorus content recorded high category in all villages and average P content was 29.91 kg ha⁻¹. Regarding potassium status recorded low to medium category and had a Similarity to N status. Available S content recorded low (6.78 ppm) to medium (14.32 ppm) category and average S content was 9.76 ppm (Table 1 & Graph 1). Depletion and loss of mobile nutrients especially Nitrogen & Potassium along with running water during the heavy pour of rainfall might be associated with relatively less built up of nutrient in hilly soil. Indiscriminate use of chemical fertilizers and non inclusion of biofertilizers might be strongly contributed to the multinutrient deficiency in Hilly soils which in turn reflected in yield loss in farmers' practice in adopted villages. Wang *et al.*, [9] reported that land cover pattern, distance of sampling and topography played a vital role in changing the available nutrient status of soils analyzed under various soil situations.



Graph 1. Soil available nutrients present in different regions


Table 1. Average nutrient status of Kollihills of Namakkal District, TN under Tribal sub plan and soil health card mission

Soil parameters	Average nutrient of Gundurnadu Panchayat of Kollihills block											
	Elangiyampatti N=28		Nathukulipatti n=15		A. Thannimathipatty N=15		Semmedu N=23		Thambapadipatti N=19		Average (100)	
Soil type	Red soil		Red soil		Red soil		Red soil		Red soil			
Soil texture	Sandy Loam		Sandy Loam		Sandy Loam		Sandy Loam		Sandy Loam			
Lime status	Non calcareous		Non calcareous		Non calcareous		Non calcareous		Non calcareous			
pH	6.98	Neutral	6.67	Neutral	7.27	Neutral	6.86	Neutral	7.09	Neutral	6.97	Neutral
EC (dSm ⁻¹)	0.089	Non saline	0.091	Non saline	0.049	Non saline	0.096	Non saline	0.098	Non saline	0.080	Non saline
Organic carbon (%)	0.79	H	0.69	M	0.56	M	0.78	H	0.54	M	0.67	M

Soil parameters	Average nutrient of Gundurnadu Panchayat of Kollihills block											
	Elangiyampatti N=28		Nathukulipatti n=15		A. Thannimathipatty N=15		Semmedu N=23		Thambapadipatti N=19		Average (100)	
Available nitrogen (kg ha ⁻¹)	279	L	245	L	203	L	302	M	198	L	245.40	L
Available Phosphorus (kg ha ⁻¹)	39.21	H	33.2	H	24.1	H	26.70	H	26.32	H	29.91	H
Available Potassium (kg ha ⁻¹)	178	M	154	M	106	L	191	M	203	L	166.40	M
Available Sulphur (mg kg ⁻¹)	14.32	M	10.23	M	9.43	L	6.78	L	8.02	L	9.76	L

Table 2. Yield attributes, yield and economics of black pepper

Treatments	Number of spikes (No.)	Length of spikes(cm)	Number of berries/spike (No.)	Yield (kg/vine)	Yield (kg/ha)	Cost of cultivation (Rs./ha)	Gross income (Rs./ha)	Net income (Rs./ha)	BC ratio
Farmers practice (check)	167	12.4	76	1.19	1321	1,34,560	3,05,010	2,03,000	2.26
KVK intervention	209	14.3	89	1.66	1844	1,46,900	4,24,120	2,77,220	2.89
CD (P=0.05)	13.50	0.84	3.69	0.130	121.81				



Graph 2. Yield attributes and economics of black pepper

3.3 Yield Attributes and Yield of Pepper

Yield and its attributes were found to be higher in KVK intervention than farmers practice. More No. of spikes was 209, length of spikes 14.3 cm, no. of berries/spike 89 and yield of dry berries/ vine was 1.66 kg observed in vines received all scientific packages of practices (Table 2 & Fig.2). Application of organic manures releases organic acids upon decomposition which mineralize the nutrients and make them available to plants. N fixing ability of bacteria, P & K solubilizing and mobilizing capacity bacteria materializes and increases the availability the nutrients from applied as well as native reservoir to the plants during requirement of various growth stage of pepper [10]. Proper nutrient supplementation to all growth stages directly contributed the yield enhancement and hence INM treatment proved its superiority in terms of recording higher yield bearing attributes than previous years and check [11].

3.4 Economics of Experiment

From this study it can be clearly stated that farmers got an additional income of Rs.74,220/ha by recording yield increase of 39.59% over check. Farmers sensitized about composting of locally available biodegradable waste, other organic manure preparation in their field itself. Hence cost involved in OM, bio agents were not included in the cost of production. Biofertilizers cost, agro chemicals and labour cost were included in both practices. Hence scientific package of practices recorded higher BC ratio. Upon field observation and survey carried out among farmers, it can be clearly stated that Quick wilt symptoms, yellowing of pepper vine and berry shedding at pin head stage was reduced upto 16.8% when farmers adopted suitable scientific management practices especially during May- June.

4. CONCLUSION

At this juncture, it can be concluded that soil fertility was comparably have less nutrient status and hence suitable management practices were suggested through KVK FLD programme. Farmers applied all the nutrient management practices and got appreciable yield bearing nature of pepper and recorded higher BC than conventional practice. Through this interventions, nearby farmers and villages were started to adopt complete scientific package of practice for pepper cultivation in hilly areas. Farmers

experienced better through Tribal sub plan and now days significant changes in farming operations had been observed. They voluntarily buy biofertilizers from KVK and inculcated the habit of application of biofertilizers for pepper cultivation largely.

ACKNOWLEDGEMENT

Authors would like to thank TANUVAS, Chennai, TNAU, Coimbatore, for technical guidance and ICAR, ATARI, Hyderabad for funding to carry out the FLD successfully under Tribal Sub plan at Namakkal district, Tamil Nadu.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Jackson ML. Soil Chemical Analysis. Prentice Hall of India Ltd., New Delhi; 1973.
2. Walkley A, Black CA. An examination of Degitijareff method for determining the organic matter and proposed modification of the chromic acid titration method. *Soil Sci.* 1934;37:29-38.
3. Subbiah BV, Asija GL. A rapid procedure for estimation of available nitrogen in soils. *Curr. Sci.* 1956;65:477-480.
4. Olsen SR, Cole CV, Watanabe PS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *U.S.D.A. Circ.* 1954;939.
5. Stanford S, English L. Use of Flame photometer in rapid soil test for K and Ca. *Agron J.* 1949;41:446-447.
6. Williams CH, Steinbergs A. Soil sulphur fractions as chemical indices of available sulphur in some Australian soils. *Aus. J. Agronomical Res.* 1959;10:340-352.
7. Kaleem Abbasi M, Ghulam Rasool. Effects of different land-use types on soil quality in the hilly area of Rawalakot Azad Jammu and Kashmir, *Acta Agriculturae Scandinavica Section B-Soil and Plant.* 2005;55(3):221-228
8. Navneet Pareek KP, Raverkar MK, Bhatt Shikhar Kaushik, Subhash Chandra, Gurvinder Singh, Hem C. Joshi. *ENVIS Bulletin Himalayan Ecology.* 2019;27:89-92.

9. Wang R, Zou R, Liu J, Liu L, Hu Y. Spatial distribution of soil nutrients in farmland in a Hilly Region of the Pearl River Delta in China Based on Geostatistics and the Inverse Distance Weighting Method. Agriculture. 2021;11:50.
10. Manju Jincy Varghese, Preethu K Paul and R. Marimuthu. Effect of integrated nutrient management on yield of black pepper. J.Krishi Vigyan. 2021;10(1):73-76.
11. Shilpa AK, Sharma Shivender Thakur, Kajal Sharma. Integrated nutrient management of peppers (*Capsicum annum* L.) in 21st Century - A Review. Int. J. Curr. Microbiol. App. Sci. 2017; 6(12):227-237.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116291>