



Associated Factors of Low Birth Weight among Newborns at Hospitals in Kisangani, Democratic Republic of Congo

Anzani Malongo ^a, Komanda Likwekwe Isaac ^b,
Gelengi Mbongo ^c, Balimo Limbele ^c, Ekili Molisho ^c,
Melangana Menama ^c, Botela Ponea ^c, Ramazani Tabora ^c,
Kalafulu Imurani ^c, Boseleka Bokumbuangole ^c,
Bokanga Ekelenge ^d, Issa Yakusu ^e,
Tshingombo Muyaya Trydon ^{f++} and Ossinga Bassandja ^{g*}

^a Higher Institute of Rural Development of Djolu, DR Congo.

^b University Clinics of Kisangani, DR Congo.

^c Higher Institute of Medical Techniques in Kisangani, DR Congo.

^d Department of Gynecology-Obstetrics, Faculty of Medicine and Pharmacy, University of Kisangani, DR Congo.

^e Cardiology Department, Cheik Anta Diop University, Senegal.

^f Faculty of Medicine and Pharmacy, University of Kisangani, DR Congo.

^g Department of Internal Medicine, Faculty of Medicine and Pharmacy, Kisangani University, DR Congo.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajpr/2024/v14i11397>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124779>

⁺⁺ Doctoral Student;

^{*}Corresponding author: Email: jacquesossinga@gmail.com;

Cite as: Malongo, Anzani, Komanda Likwekwe Isaac, Gelengi Mbongo, Balimo Limbele, Ekili Molisho, Melangana Menama, Botela Ponea, Ramazani Tabora, Kalafulu Imurani, Boseleka Bokumbuangole, Bokanga Ekelenge, Issa Yakusu, Tshingombo Muyaya Trydon, and Ossinga Bassandja. 2024. "Associated Factors of Low Birth Weight Among Newborns at Hospitals in Kisangani, Democratic Republic of Congo". *Asian Journal of Pediatric Research* 14 (11):8-16. <https://doi.org/10.9734/ajpr/2024/v14i11397>.

ABSTRACT

Background: Low birth weight remains a major public health problem in every country in the world. It is a predictor of early neonatal mortality, morbidity and long-term health. The aim of this study was to determine the prevalence of low birth weight and to identify the associated factors in a hospital setting in Kisangani.

Material and Methods: An unmatched case-control study was used including all live and full-term births was conducted from 1 December 2023 to 31 May 2024 in the maternity wards of Makiso-Kisangani General Referral Hospital, Mangobo General Referral Hospital, Tshopo General Referral Hospital and Alwaleed Health Centre. A sample of 646 newborns was selected: 86 cases (newborns weighing less than 2500 grams) and 560 controls (newborns weighing 2500 grams or more). Factors associated with prehypertension were identified using logistic regression.

Results: The prevalence of low birth weight was 13.4%. Factors associated with low birth weight were maternal age less than 18 years, intergenital interval less than 2 years, maternal infection during pregnancy, high blood pressure, level of primary education and urban residential environment.

Conclusion: This study shows that low birth weight remains a public health problem in Kisangani. In addition to shedding light on the other contributing variables and the connections between them, prospective national research would enable the development of preventative measures.

Keywords: Prevalence; associated factors; low birth weight; Kisangani; Democratic Republic of Congo.

1. INTRODUCTION

Low birth weight (LBW) is defined as a weight of newborns at birth less than 2500 gram by World Health Organization. More than 20 million infants are born with low birth weight in worldwide (Weise, 2012; United Nations, 2015).

LBW continues to remain a major public health problem worldwide, especially in the developing countries. The birth weight of an infant is the single most important determinant of its chances of survival, healthy growth, and development (Bendhari & Haralkar, 2015). There is widespread agreement that having LBW at birth disadvantages the child. LBW accounts for 60 to 80% of all newborn deaths. Along with short- and long-term infant and childhood morbidity, it is a significant cause of perinatal mortality. Infants born with birthweights of less than 2500 g had a mortality rate that was up to 40 times greater, and they had a significantly increased chance of developing long-term disabilities (Ghimire et al., 2014; Khan et al., 2016; Ramakrishnan, 2004; Barker & Godfrey, 2004; World Health Organization, 2011; Goldenberg & Culhane, 2007; Torres-Arreola et al., 2005).

In 2020, the prevalence of LBW was greater than 10 % in most African countries, and more

specifically greater than 15 % in 16 countries, with the highest rate (23 %) recorded in the Comoros. The prevalence of LBW was around 10 % or less in eight countries (Algeria, Eswatini, Kenya, Democratic Republic of Congo, Rwanda, Sierra Leone, Tunisia and United Republic of Tanzania) (UNICEF & World Health Organization, 2019).

National measures to enhance mother and child health have been implemented, but the Democratic Republic of the Congo (DRC) still faces a public health crisis related to LBW. For a number of reasons, including the fact that the underlying mechanisms causing LBW remain poorly understood, it is challenging to prevent risk factors. Very little is known about LBW and its contributing causes in Kisangani. This is the context for our study, the first of its kind in our community, which aims to better target and combine interventions to prevent LBW.

2. MATERIALS AND METHODS

2.1 Study Design

An unmatched case-control study was used.

2.2 Setting and Study Period

This study was conducted in the maternity wards of Makiso-Kisangani General Hospital, Mangobo General Hospital, Tshopo General Hospital and Alwaleed Health Centre. These maternity units were chosen because of the high number of births recorded and their easy geographical accessibility. This work was carried out over a period running from 1 December 2023 to 30 May 2024.

2.3 Study Target Population

The target population for this study was newborns born live at term in the above-mentioned health facilities during the study period.

2.4 Sampling and Sample Size

Non-probability convenience sampling was used. A total of 646 respondents were enumerated for this study.

2.5 Selection of Respondents (Inclusion and Exclusion Criteria)

Live newborns born at term with no known risk factors (i.e. intrauterine growth retardation) for low birth weight were included in the study.

Mothers who gave birth prematurely (before 37 completed weeks of gestation) and newborns from twin pregnancies were excluded from the study.

Mothers who gave birth to newborns weighing less than 2500 grams were cases and newborns ≥ 2500 grams were controls.

2.6 Study Variables

For the variables studied, cases comprising newborns with LBW were compared with controls, comprising all full-term newborns with a birth weight of between 2500 and 4000g. The dependent variable was low birth weight. The independent variables were the mother's socio-demographic data (age, profession and socio-economic level), parity and medical history, as well as the course of the last pregnancy (pathologies during pregnancy, follow-up, term, gestational age), the characteristics of the delivery (mode) and the characteristics of the newborn (term, weight, size, sex, malformations

and transfer to the neonatology department). Anaemia in pregnant women was defined as an Hb level of less than 10g/dl. Arterial hypertension was defined as a blood pressure greater than 135/85mmHg. Maternal infection was defined as any bacterial, viral, parasitic or mycotic infection occurring during gestation. The nutritional status of the mothers was assessed by calculating the body mass index (BMI) determined by the ratio of body weight (in kilograms)/square of height (in metres). A BMI of between 18.5 and 24.9 kg/m² corresponded to good nutritional status. Malnutrition was defined as a BMI of less than 18.5 kg/m².

2.7 Data Collection

Data were collected retrospectively, using a documentary analysis technique. We used women's medical delivery records and registers using a pre-established data collection form containing the study variables.

2.8 Data Processing and Analysis

The data were entered and processed using SPSS software (IBM SPSS Statistics 22).

Frequencies, percentages, means and standard deviations were used to describe maternal sociodemographic and obstetric characteristics. Factors associated with prehypertension were identified using logistic regression.

3. RESULTS

3.1 General Characteristics of Term Newborns in the Study Sites

The data from this study indicate that the majority of newborns were male, both among those with LBW and those of normal weight. The mean weight of term newborns ranged from 1958 \pm 339 g among LBW births to 3329 \pm 480 g among normal weight births. The above data are presented in Table 1.

3.2 Socio-demographic Data of the Study Participants

These data show that most of the study participants were aged between 18 and 35, were Christian, had no occupation, were illiterate and lived in urban areas. These data are presented in Table 2.

Table 1. General characteristics of term newborns in the study sites

General characteristics	Weight	
	< 2500 grams n (86)	≥ 2500 grams n (560)
Sex of newborn		
Male[n (%)]	45 (52,3)	287 (51, 2)
Female [n (%)]	41 (47,7)	273 (48,8)
Newborn weight [mean ± SD]	1958±329 grams	3329±480 grams

Table 2. Socio-demographic data of mothers taking part in the study

Socio-demographic data	N	%
Age (years)		
< 18	29	4,5
18-35	517	80
>35	100	15,5
Residence		
Urban environment	581	90
Rural	65	10
Religion		
Christian	587	90,8
Muslim	32	4,9
Kimbanguist	19	3
Other	8	1,5
Profession		
With profession	529	81,8
No profession	117	18,2
Level of education		
Illiterate	387	58,5
Primary	129	20
Secondary	77	12
University	53	9,5

3.3 Obstetrical Characteristics of Study Participants

The obstetric data provided the following information:

- the majority of the women surveyed had an intergenital interval of less than 24 months and a normal BMI;
- Most of them had no history of abortion or stillbirth and had given birth vaginally;
- The majority were multiparous, normotensive and had not contracted any infection during pregnancy;
- Anaemia was found in 53.5% of mothers who had given birth to LBW babies, compared with 25% of those who had given birth to babies of normal weight. These data are given in Table 3.

3.4 Prevalence of Low Birth Weight in the Study Sites

Of the 646 full-term newborns in the study sites, 86 were of low birthweight, representing a prevalence of 13.4%.

3.5 Association between Maternal, Obstetric and Neonatal Characteristics and Newborn Birth Weight

Multivariate analysis indicated that maternal age less than 18 years, interpregnancy interval less than 2 years, maternal infection during pregnancy, arterial hypertension, primary education level, urban residential environment and male sex of the newborn were significant predictors of LBW. These data are presented in Table 4.

4. DISCUSSION

4.1 General Characteristics of the Study Population

The aim of this unmatched case-control study was to determine the prevalence of LBW in hospitals in Kisangani and to identify associated factors. Previous studies have matched one case to one control (Camara et al., 1996; Mabilia-Babela et al., 2007), while other authors have matched more than one control to a case. In the present study, we matched 6 controls to one case (Bwana et al., 2014; Demelash et al., 2015; Hassoune et al., 2015).

4.2 Prevalence of Low Birth Weight

Our study revealed a prevalence of low birth weight of 13.4%. This prevalence is close to that found in studies conducted in other regions of DR Congo, i.e. 11.6% and 11.8% respectively (Mamba et al., 2021; Saasita et al., 2022). However, it remains high compared with the national rate according to the demographic and health survey, which reported a rate of 7% and 6.3% for Katanga province (MPSMRM, MSP & ICF International, 2013-2014).

High prevalence rates have also been reported in studies carried out in Zimbabwe and Morocco, where they were 19.9% and 22% respectively (Elaabsi et al., 2022; Feresu et al., 2004). The differences in prevalence observed in these

studies could be explained by the multiplicity of risk factors, in particular malnutrition, inadequate monitoring of pregnancy, malaria and repeated urogenital infections during pregnancy (Kakudji et al., 2015).

4.3 Factors Associated with LBW

4.3.1 Maternal age under 18 years

According to the results of studies carried out in Tunisia and DR Congo, maternal age under 18

years was a factor in the occurrence of LBW (Bwana et al., 2014; Kakudji et al., 2015; Letaief et al., 2001). Our results concur with these studies, with a statistically significant difference (p=0.003). Several factors could explain this finding, including competition for nutrients between the growing adolescent and the foetus, which appears to be at the root of the problem, and competition between pregnancy and growth, which has a particularly unfavourable effect on the micronutrient status of adolescent girls (Bwana et al., 2014; Kakudji et al., 2015).

Table 3. Obstetric characteristics of study participants (N= 646). Bivariate analysis.

Obstetrical characteristics	Case (n= 86)	Control (n= 560)	Pearson Chi-square
Intergenic interval			0.000
< 24 months [n (%)]	69 (80,2)	406 (72,5)	
≥ 24 months [n (%)]	17 (19,8)	154 (27,5)	
Hypertension			0.007
Yes [n (%)]	4 (4,6)	14 (2,5)	
No [n (%)]	82 (95,4)	546 (97,5)	
Anemia			0.002
Yes [n (%)]	46 (53,5)	140 (25)	
No [n (%)]	40 (46,5)	420 (75)	
Maternal infection during pregnancy			0.000
Yes [n (%)]	27 (31,3)	168 (30)	
No [n (%)]	59 (68,7)	392 (70)	
BMI			0.453
< 18,5 [n (%)]	9 (10,4)	39 (6,9)	
18,5 - 24,9 [n (%)]	58 (67,4)	392 (70)	
25 - 29,9 [n (%)]	13 (15,1)	95 (17)	
≥ 30 [n (%)]	6 (7,1)	34 (6,1)	
Number of pregnancies			0.089
1 [n (%)]	29 (33,7)	201 (35,8)	
2 [n (%)]	18 (21)	128 (22,8)	
3 [n (%)]	39 (45,3)	231 (41,4)	
History of stillbirth			0.167
Yes [n (%)]	3 (3,4)	9 (1,6)	
No [n (%)]	83 (96,6)	551 (98,4)	
Route of delivery			0.345
Vaginal delivery [n (%)]	77 (89,5)	498 (88,9)	
Caesarean section [n (%)]	9 (10,5)	62 (11,1)	
History of abortion			0.875
Yes	13 (15)	67 (12)	
No	73 (85)	493 (88)	

Table 4. Association between maternal, obstetric and neonatal characteristics and birth weight of newborns

Characteristics	Adjusted OR (95% CI)	p
Maternal age under 18	1.045 (1.013-1.070)	0.003
Intergenic interval less than 2 years	1.810 (1.264-2.602)	0.002
Maternal infection during pregnancy	3.960 (2.718-5.565)	0.000
Hypertension	3.324 (1.697-6.468)	0.000
Primary education level	0.453 (0.279- 0.747)	0.002
Urban residential environment	0.523 (0.334- 0.814)	0.004
Anemia	1,22 (1,07-1,47)	0,002

4.3.2 Intergenerational interval of less than 2 years

It appears that an intergenerational interval of less than 2 years was associated with low birth weight in the maternity units visited in Kisangani during the present study ($p= 0.002$). This result is similar to that of Letaief et al., 2001 and Mamba et al., 2021. Maternal exhaustion syndrome describes what happens when a woman's body is used without rest and without time to recover. Other factors, such as the energy costs of pregnancy and lactation, particularly in the context of close reproductive cycles with no recovery time for the woman, lead to a cumulative deterioration in maternal nutritional status, which can lead to LBW (Bwana et al., 2014).

4.3.3 Maternal infection during pregnancy

Our study showed that maternal infection during pregnancy was identified as a predictor of LBW. This result corroborates the observations made by other authors on this subject (Hassoune et al., 2015; Mamba et al., 2021; Siza, 2008; Zheng et al., 2015). However, a study from Benin found no correlation between infection during pregnancy and the birth of a newborn with LBW (Zheng et al., 2015). The normal course of pregnancy can be disrupted by infections, firstly because of the inflammatory reaction that accompanies any infection and secondly because, during an infection, the woman is weakened and obliged to draw on her own nutritional and energy reserves (Setondji, 2014).

4.3.4 Hypertension

The bivariate and multivariate analyses (OR =0.453 ; 0.277-0.747 ; $p= 0.002$) performed with our data indicate that there is a correlation between hypertension and LBW. This result is similar to that of Mabiála-Babela et al., 2007 and Nkwabong et al., 2015. High blood pressure is thought to disrupt blood circulation and consequently placental exchanges, leading to a reduction in both oxygen and micronutrient supply from the mother to the foetus during pregnancy (Mamba et al., 2021).

4.3.5 Level of primary education

According to the results of this study, mothers with a high level of education were less likely to give birth to a low birth weight baby than those

with a primary education. This observation is consistent with several studies associating LBW with maternal education (Tampah-Naah et al., 2016; Amosu et al., 2014).

4.3.6 The urban residential environment

In this study, the probability of giving birth to a low birth weight baby was significantly higher among urban women, which is consistent with studies conducted in Ethiopia and Bangladesh (Tema, 2006; Azimul et al., 2009).

According to another study conducted in Zambia, women living in urban areas were unaware of the benefits of starting antenatal care early, and lacked sufficient knowledge about such care and early antenatal attendance (Banda et al., 2012). This constatation confirms the observation made in this study (OR 0.523 (0.335- 0.814; $p=0.004$).

4.3.7 Anemia

According to this research, it has been shown that there is a relationship between anaemia and LBW.

Similar results have been obtained by other researchers (Figueiredo et al., 2018; Rahmati et al., 2017). Gestational anaemia can lead to a number of complications, one of the most significant of which could be the development of LBW due to poor intrauterine growth associated with disruption of foeto-placental blood flow, resulting in reduced micronutrient and oxygen intake (Setondji, 2014).

5. LIMITATIONS

In this study, the main limitation lies in the data used to carry out the analyses. It was collected primarily for routine healthcare rather than for a specific intervention. It is therefore possible that errors were made when documenting the records. Lastly, this study's analysis of the data only captures the reality of a small number of public health facilities. Consequently, the findings cannot be extrapolated to moms who gave delivery at home or in private healthcare facilities.

6. CONCLUSION

This study shows that low birth weight remains a public health problem in Kisangani. The research was carried out in four maternity units in the city

and revealed certain factors predictive of LBW, such as maternal age under 18, birth interval under 2 years, maternal infection during pregnancy, hypertension, primary education level, urban residential environment and anemia. In addition to shedding light on the other contributing variables and the connections between them, prospective national research would enable the development of preventative measures.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ETHICAL APPROVAL

Before starting the survey, a request for authorisation was sent to the medical-health and administrative authorities of the health facilities concerned. The anonymity of the questionnaires ensured the confidentiality of the data.

CONSENT

As per international standards, parental written consent has been collected and preserved by the author(s).

ACKNOWLEDGEMENTS

The authors wish to thank student Kengakoy Bukuka for actively participating in the data collection and the first frame of this work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

Amosu, A. M., Degun, A. M., & Ter, G. D. (2014). Maternal socio-demographic characteristics as correlates of newborn birth weight in Abeokuta city, Nigeria. *Biomedical Research*, 25(4), 612-616.

Azimul, S. K., Matin, A., Shabnam, S., Shamianaz, S., & Bannerje. (2009). Maternal factors affecting low birth weight in urban area of Bangladesh. *Journal of Dhaka Medical College*, 18(1), 64-69.

Banda, I., Michelo, C., & Hazemba, A. (2012). Factors associated with late antenatal care attendance in selected rural and urban communities in the Copperbelt Province of Zambia. *Medical Journal of Zambia*, 39(3), 29-36.

Barker, D. J. P., & Godfrey, K. M. (2004). Maternal nutrition, fetal programming and adult chronic disease. In M. J. Gibney, B. M. Margetts, J. M. Kearney, & L. Arab (Eds.), *Public health nutrition*. Oxford: Blackwell Science.

Bendhari, M. L., & Haralkar, S. J. (2015). Study of maternal risk factors for low birth weight neonates: A case-control study. *International Journal of Medical Science and Public Health*, 4, 987-990.

Bwana, K. I., Kilolo, N. U. E., Kabamba, N. M., & Kalenga, M. K. P. (2014). Risk factors for low birth weight in semi-rural Kamina, Democratic Republic of Congo. *The Pan African Medical Journal*, 23(17), 220.

Camara, B., Diack, B., Diouf, S., Signate Sy, H., Sall, M. G., & Ba, M. (1996). Low birth weight: Frequency and risk factors in the Guediawaye district (suburb of Dakar – Senegal). *Médecine d'Afrique Noire*, 43(5), 260-265. PubMed | Google Scholar

Demelash, H., Motbainor, A., Nigatu, D., Gashaw, K., & Melese, A. (2015). Risk factors for low birth weight in Bale zone hospitals, South-East Ethiopia: A case-control study. *BMC Pregnancy and Childbirth*, 15, 264. PubMed | Google Scholar

Elaabsi, M., Loukid, M., & Lamtali, S. (2022). Cross-sectional study of factors associated with low birth weight in the Marrakech region (Morocco). *PAMJ - One Health*, 9(7). <https://doi.org/10.11604/pamjoh.2022.9.7.33034>

Feresu, S. A., Harlow, S. D., Welch, K., & Gillespie, B. W. (2004). Incidence of and socio-demographic risk factors for stillbirth, preterm birth and low birth weight among Zimbabwean women. *Paediatric and Perinatal Epidemiology*, 18(2), 154-163. PubMed | Google Scholar

Figueiredo, A. C. M. G., Gomes-Filho, I. S., Silva, R. B., Pereira, P. P. S., Mata, F. A. F., Lyrio, A. O., et al. (2018). Maternal anemia and low birth weight: A systematic review and meta-analysis. *Nutrients*, 10, 601.

Ghimire, R., Phalke, D. B., Phalke, V. D., Banjade, B., & Singh, A. K. (2014).

- Determinants of low birth-weight: A case control study in Pravara rural hospital in western Maharashtra, India. *International Journal of Scientific Research*, 3(7), 2277-8179.
- Goldenberg, R. L., & Culhane, J. F. (2007). Low birth-weight in the United States. *American Journal of Clinical Nutrition*, 85(2), 5845-905.
- Hassoune, S., Bassel, S., Nani, S., Elbouri, H., Zine, K., & Maaroufi, A. (2015). Maternal factors associated with low birth weight: A case-control study in a Moroccan public hospital. *Pan African Medical Journal*, 20, 303. PubMed | Google Scholar
- Kakudji, L. P., Mukuku, O., & Kalenga, M. K. P. (2015). Study of low birth weight associated with maternal age and parity in a mother-child couple population followed in Lubumbashi. *Pan African Medical Journal*, 20, 246.
- Khan, A., Deeba, F., & Jaleel, R. (2016). Frequency and risk factors of low birth-weight in term pregnancy. *Pakistan Journal of Medical Sciences*, 32(1), 138-142.
- Letaief, M., Soltani, M. S., Ben, S. K., & Bchir, A. (2001). Epidemiology of low birth weight in the Tunisian Sahel. *Public Health*, 13(4), 359-366.
- Mabiala-Babela, J. R., Matingou, V. C., & Senga, P. (2007). Risk factors for low birth weight in Brazzaville, Congo. *Journal de Gynécologie Obstétrique et Biologie de la Reproduction*, 36(8), 795-798. PubMed | Google Scholar
- Mamba, T. C., Mbusa, K. R., Kapiteni, M. W., & Mungo, M. B. (2021). Factors associated with low birth weight of term newborns in Bukavu, Democratic Republic of Congo. *International Journal of Innovation and Scientific Research*, 53(2), 123-131.
- Ministry of Planning and Monitoring of the Implementation of the Revolution of Modernity (MPSMRM), Ministry of Public Health (MSP), & ICF International. (2013-2014). Demographic and health survey in the Democratic Republic of Congo. Google Scholar.
- Nkwabong, E., Nounemi, N. K., Sando, Z., Mbu, R. E., & Mbede, J. (2015). Risk factors and placental histopathological findings of term born low birth weight neonates. *Placenta*, 36, 138-141.
- Rahmati, S., Delpishe, A., Azami, M., Ahmadi, M., & Sayehmiri, K. (2017). Maternal anemia during pregnancy and infant low birth weight: A systematic review and meta-analysis. *International Journal of Reproductive BioMedicine*, 15(3), 125-134.
- Ramakrishnan, U. (2004). Nutrition and low birth-weight: From research to practice. *American Journal of Clinical Nutrition*, 79(1), 17-21.
- Saasita, K. A., Mbahweka, K. F., Wahangire, K. J., Katsonger, K. A., Kyoghero, K. J., Muyisa, M. R., & Kasomo, P. J. (2022). Epidemiological profile and factors of low birth weight in Katwa General Referral Hospital in the North-East of the DR. *Kisangani Medical Journal*, 12(2), 578-583.
- Setondji, G. R. P. (2014). Low birth weight, prematurity and intrauterine growth retardation: Risk factors and consequences on growth from birth to 18 months of life in Beninese newborns. *Public Health and Epidemiology, Pierre and Marie Curie University-Paris VI, Paris*.
- Siza, J. E. (2008). Risk factors associated with low birth weight of neonates among pregnant women attending a referral hospital in northern Tanzania. *Tanzania Journal of Health Research*, 10(1).
- Tampah-Naah, A. M., Yendaw, E., & Tampahnaah, A. M., Anzagra, L., & Yendaw, E. (2016). Factors correlate with low birth weight in Ghana. *British Journal of Medicine and Medical Research*, 4(16), 1-8.
- Tema, T. (2006). Prevalence and determinants of low birthweight in the Jimma Zone, southwest Ethiopia. *East African Medical Journal*, 83(7), 366-371. PubMed.
- Torres-Arreola, L. P., Constantino-Casas, P., Flores-Hernandez, J. P. V.-B., & Rendon, M. E. (2005). Socioeconomic factors and low birth-weight in Mexico. *BMC Public Health*, 5(20), 1-7.
- UNICEF & World Health Organization. (2019). Joint estimates of low birthweight. 2023 edition [Accessed 06 October 2024]. www.who.int/teams/nutrition-and-food-safety/monitoring-nutritional-status-and-food-safety-and-events/joint-low-birthweight-estimates
<https://doi.org/10.4060/CC8743EN-fig23>
- United Nations. (2015). Levels and trends in child mortality: Report 2015. January 2019.
- Weise, A. (2012). WHA global nutrition targets 2025: Low birth weight policy brief. WHO Publications, 1-7. January 2019.

- World Health Organization. (2011). WHO guideline on optimal feeding of low birth-weight in low and middle income countries. Geneva: World Health Organization.
- Zheng, J., Xiao, X., Zhang, Q., Mao, L., Yu, M., & Xu, J. (2015). The placental microbiome varies in association with low birth weight in full-term neonates. *Nutrients*, 7, 6924-6937.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/124779>