



The Effect of Black Soldier Fly (*Hermetia illucens*) Larvae Feeding on the Growth of Mutiara Catfish Seeds (*Clarias gariepinus*)

Muhammad Mahfud Ashqolani ^{a*}, Kiki Haetami ^a,
Ine Maulina ^a and Yuniar Mulyani ^a

^a Fisheries Study Program, Faculty of Fisheries and Marine Sciences,
Padjadjaran University, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v21i4548

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95808>

Original Research Article

Received: 12/11/2022
Accepted: 18/01/2023
Published: 25/02/2023

ABSTRACT

Aims: The purpose of this study was to determine the level of feeding in the form of black soldier fly larvae (*Hermetia illucens*) to increase the growth of the (*Clarias gariepinus*) mutiara catfish seeds.

Study Design: This study used an experimental method with a completely randomized design (CRD) consisting of four treatments and four replications. The treatments are a combination of commercial feed and maggots consisting of A (control, 100% commercial feed), B (75% commercial feed + 25% dry maggot), C (50% commercial feed + 50% dry maggot) and D (100% dry maggot).

Place and Duration of Study: This research was conducted from August 2022 to September 2022 at the Cipancuh Fish Seed Center, Indramayu Regency. Mutiara catfish seeds were obtained from the Cipancuh Fish Seed Center, commercial feed and dried maggot were obtained from online stores whose quality was guaranteed.

*Corresponding author: Email: muhammad18255@mail.unpad.ac.id;

Methodology: The parameters observed in this study were survival, specific growth rate and water quality. The water quality observed included temperature, pH, DO and ammonia levels. Mutiara catfish fry were maintained for 40 days by feeding them three times a day. Length data collection and survival value are carried out every ten days as well as water quality testing using certain tools.

Results: The results showed that the combination of commercial feed and dry maggot had an effect on the specific length growth rate and survival of mutiara catfish seeds. the combination of 50% commercial feed and 50% dry maggot produced the highest specific length growth rate of 16.19%, 75% commercial feed and 25% dry maggot produced a specific length growth rate of 13.36%, with 100% commercial feed % resulted in a specific growth rate of 11.83% and by feeding 100% dry maggot produced the lowest growth rate of 8.03%. The highest survival rate was in treatment D by feeding 100% dry maggot with a value of 91%, the survival rate in the treatment given a combination of 50% commercial feed and 50% dry maggot resulted in a value of 84%, with 75% commercial feeding and maggot dry 25% resulted in a survival rate of 79.50% and the lowest survival rate at 100% commercial feeding was 78.50%. the water quality of the maintenance media is in accordance with the effective requirements for the maintenance of SNI mutiara catfish seeds.

Conclusion: Giving a combination of 50% commercial feed and 50% dry maggot resulted in the highest specific length growth rate of 16.19%. The highest survival rate of mutiara catfish seeds was obtained by feeding 100% dry maggot, which was 91%.

Keywords: Growth; mutiara catfish seeds; *Clarias gariepinus*; black soldier fly maggot; dry maggot; *Hemeticia illucens*.

1. INTRODUCTION

Mutiara catfish (*Clarias gariepinus*) is a type of catfish selected by the Sukamandi Fish Breeding Research Institute (BRPI) which has been set for release based on the Decree of the Minister of Maritime Affairs and Fisheries Number 77 KEPMEN-KP/2015. Mutiara catfish is a product superior results of selection from African catfish crossed with four strains of African catfish in Indonesia, namely catfish Egypt, paiton, sangkuriang and dumbo as their parent populations [1]. The advantages of mutiara catfish are higher growth, high feed efficiency, uniform size, resistance to disease, resistance to environmental stress conditions, high productivity, and higher cultivation benefits because the proportion of meat is relatively higher [2].

Mutiara catfish included in the class of all-eating or omnivorous but tend to eat meat. Catfish is a type of fish that has a habit of eating at the bottom of waters or ponds (*bottom feeder*) [3]. Mutiara catfish are *nocturnal*, that is, they have a tendency to be active and look for food at night, but in aquaculture they will adapt as a diurnal. During the day catfish prefer to stay or take shelter in the dark parts of the waters. In maintenance ponds, especially in intensive cultivation, catfish can be used to being fed pellets in the morning or during the day. That catfish is a fish that eats all food ingredients (omnivores), both animal and vegetable

ingredients. Mutiara catfish's natural food is small animals, such as water fleas from the *daphnia*, *cladocera* or *copepod* groups [4].

Cannibalism is the act of killing and consuming all or part of the body of individuals of the same species. The nature of cannibalism in catfish is one of the factors causing death, this is due to the non-uniform size of the fish along with the ongoing rearing process [5].

The problem that often arises in the cultivation of sangkuriang catfish (*Clarias* sp) is the high mortality rate of sangkuriang catfish (*Clarias* sp) seeds due to cannibalism in hatchery activities. The mortality rate of sangkuriang catfish (*Clarias* sp) fry due to cannibalism under cultivation conditions can range from 15% -90% mortality [6]. To overcome the problem of cannibalism of mutiara catfish seeds, there are several efforts, one of which is by providing highly nutritious feed.

In fish farming activities, feed is very important in increasing production. In intensive aquaculture, almost 70% of production costs are spent only on procuring feed. To increase profits, fish cultivators must make production costs more efficient, one of which is by reducing feed costs by utilizing natural feeds available in the environment [7].

Natural feed is a kind of fish feed in the form of aquatic organisms which are primary producers

in the food chain. Natural food is able to breed in waters in accordance with its life tolerance. Several types of natural fish feed that can be mass-produced include *infusoria*, *daphnia*, *chlorella*, *tetraselmis*, *rotifers*, *artemia*, to *tubifex* worms or silk worms. In the cultivation of natural feed is needed because it has a high nutritional content and is complete and easy to digest. Natural food has high protein to support the growth of fish such as silk worms which have high protein. The use of natural feed also aims to cut capital for relatively expensive pellet buyers.

Black soldier fly (BSF) larvae or commonly called *maggot* as animal feed has direct and indirect benefits. That prepupa is ideal for use as a mixture of feed or pellet raw materials because it is able to meet the quantity of production [8]. Prepupae are more suitable for direct fish feed, because their small shape matches the size of the fish's mouth. Maggot has *potential* as feed because of its high protein content of 44.26% and fat content of 29.65%. The value of amino acids, fatty acids and minerals contained in the larvae is also not inferior to other protein sources, so that BSF larvae are ideal raw materials that can be used as feed. *Black soldier fly* or *maggot* larvae can also be used as alternative feed or additional feed for cultivated fish, especially mutiara catfish [9].

The best treatment uses a combination of commercial feed and *maggot* (BSF) at a ratio of 50% + 50% by giving the highest daily growth rate value of 1.86%, a length ratio value of 1.40%, a feed efficiency value of 65.09% and the lowest feed conversion ratio value is 1.54 [10].

The growth in length and weight of sangkuriang catfish (*Clarias sp*) in the *maggot* feeding treatment gave better results than the other treatments (treatment with chicken intestine, golden snail and trash fish) with an average average length reaches 7.87 cm, and weight reaches 78.23 grams [11].

The provision of alternative feed in the form of BSF *maggot* had a good effect on the growth of tilapia and the treatment that had the best effect on the growth of tilapia seeds was at the dose of giving a combination of 50% pellet + 50% *maggot feed*, with an absolute growth yield of 5, 5 gr, daily growth of 3.7%, relative growth of 139.4% [12].

The purpose of this study was to determine the level of feeding in the form of *black soldier fly*

(*Hermetia illucens*) larvae to increase the growth of mutiara catfish (*Clarias gariepinus*). It is hoped that by providing feed in the form of black soldier fly larvae (*Hermetia illucens*) it can increase the growth of mutiara catfish (*Clarias gariepinus*).

2. MATERIALS AND METHODS

This research is explorative in nature by taking data from the field. The test fish used in this study were mutiara catfish seeds obtained from the Cipancuh Fish Seed Center. The number of catfish used during this study was 800 fish, measuring 3-5 cm/head with a stocking density of 50 fish/pond. Pool size 100 x 60 x 40 cm³ totaling 16 pools. The research method used was an experimental method with an experimental design using a completely randomized design (CRD). This study used 4 treatments with 4 repetitions which were observed for 40 days. The treatment carried out in this study was Treatment A 100% Commercial Feed (Control), Treatment B Combination of 75% commercial feed + 25% dry black soldier fly larvae, Treatment C 50% commercial feed combination + 50 dried black soldier fly larvae % and Treatment D. Feed in the form of larvae was added with dry black soldier fly larvae as much as 100%. Feeding is done twice a day.

Observations were carried out for 40 days with observation parameters including the degree of survival and the rate of increase in specific length and water quality which were observed every ten days. The survival rate is calculated using the formula [13].

$$SR = (Nt/No) \times 100 \%$$

Information:

SR : Degree of survival (%)
 Nt : Number of live fish at the end of maintenance (tails)
 No : Number of fish at the beginning of rearing (tails)

Calculation of specific length growth as follows:

$$LPPS = (LnLt - LnLo) / t \times 100 \%$$

Information:

LPPS : Specific Length Growth Rate (%)
 Lt : Average length of fish at the end of treatment (cm)
 Lo : Average length of fish at the beginning of treatment (cm)
 t : Maintenance period (days)

Water quality measurement parameters observed included temperature, pH, DO and ammonia levels, for measuring temperature using a thermometer dipped directly into the treatment water, measuring pH using litmus paper, measuring DO using a DO meter and measuring ammonia levels using the a spectrophotometer and additional materials.

The data obtained was tested with statistical and descriptive analysis. Statistical analysis used Analysis of Variance (ANOVA) at a 95% confidence interval. To see the difference in treatment, a further test was carried out with the Duncan test [14].

3. RESULTS AND DISCUSSION

3.1 Survival Rate

Based on the *Analysis of Variance* (ANOVA) test with a 95% confidence level, it showed that the combination of commercial feed and dry *maggot* had a significant effect ($P < 0.05$) on the survival of mutiara catfish seeds with a rearing period of 40 days. For 40 days the maintenance of mutiara catfish seeds has a high survival value. The survival value of catfish seeds was above 80% using a combination of artificial feed and *maggot* [10]. The survival value of catfish seeds at a size of 3-5 cm is said to be good if the value is greater than 80%. Data on the survival of catfish seeds given a combination of artificial feed and *maggot* can be seen in Table 1.

In Table 1 it can be seen that the highest survival value was obtained in treatment D, namely feeding 100% dry *maggot* with a value of 91%,

while the smallest survival value was obtained in treatment A, namely feeding in the form of 100% commercial feed with a value of 78.5%. This is influenced by the cannibalism of catfish and the quality of the water. The death of catfish seeds in this study was caused by the cannibalism of catfish and water quality. Cannibalism is influenced by fish density, fish age, fish size and weight ratio of individual predators. Cannibalism can be reduced by the presence of other foods in the rearing medium. Cannibalism can be reduced by the presence of other foods in the rearing medium. Adequate nutrition from feed and the use of natural feed can increase the survival value of pearl catfish fry and reduce the cannibalism rate of pearl catfish fry. With the high survival rate and low cannibalism of pearl catfish seeds, the uniformity of seed size can be well maintained [15].

In addition to the cannibalism of catfish, the survival rate of catfish seeds can also be affected by water quality. That *maggot* has the ability to produce natural enzymes that stimulate fish appetite so that it can increase the digestibility of fish for feed and in the end there is no BSF *maggot* which is not eaten by fish and does not leave leftover feed which can affect quality water [16].

3.2 Specific Length Growth Rate

The value of the specific length growth rate in each treatment has a different value. This is because the nutritional components in the feed for each treatment are different. Data on the value of the specific elongation rate of mutiara catfish seeds can be seen in Table 2.

Table 1. Survival of Catfish Seeds

Treatment	Life sustainability (%)
A (100% commercial feed)	78.50a
B (75% commercial feed + 25% dry <i>maggot</i>)	79.50ab
C (50% commercial feed + 50% dry <i>maggot</i>)	84b
D (100% dry <i>maggot</i>)	91c

Table 2. Value of Growth Rate of Specific Length of Catfish Seeds

Treatment	LPBS(%)
D (100% dry <i>maggot</i>)	8.03a
A (100% commercial feed)	11,83b
B (75% commercial feed + 25% dry <i>maggot</i>)	13,36c
C (50% commercial feed + 50% dry <i>maggot</i>)	16,19d

Table 3. Water quality value during maintenance

Treatment	Observed observations			
	Temperature	pH	DO	ammonia
A	25-26	7	7.0-7.6	<0.03-0.1 mg/l
B	25-26	7	6,8-7,2	<0.03-0.1 mg/l
C	25-26	7	6.5-7.1	<0.03-0.1 mg/l
D	25-27	7	6.4-7.0	<0.03-0.05 mg/l

The highest growth rate for specific length of mutiara catfish seeds was found in treatment C by providing a combination of feed in the form of 50% commercial feed + 50% dry *maggot* with a value of 16.19% and the lowest treatment was treatment D by providing feed in the form of 100% dry *maggot* with a value 8.03%. All treatments had significant differences between treatments on Duncan's test. These differences are influenced by differences in the type and combination of feed given.

To produce optimal fish growth, a balance of feed composition components is needed [17]. Giving a combination of feed will affect the nutritional composition of the feed, because the more complete the nutritional composition of the feed given, the better the quality of growth, which is the difference in the combination of feed will affect the growth of each treatment. Each treatment was given a different feed combination, therefore each treatment had a specific length growth rate that differed from one another.

Treatment D with feeding in the form of 100% dry *maggot* had the lowest specific length growth rate. This is due to the limiting factor in *maggot*, namely chitin, which is difficult for fish to digest because it is not dissolved by strong acids, which interferes with fish growth [18].

3.3 Water Quality

The temperature and pH observations of each treatment showed an almost the same range, namely 25-27 °C and 7 respectively. States that catfish can live and grow well in the temperature range of 25-30 °C and pH 6.5-8. Temperature and pH values can affect the growth and survival of mutiara catfish seeds, because if the temperature and pH values are not appropriate, it will interfere with appetite and can result in death [19].

In observing the value of *dissolved oxygen* (DO) in each treatment showed relatively the same value, namely the range of 6.4-7.96. This value is

affected because the fish rearing container is given sufficient aeration to increase DO in each maintenance container. States that catfish can live and grow optimally if the DO value is > 3 mg/l [19].

The value of ammonia levels is one of the important factors in the growth and survival of catfish, if the value is too high from the optimal limit it will cause death. Value of ammonia levels in each treatment range <0.03-1 mg/l. This value is still in the optimal range for live catfish. Catfish are able to live in waters with ammonia levels <0.1 mg/l [19].

Feeding in the form of a combination of commercial feed and dry *maggot* does not affect the water quality in the rearing container. This is shown by the values of temperature, pH, DO and ammonia levels which are in accordance with SNI so that fish can maintain their survival and their growth can run optimally.

4. CONCLUSION

The results showed that the combination of commercial feed and dry *maggot* had an effect on the specific length growth rate and survival of pearl catfish seeds. the combination of 50% commercial feed and 50% dry *maggot* produced the highest specific length growth rate of 16.19%, 75% commercial feed and 25% dry *maggot* produced a specific length growth rate of 13.36%, with 100% commercial feed % produced in specific growth rate of 11.83% and feeding 100% dry *maggot* resulted in the lowest growth rate of 8.03%. The highest survival rate was in treatment D by feeding 100% dry *maggot* with a value of 91%, the survival rate in the treatment given a combination of 50% commercial feed and 50% dry *maggot* resulted in a value of 84%, with a value of 75% feeding commercial and *maggot* 25% dry resulted in a survival rate of 79.50% and the lowest survival rate at 100% commercial feeding was 78.50%. the water quality of the maintenance media is in accordance with the effective requirements for maintaining SNI

Mutiara catfish seeds. In the maintenance stage of size 3-5 Mutiara catfish seeds, a combination of commercial feed and dry maggot with a ratio of 50%: 50% should be given so that growth is more optimal.

ACKNOWLEDGEMENTS

The author would like to thank the Ministry of Maritime Affairs and Fisheries of Indramayu Regency, Cipancuh Fish Seed Center and to all those who have helped implementation of this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. BPPI. Fish breeding research bulletin. Subang 2014;3(2).
2. Matasina SZ, Tangguda S.. Growth of Mutiara Catfish (*Clarias gariepinus*) Seeds at Pt. Indosco Dwi Jaya (Farm Fish Booster Centre) Sidoarjo, East Java. Journal of Indonesian Swamp Aquaculture . 2020;8(2):123–128.
3. Mahyudin. Complete Guide to Catfish Agribusiness. Self-help Spreader. Jakarta; 2008.
4. Kordi, KMGH. Cultivating catfish in tarpaulin ponds. Andi. Yogyakarta. Thing. 2010:1-22
5. Sopha Soraya, Santoso Limin, Putri Berta. The Effect of Partial Substitution of Fish Meal with Bone Meal on the Growth of Sangkuriang Catfish (*Clarias gariepinus*). e-Journal of Aquaculture Engineering and Technology. 2015;III(2)
6. Sylvawan, Hasan H, Sunarto.. The Effectiveness of Noni (*Morinda citrifolia*) Fruit Extract to Reduce the Cannibalism Rate of Sangkuriang Catfish (*Clarias* sp) Seeds with the Bioencapsulation Method. Ruaya Journal. 2014;2:44-57.
7. Herlina S. Effect of Different Feed Types on Growth and Survival of Snakehead Fish (*Channa striata*) Seeds . Journal of Tropical Animal Science. 2016;5(2):64–67.
8. Rachmawati R, Buchori D, Hidayat P, Hem S, Fahmi MR. Development and Nutritional Content of *Hermetia illucens* (Linnaeus) Larvae (Diptera: Stratiomyidae) in Oil Palm Cake. Journal of Indonesian Entomology. 2015;7(1):28
9. Rini Fahmi M, Hem S, Wayan Subamia, Subamia I, Wayan. Potential of Maggot for Increasing Growth and Health Status of Fish. Aquaculture Research Journal . 2009;4(2):221–232.
10. Habibie MR, Yustiati A, Haetami K, Suryadi IB. Growth Performance of Pearl Catfish (*Clarias gariepinus*) with a Combination of Commercial Feed and Black Soldier Fly Maggot (*Hermetia illucens*). Asian Journal of Fisheries and Aquatic Research. 2020;9(4):27–36.
11. Irawan D, Helmizuryanu.. Analysis of Differences in the Type of Feed as a Substitute for Pellets on the Growth and Survival of Sangkuriang Catfish (*Clarias gariepinus*). Fiseries. 2014;3(1): 24 – 24.
12. Sepang Daniella A, Mudeng Jopy D, Monijung Revol D, Sambali Hariyani, Mokolensang Jeffrie F. Growth of tilapia (*Oreochromis niloticus*) fed a combination of pellets and dry maggot (*Hermetia illucens*) with different presentations. Sam Ratulangi University. Aquaculture. 2021;9(1): 33 – 44
13. Effendi MI. Fisheries Biology. Pustaka Nusantara Foundation, Yogyakarta; 2002.
14. Gaspersz V. Experiment Design Methods. CV. ARMICO; 1991.
15. Fessehaye Y, Kabir A, Bovenhuis H, Komen H. Prediction of cannibalism in juvenile *Oreochromis niloticus* based on predator to prey weight ratio and effects of age and stocking density. Aquaculture. 2005;20:9-17
16. Fauzi, Rizal UA, Sari Eeka RN. Analysis of Maggot Cultivation Business as an Alternative to Catfish Feed. Journal of Agro-industry Technology and Management. 2018;1:39-46.
17. Dani NP, Budiharjo A, Listyawati S.. Composition of Artificial Feed to Increase Growth and Protein Content of Tawes Fish (*Puntius javanicus* Blkr.). Journal of Bio Smart. 2005;7(2):82-90
18. Suarjuniarta I Kadek A, Juliantoro Pande GS, Kartika I Wayan D. Feed Conversion Ratio, Growth and Survival of Catfish (*Clarias* Sp.) Fed Commercial Pellets and

BSF Maggot Black Soldier Fly (*Hermetia illucens*). Current Trends in Aquatic Sciences. 2021;IV(2):152-158

19. Indonesian National Standard (SNI). Standards for Dumbo Catfish (*Clarias sp.*). SNI. 2014;6484.3:2014.

© 2023 Ashqolani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95808>